

Table 2.57C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	595	597	596	602	563	587
NOT HISPANIC OR LATINO¹	548	561	564	566	534	546
White	447	443	509	505	479	484
Black or African American	250	222	189	194	181	189
American Indian or Alaska Native	33	28	35	39	35	36
Native Hawaiian or Other Pacific Islander	29	28	35	26	27	24
Chinese	79	69	36	30	31	25
Filipino	75	78	56	45	47	39
Japanese	37	45	24	21	23	20
Asian Indian	69	62	52	38	50	34
Korean	41	37	33	24	32	22
Vietnamese	37	32	29	30	28	30
HISPANIC OR LATINO¹	206	217	193	179	185	176
Mexican	166	174	141	146	139	138
Puerto Rican	57	68	67	65	67	66
Central or South American	91	89	75	61	68	58
Cuban	45	40	47	40	39	37

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.57D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.25	0.25	0.25	0.25	0.24	0.25
NOT HISPANIC OR LATINO¹	0.26	0.27	0.27	0.27	0.26	0.26
White	0.27	0.27	0.31	0.30	0.29	0.29
Black or African American	0.92	0.80	0.69	0.70	0.67	0.68
American Indian or Alaska Native	2.38	2.17	2.56	3.08	2.54	2.82
Native Hawaiian or Other Pacific Islander	4.44	5.00	5.31	4.67	4.09	4.28
Chinese	3.75	2.99	1.71	1.30	1.46	1.07
Filipino	3.58	3.55	2.67	2.06	2.24	1.76
Japanese	4.43	5.25	2.89	2.45	2.67	2.34
Asian Indian	3.44	2.75	2.60	1.69	2.48	1.50
Korean	5.23	5.05	4.17	3.23	4.05	2.96
Vietnamese	4.47	4.51	3.51	4.22	3.41	4.15
HISPANIC OR LATINO¹	0.70	0.71	0.65	0.59	0.63	0.58
Mexican	0.85	0.87	0.72	0.73	0.71	0.69
Puerto Rican	1.80	2.20	2.12	2.09	2.10	2.14
Central or South American	2.26	2.02	1.86	1.39	1.69	1.31
Cuban	3.52	2.99	3.75	2.99	3.05	2.80

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.57P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0001		0.0098		0.0203	
NOT HISPANIC OR LATINO¹	0.0078		0.0240		0.0509	
White	0.0793		0.3255		0.4358	
Black or African American	0.1496		0.1419		0.1624	
American Indian or Alaska Native	0.5856		0.1889		0.3065	
Native Hawaiian or Other Pacific Islander	0.1594		0.9036		0.9518	
Chinese	0.9048		0.7333		0.6398	
Filipino	0.9904		0.0012		0.0122	
Japanese	0.6565		0.2320		0.2191	
Asian Indian	0.5068		0.0580		0.1216	
Korean	0.0154		0.0543		0.0518	
Vietnamese	0.9569		0.9569		0.8619	
HISPANIC OR LATINO¹	0.0018		0.2016		0.1847	
Mexican	0.0093		0.3365		0.1877	
Puerto Rican	0.2075		0.8027		0.6486	
Central or South American	0.0555		0.0394		0.1115	
Cuban	0.6173		0.7110		0.7427	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.58C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	80	74	64	63	54	53
NOT HISPANIC OR LATINO¹	73	66	59	57	50	49
White	64	57	53	52	45	45
Black or African American	26	27	19	19	15	14
American Indian or Alaska Native	6	8	6	6	5	6
Native Hawaiian or Other Pacific Islander	*	*	*	2	2	2
Chinese	5	6	4	4	3	3
Filipino	6	*	5	4	3	3
Japanese	*	*	*	*	*	*
Asian Indian	6	7	5	5	3	4
Korean	4	*	*	*	1	*
Vietnamese	5	*	2	3	1	*
HISPANIC OR LATINO¹	33	35	26	26	19	20
Mexican	30	31	22	23	17	18
Puerto Rican	9	11	8	9	8	6
Central or South American	10	11	8	7	6	5
Cuban	6	5	5	4	5	3

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.58D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.32	0.30	0.26	0.25	0.22	0.21
NOT HISPANIC OR LATINO¹	0.35	0.32	0.28	0.27	0.24	0.23
White	0.41	0.37	0.34	0.33	0.29	0.29
Black or African American	0.72	0.71	0.52	0.51	0.40	0.38
American Indian or Alaska Native	3.43	4.40	3.19	3.54	2.82	3.35
Native Hawaiian or Other Pacific Islander	*	*	*	3.38	2.85	3.08
Chinese	2.33	2.70	1.94	2.16	1.30	1.53
Filipino	3.74	*	2.93	2.47	2.04	2.03
Japanese	*	*	*	*	*	*
Asian Indian	2.99	3.06	2.58	2.00	1.66	1.78
Korean	4.72	*	*	*	0.95	*
Vietnamese	4.28	*	2.01	2.73	1.00	*
HISPANIC OR LATINO¹	0.83	0.84	0.64	0.63	0.48	0.48
Mexican	1.04	1.07	0.78	0.81	0.58	0.61
Puerto Rican	1.98	2.28	1.77	1.81	1.59	1.26
Central or South American	2.41	2.40	1.93	1.57	1.33	1.20
Cuban	4.76	3.56	4.19	2.95	3.95	2.11

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.58P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0000		0.0002		0.0182	
NOT HISPANIC OR LATINO¹	0.0000		0.0005		0.0300	
White	0.0000		0.0046		0.0358	
Black or African American	0.0304		0.1532		0.4426	
American Indian or Alaska Native	0.1395		0.2725		0.1747	
Native Hawaiian or Other Pacific Islander	*		*		0.6409	
Chinese	0.4692		0.9091		0.4998	
Filipino	*		0.3683		0.8355	
Japanese	*		*		*	
Asian Indian	0.2019		0.0718		0.4252	
Korean	*		*		*	
Vietnamese	*		0.6976		*	
HISPANIC OR LATINO¹	0.0120		0.2282		0.3122	
Mexican	0.1023		0.4966		0.5771	
Puerto Rican	0.0350		0.2903		0.3408	
Central or South American	0.2120		0.0909		0.2559	
Cuban	0.0093		0.2363		0.1074	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.59C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	106	104	110	108	107	108
NOT HISPANIC OR LATINO¹	92	88	92	93	91	96
White	71	72	77	84	78	85
Black or African American	38	36	36	34	33	35
American Indian or Alaska Native	5	5	7	6	7	6
Native Hawaiian or Other Pacific Islander	6	7	8	7	8	7
Chinese	12	15	9	13	9	11
Filipino	10	11	11	10	11	9
Japanese	*	*	*	*	*	*
Asian Indian	10	10	10	10	9	8
Korean	10	9	9	10	9	10
Vietnamese	8	9	8	7	7	7
HISPANIC OR LATINO¹	48	51	51	52	48	50
Mexican	41	44	45	43	41	40
Puerto Rican	14	16	16	15	15	15
Central or South American	17	19	15	20	15	18
Cuban	7	6	7	7	7	7

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.59D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.34	0.33	0.35	0.34	0.34	0.34
NOT HISPANIC OR LATINO¹	0.36	0.33	0.36	0.35	0.35	0.36
White	0.36	0.36	0.39	0.43	0.40	0.43
Black or African American	0.92	0.83	0.86	0.80	0.79	0.80
American Indian or Alaska Native	2.54	2.98	3.49	3.51	3.60	3.64
Native Hawaiian or Other Pacific Islander	3.93	4.69	5.17	5.32	4.89	4.68
Chinese	4.22	4.17	3.25	3.66	3.07	3.06
Filipino	3.74	4.06	3.86	3.94	3.95	3.57
Japanese	*	*	*	*	*	*
Asian Indian	3.38	3.26	3.47	2.98	3.06	2.53
Korean	5.20	5.20	4.96	5.72	5.08	5.72
Vietnamese	5.39	5.51	5.11	4.39	4.31	3.94
HISPANIC OR LATINO¹	0.85	0.90	0.91	0.91	0.85	0.88
Mexican	1.05	1.09	1.13	1.07	1.04	0.99
Puerto Rican	2.50	3.01	2.90	2.97	2.66	2.93
Central or South American	2.29	2.55	2.04	2.65	2.06	2.44
Cuban	4.58	4.20	4.94	4.71	4.75	4.94

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.59P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0001		0.0322		0.0746	
NOT HISPANIC OR LATINO¹	0.0001		0.0064		0.0165	
White	0.0005		0.0051		0.0184	
Black or African American	0.2830		0.5573		0.4805	
American Indian or Alaska Native	0.8701		0.6479		0.6568	
Native Hawaiian or Other Pacific Islander	0.6562		0.7646		0.5330	
Chinese	0.4213		0.2636		0.7594	
Filipino	0.4670		0.3335		0.5727	
Japanese	*		*		*	
Asian Indian	0.4015		0.5485		0.8744	
Korean	0.1032		0.1750		0.1571	
Vietnamese	0.5287		0.6604		0.7003	
HISPANIC OR LATINO¹	0.3916		0.6138		0.5049	
Mexican	0.7697		0.3341		0.5249	
Puerto Rican	0.5447		0.6395		0.8621	
Central or South American	0.2769		0.2015		0.4662	
Cuban	0.9371		0.7439		0.4521	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.60C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	570	556	549	561	521	548
NOT HISPANIC OR LATINO¹	522	526	521	532	496	515
White	429	417	461	470	438	454
Black or African American	240	205	182	189	176	182
American Indian or Alaska Native	30	24	33	37	33	33
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Chinese	76	66	32	24	27	19
Filipino	73	77	53	43	45	37
Japanese	36	43	24	20	22	19
Asian Indian	67	60	50	34	48	31
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	201	210	181	165	172	161
Mexican	163	167	130	131	126	126
Puerto Rican	55	63	66	63	65	64
Central or South American	88	85	72	59	67	57
Cuban	47	40	50	40	41	37

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.60D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.32	0.31	0.30	0.31	0.29	0.30
NOT HISPANIC OR LATINO¹	0.33	0.33	0.33	0.33	0.31	0.32
White	0.33	0.32	0.35	0.36	0.34	0.35
Black or African American	1.25	1.04	0.95	0.96	0.91	0.93
American Indian or Alaska Native	3.01	2.55	3.39	3.94	3.39	3.57
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Chinese	4.75	3.80	1.99	1.36	1.70	1.07
Filipino	4.41	4.37	3.19	2.44	2.69	2.10
Japanese	4.95	5.66	3.27	2.57	3.05	2.47
Asian Indian	4.46	3.53	3.31	2.01	3.20	1.83
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	1.01	1.02	0.91	0.80	0.87	0.78
Mexican	1.29	1.28	1.02	1.01	1.00	0.96
Puerto Rican	2.56	3.01	3.05	3.01	3.02	3.05
Central or South American	3.05	2.64	2.49	1.85	2.31	1.77
Cuban	4.66	3.82	5.02	3.75	4.10	3.50

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.60P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0179		0.0504		0.0558	
NOT HISPANIC OR LATINO¹	0.1814		0.1117		0.1366	
White	0.5485		0.7140		0.7575	
Black or African American	0.3437		0.1393		0.1325	
American Indian or Alaska Native	0.7652		0.2964		0.3857	
Native Hawaiian or Other Pacific Islander	*		*		*	
Chinese	0.7271		0.2743		0.3598	
Filipino	0.8668		0.0018		0.0131	
Japanese	0.8005		0.1902		0.1578	
Asian Indian	0.6767		0.0961		0.1345	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0086		0.2026		0.1497	
Mexican	0.0175		0.2349		0.1316	
Puerto Rican	0.5050		0.9259		0.7772	
Central or South American	0.0821		0.1077		0.1790	
Cuban	0.9286		0.8179		0.8500	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.61C Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	585	578	583	592	554	576
NOT HISPANIC OR LATINO¹	536	540	552	559	524	537
White	440	429	498	499	469	479
Black or African American	247	215	187	191	180	185
American Indian or Alaska Native	30	25	34	39	34	36
Native Hawaiian or Other Pacific Islander	27	28	33	26	27	24
Chinese	79	68	36	29	31	24
Filipino	76	77	58	45	48	39
Japanese	37	44	24	21	23	20
Asian Indian	68	62	51	38	49	33
Korean	*	36	33	25	32	22
Vietnamese	37	31	29	31	28	*
HISPANIC OR LATINO¹	210	220	193	180	183	175
Mexican	169	177	142	144	137	136
Puerto Rican	55	66	66	65	66	67
Central or South American	91	89	74	61	68	58
Cuban	45	39	48	40	39	38

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.61D Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.28	0.27	0.28	0.28	0.26	0.27
NOT HISPANIC OR LATINO¹	0.29	0.29	0.30	0.30	0.28	0.29
White	0.29	0.28	0.33	0.33	0.31	0.32
Black or African American	1.05	0.90	0.80	0.80	0.77	0.77
American Indian or Alaska Native	2.52	2.27	2.90	3.52	2.85	3.23
Native Hawaiian or Other Pacific Islander	4.66	5.71	5.73	5.29	4.70	4.83
Chinese	4.14	3.23	1.88	1.38	1.61	1.17
Filipino	3.93	3.79	2.98	2.23	2.50	1.91
Japanese	4.64	5.37	3.08	2.56	2.83	2.44
Asian Indian	3.79	3.05	2.84	1.86	2.73	1.65
Korean	*	5.50	4.70	3.75	4.54	3.43
Vietnamese	5.29	5.08	4.08	5.03	3.90	*
HISPANIC OR LATINO¹	0.82	0.84	0.76	0.68	0.72	0.67
Mexican	1.01	1.04	0.85	0.84	0.82	0.80
Puerto Rican	2.05	2.52	2.45	2.51	2.43	2.55
Central or South American	2.51	2.24	2.06	1.55	1.89	1.47
Cuban	3.95	3.30	4.20	3.38	3.41	3.15

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.61P Cigarette Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0027		0.0294		0.0368	
NOT HISPANIC OR LATINO¹	0.0472		0.0571		0.0769	
White	0.2471		0.4772		0.5433	
Black or African American	0.2560		0.1850		0.1831	
American Indian or Alaska Native	0.7824		0.2689		0.4360	
Native Hawaiian or Other Pacific Islander	0.1403		0.9744		0.9230	
Chinese	0.8154		0.7044		0.5763	
Filipino	0.8372		0.0018		0.0128	
Japanese	0.7560		0.2110		0.2157	
Asian Indian	0.5973		0.0840		0.1343	
Korean	*		0.0439		0.0243	
Vietnamese	0.8775		0.9530		*	
HISPANIC OR LATINO¹	0.0070		0.2741		0.2231	
Mexican	0.0188		0.3838		0.1969	
Puerto Rican	0.4233		0.9926		0.7953	
Central or South American	0.0614		0.0551		0.1208	
Cuban	0.9308		0.8189		0.9008	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.62C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	457	466	655	644	688	672
NOT HISPANIC OR LATINO¹	416	430	613	620	651	639
White	318	332	526	548	578	570
Black or African American	168	190	214	206	195	220
American Indian or Alaska Native	40	29	42	48	48	37
Native Hawaiian or Other Pacific Islander	25	30	28	29	28	25
Chinese	58	72	74	78	69	68
Filipino	66	79	83	110	75	97
Japanese	34	30	44	44	45	47
Asian Indian	63	66	66	65	60	62
Korean	38	39	40	39	40	38
Vietnamese	45	38	42	37	36	32
HISPANIC OR LATINO¹	176	189	240	230	233	212
Mexican	139	149	187	188	179	164
Puerto Rican	57	58	77	71	73	67
Central or South American	70	79	88	93	84	81
Cuban	28	36	37	48	43	44

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.62D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.19	0.20	0.28	0.27	0.29	0.28
NOT HISPANIC OR LATINO¹	0.20	0.21	0.30	0.30	0.31	0.31
White	0.19	0.20	0.32	0.33	0.35	0.34
Black or African American	0.62	0.68	0.79	0.74	0.72	0.79
American Indian or Alaska Native	2.93	2.31	3.05	3.74	3.53	2.93
Native Hawaiian or Other Pacific Islander	3.85	5.40	4.37	5.15	4.26	4.50
Chinese	2.76	3.11	3.49	3.39	3.28	2.97
Filipino	3.13	3.59	3.96	5.01	3.60	4.40
Japanese	4.07	3.45	5.23	5.07	5.35	5.44
Asian Indian	3.15	2.93	3.31	2.86	3.02	2.73
Korean	4.82	5.32	5.08	5.32	5.04	5.09
Vietnamese	5.44	5.23	5.17	5.12	4.39	4.40
HISPANIC OR LATINO¹	0.60	0.62	0.82	0.76	0.79	0.70
Mexican	0.71	0.75	0.96	0.94	0.92	0.82
Puerto Rican	1.79	1.87	2.42	2.30	2.31	2.17
Central or South American	1.74	1.81	2.17	2.11	2.06	1.84
Cuban	2.22	2.67	2.94	3.60	3.36	3.29

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.62P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0755		0.0744		0.2649	
NOT HISPANIC OR LATINO¹	0.3841		0.3627		0.6079	
White	0.8182		0.9956		0.7777	
Black or African American	0.1057		0.0791		0.0630	
American Indian or Alaska Native	0.8781		0.2055		0.2258	
Native Hawaiian or Other Pacific Islander	0.2028		0.6545		0.7910	
Chinese	0.7665		0.8269		0.4694	
Filipino	0.4241		0.8926		0.3238	
Japanese	0.4987		0.1781		0.0820	
Asian Indian	0.7521		0.4160		0.6618	
Korean	0.6322		0.3582		0.4473	
Vietnamese	0.5200		0.8251		0.8197	
HISPANIC OR LATINO¹	0.0468		0.0197		0.0720	
Mexican	0.2321		0.0346		0.0560	
Puerto Rican	0.0845		0.7825		0.9472	
Central or South American	0.5124		0.3739		0.2878	
Cuban	0.1900		0.8784		0.6405	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.63C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	80	76	76	71	60	55
NOT HISPANIC OR LATINO¹	72	68	69	64	54	50
White	62	61	57	58	47	44
Black or African American	29	31	26	26	19	18
American Indian or Alaska Native	7	7	6	5	5	4
Native Hawaiian or Other Pacific Islander	5	4	5	4	3	*
Chinese	8	7	7	7	4	6
Filipino	5	7	5	5	5	5
Japanese	*	*	*	*	2	*
Asian Indian	7	9	7	9	5	6
Korean	*	*	*	*	3	3
Vietnamese	6	*	*	*	3	*
HISPANIC OR LATINO¹	36	38	33	35	25	29
Mexican	33	33	29	29	21	24
Puerto Rican	11	12	11	10	8	9
Central or South American	11	11	10	11	9	10
Cuban	5	6	5	6	5	4

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.63D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.32	0.30	0.30	0.28	0.24	0.22
NOT HISPANIC OR LATINO¹	0.35	0.33	0.33	0.31	0.26	0.24
White	0.40	0.39	0.37	0.37	0.30	0.28
Black or African American	0.80	0.82	0.69	0.68	0.53	0.48
American Indian or Alaska Native	3.85	3.84	3.17	3.02	2.73	2.17
Native Hawaiian or Other Pacific Islander	5.80	5.71	5.85	5.02	4.09	*
Chinese	3.72	3.58	3.35	3.24	2.07	2.67
Filipino	3.08	4.17	3.08	3.18	2.90	2.84
Japanese	*	*	*	*	3.19	*
Asian Indian	3.30	3.92	3.15	3.89	2.22	2.48
Korean	*	*	*	*	3.27	3.61
Vietnamese	5.48	*	*	*	2.69	*
HISPANIC OR LATINO¹	0.91	0.91	0.83	0.84	0.64	0.69
Mexican	1.16	1.13	1.02	1.01	0.73	0.81
Puerto Rican	2.26	2.40	2.25	2.13	1.69	1.89
Central or South American	2.52	2.39	2.38	2.38	2.13	2.23
Cuban	4.45	4.56	4.57	4.48	4.38	3.11

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.63P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0236		0.2125		0.9274	
NOT HISPANIC OR LATINO¹	0.0445		0.2333		0.5136	
White	0.1023		0.3556		0.7334	
Black or African American	0.1834		0.4082		0.2204	
American Indian or Alaska Native	0.1943		0.2971		0.4271	
Native Hawaiian or Other Pacific Islander	0.7288		0.1887		*	
Chinese	0.0723		0.2555		0.1202	
Filipino	0.0144		0.2155		0.4038	
Japanese	*		*		*	
Asian Indian	0.8204		0.7875		0.7916	
Korean	*		*		0.7882	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.2476		0.6647		0.2903	
Mexican	0.5332		0.8448		0.2582	
Puerto Rican	0.1226		0.1198		0.2551	
Central or South American	0.4972		0.7889		0.8067	
Cuban	0.0814		0.2481		0.3225	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.64C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	77	74	99	92	124	116
NOT HISPANIC OR LATINO¹	68	65	84	78	109	101
White	53	50	65	62	87	83
Black or African American	28	30	33	35	36	39
American Indian or Alaska Native	4	5	7	6	7	6
Native Hawaiian or Other Pacific Islander	7	5	8	6	8	7
Chinese	9	10	9	11	12	14
Filipino	7	8	12	12	10	9
Japanese	*	*	*	*	*	*
Asian Indian	11	11	11	11	11	11
Korean	8	7	8	8	10	9
Vietnamese	7	9	8	9	*	9
HISPANIC OR LATINO¹	35	38	45	46	47	48
Mexican	31	34	39	38	40	42
Puerto Rican	9	12	11	12	14	15
Central or South American	13	13	15	18	17	17
Cuban	3	3	4	4	5	6

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.64D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.25	0.23	0.31	0.29	0.40	0.36
NOT HISPANIC OR LATINO¹	0.26	0.25	0.32	0.30	0.42	0.39
White	0.27	0.25	0.33	0.31	0.45	0.42
Black or African American	0.68	0.70	0.80	0.82	0.85	0.90
American Indian or Alaska Native	2.10	2.83	3.66	3.42	3.51	3.49
Native Hawaiian or Other Pacific Islander	4.20	3.78	4.73	4.49	4.87	4.72
Chinese	3.28	2.77	3.12	3.06	4.11	4.07
Filipino	2.59	3.06	4.34	4.79	3.80	3.64
Japanese	*	*	*	*	*	*
Asian Indian	3.80	3.42	3.90	3.37	3.89	3.52
Korean	4.36	4.07	4.55	4.53	5.61	5.58
Vietnamese	4.20	5.33	5.40	5.47	*	5.64
HISPANIC OR LATINO¹	0.63	0.67	0.81	0.80	0.84	0.85
Mexican	0.78	0.85	0.99	0.95	1.02	1.03
Puerto Rican	1.60	2.23	1.94	2.23	2.50	2.86
Central or South American	1.85	1.78	2.10	2.38	2.28	2.27
Cuban	2.06	2.47	2.57	2.58	3.49	4.20

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.64P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.3292		0.7494		0.9921	
NOT HISPANIC OR LATINO¹	0.4804		0.2949		0.6358	
White	0.6186		0.2152		0.3003	
Black or African American	0.0367		0.8752		0.9046	
American Indian or Alaska Native	0.8814		0.6534		0.5355	
Native Hawaiian or Other Pacific Islander	0.9321		0.8993		0.3448	
Chinese	0.2007		0.2203		0.7405	
Filipino	0.4380		0.7116		0.6021	
Japanese	*		*		*	
Asian Indian	0.9881		1.0000		0.5223	
Korean	0.5322		0.8521		0.4225	
Vietnamese	0.0823		0.7337		*	
HISPANIC OR LATINO¹	0.4822		0.2424		0.3107	
Mexican	0.6470		0.4848		0.3609	
Puerto Rican	0.1244		0.4129		0.5396	
Central or South American	0.1938		0.8910		0.7294	
Cuban	0.3761		0.6734		0.9289	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.65C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	434	438	636	630	661	659
NOT HISPANIC OR LATINO¹	393	409	598	611	627	625
White	298	318	518	543	561	563
Black or African American	159	174	207	197	185	207
American Indian or Alaska Native	35	26	37	48	46	37
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Chinese	53	69	70	74	66	68
Filipino	64	79	81	106	77	96
Japanese	*	28	*	42	*	*
Asian Indian	59	63	62	62	58	60
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	175	180	234	223	228	207
Mexican	143	139	183	182	172	161
Puerto Rican	54	59	73	72	72	67
Central or South American	69	76	85	91	81	78
Cuban	28	35	38	48	42	43

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.65D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.24	0.24	0.35	0.35	0.37	0.36
NOT HISPANIC OR LATINO¹	0.25	0.25	0.37	0.38	0.39	0.39
White	0.23	0.24	0.40	0.41	0.43	0.43
Black or African American	0.83	0.89	1.07	1.00	0.96	1.06
American Indian or Alaska Native	3.57	2.80	3.80	5.12	4.64	4.02
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Chinese	3.33	3.97	4.38	4.26	4.12	3.93
Filipino	3.89	4.47	4.85	5.98	4.66	5.42
Japanese	*	3.63	*	5.56	*	*
Asian Indian	3.91	3.72	4.11	3.64	3.83	3.51
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	0.88	0.87	1.17	1.08	1.15	1.00
Mexican	1.12	1.07	1.44	1.39	1.35	1.23
Puerto Rican	2.53	2.81	3.38	3.42	3.34	3.20
Central or South American	2.38	2.36	2.94	2.84	2.81	2.44
Cuban	2.76	3.32	3.80	4.57	4.20	4.04

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.65P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.1691		0.0700		0.2310	
NOT HISPANIC OR LATINO¹	0.5610		0.2940		0.5207	
White	0.8698		0.8562		0.9485	
Black or African American	0.2769		0.0840		0.0641	
American Indian or Alaska Native	0.6508		0.2160		0.3066	
Native Hawaiian or Other Pacific Islander	*		*		*	
Chinese	0.3563		0.5201		0.3174	
Filipino	0.3312		0.9402		0.2598	
Japanese	*		*		*	
Asian Indian	0.7212		0.3834		0.7215	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0966		0.0389		0.0808	
Mexican	0.2965		0.0461		0.0516	
Puerto Rican	0.2881		0.8003		0.7103	
Central or South American	0.3443		0.3516		0.2602	
Cuban	0.4533		0.6478		0.4574	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.66C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	442	445	651	640	679	667
NOT HISPANIC OR LATINO¹	404	418	612	622	643	637
White	307	321	525	548	569	569
Black or African American	164	179	214	204	190	216
American Indian or Alaska Native	36	25	40	49	47	38
Native Hawaiian or Other Pacific Islander	26	*	29	29	28	25
Chinese	56	70	73	75	69	67
Filipino	64	79	83	110	75	96
Japanese	*	30	45	43	46	46
Asian Indian	61	66	64	64	59	62
Korean	*	*	*	*	39	36
Vietnamese	*	*	*	35	38	31
HISPANIC OR LATINO¹	174	182	238	229	231	211
Mexican	141	143	186	189	174	165
Puerto Rican	56	58	75	71	74	68
Central or South American	69	77	86	91	84	80
Cuban	28	35	39	48	43	43

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.66D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.21	0.21	0.31	0.30	0.32	0.31
NOT HISPANIC OR LATINO¹	0.22	0.22	0.33	0.33	0.35	0.34
White	0.20	0.21	0.35	0.36	0.38	0.38
Black or African American	0.70	0.75	0.91	0.85	0.81	0.90
American Indian or Alaska Native	3.07	2.30	3.36	4.45	3.95	3.46
Native Hawaiian or Other Pacific Islander	4.45	*	5.01	5.90	4.84	5.16
Chinese	2.95	3.33	3.82	3.58	3.62	3.20
Filipino	3.31	3.87	4.30	5.39	3.90	4.72
Japanese	*	3.59	5.62	5.26	5.73	5.60
Asian Indian	3.39	3.28	3.58	3.18	3.28	3.08
Korean	*	*	*	*	5.62	5.54
Vietnamese	*	*	*	5.73	5.31	5.01
HISPANIC OR LATINO¹	0.68	0.69	0.93	0.87	0.91	0.80
Mexican	0.85	0.84	1.12	1.11	1.05	0.97
Puerto Rican	2.07	2.23	2.78	2.72	2.74	2.60
Central or South American	1.91	1.95	2.39	2.31	2.33	2.03
Cuban	2.45	2.97	3.38	4.03	3.79	3.63

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.66P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.1295		0.0911		0.2513	
NOT HISPANIC OR LATINO¹	0.4954		0.4001		0.6032	
White	0.9309		0.9744		0.8016	
Black or African American	0.1591		0.0981		0.0808	
American Indian or Alaska Native	0.6840		0.2535		0.2630	
Native Hawaiian or Other Pacific Islander	*		0.5406		0.8368	
Chinese	0.5016		0.6630		0.3601	
Filipino	0.3397		0.9278		0.3119	
Japanese	*		0.2218		0.1139	
Asian Indian	0.7226		0.3927		0.6452	
Korean	*		*		0.3985	
Vietnamese	*		*		0.8342	
HISPANIC OR LATINO¹	0.0684		0.0235		0.0504	
Mexican	0.2408		0.0341		0.0306	
Puerto Rican	0.1879		0.9463		0.8511	
Central or South American	0.4688		0.3284		0.2315	
Cuban	0.3916		0.7203		0.4989	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.67C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	688	672	512	483	286	274
NOT HISPANIC OR LATINO¹	651	639	493	461	265	256
White	578	570	430	414	239	241
Black or African American	195	220	161	164	89	85
American Indian or Alaska Native	48	37	31	33	20	21
Native Hawaiian or Other Pacific Islander	28	25	20	22	10	9
Chinese	69	68	37	38	9	20
Filipino	75	97	40	44	14	18
Japanese	45	47	22	22	5	5
Asian Indian	60	62	36	35	14	8
Korean	40	38	28	25	22	12
Vietnamese	36	32	19	12	11	7
HISPANIC OR LATINO¹	233	212	184	171	94	91
Mexican	179	164	153	143	74	80
Puerto Rican	73	67	60	58	36	34
Central or South American	84	81	65	64	24	20
Cuban	43	44	42	40	35	14

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.67D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.29	0.28	0.22	0.20	0.12	0.11
NOT HISPANIC OR LATINO¹	0.31	0.31	0.24	0.22	0.13	0.12
White	0.35	0.34	0.26	0.25	0.14	0.14
Black or African American	0.72	0.79	0.59	0.59	0.33	0.30
American Indian or Alaska Native	3.53	2.93	2.30	2.59	1.44	1.64
Native Hawaiian or Other Pacific Islander	4.26	4.50	3.01	3.98	1.53	1.65
Chinese	3.28	2.97	1.74	1.66	0.43	0.86
Filipino	3.60	4.40	1.89	1.99	0.67	0.82
Japanese	5.35	5.44	2.63	2.59	0.57	0.63
Asian Indian	3.02	2.73	1.82	1.55	0.72	0.34
Korean	5.04	5.09	3.47	3.37	2.74	1.68
Vietnamese	4.39	4.40	2.29	1.72	1.38	0.93
HISPANIC OR LATINO¹	0.79	0.70	0.63	0.56	0.32	0.30
Mexican	0.92	0.82	0.79	0.72	0.38	0.40
Puerto Rican	2.31	2.17	1.90	1.88	1.14	1.09
Central or South American	2.06	1.84	1.59	1.47	0.58	0.45
Cuban	3.36	3.29	3.31	2.97	2.80	1.08

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.67P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.2649		0.7770		0.3893	
NOT HISPANIC OR LATINO¹	0.6079		0.9413		0.1789	
White	0.7777		0.4184		0.1039	
Black or African American	0.0630		0.0269		0.9419	
American Indian or Alaska Native	0.2258		0.6876		0.7932	
Native Hawaiian or Other Pacific Islander	0.7910		0.5213		0.2485	
Chinese	0.4694		0.2762		0.1181	
Filipino	0.3238		0.9840		0.3933	
Japanese	0.0820		0.8472		0.6363	
Asian Indian	0.6618		0.7916		0.5214	
Korean	0.4473		0.8027		0.4667	
Vietnamese	0.8197		0.8152		0.7034	
HISPANIC OR LATINO¹	0.0720		0.5389		0.3452	
Mexican	0.0560		0.2181		0.4760	
Puerto Rican	0.9472		0.2359		0.9966	
Central or South American	0.2878		0.5234		0.5087	
Cuban	0.6405		0.9688		0.5560	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.68C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	60	55	49	50	23	25
NOT HISPANIC OR LATINO¹	54	50	44	45	21	22
White	47	44	40	42	20	21
Black or African American	19	18	14	13	4	4
American Indian or Alaska Native	5	4	5	4	2	2
Native Hawaiian or Other Pacific Islander	3	*	2	*	*	*
Chinese	4	6	3	4	*	*
Filipino	5	5	3	3	*	*
Japanese	2	*	0	*	0	*
Asian Indian	5	6	3	4	*	*
Korean	3	3	2	2	*	*
Vietnamese	3	*	*	*	*	*
HISPANIC OR LATINO¹	25	29	19	25	8	10
Mexican	21	24	16	20	8	9
Puerto Rican	8	9	6	6	1	3
Central or South American	9	10	7	9	3	3
Cuban	5	4	*	3	1	1

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.68D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.24	0.22	0.20	0.20	0.09	0.10
NOT HISPANIC OR LATINO¹	0.26	0.24	0.21	0.21	0.10	0.11
White	0.30	0.28	0.26	0.27	0.13	0.13
Black or African American	0.53	0.48	0.38	0.35	0.11	0.10
American Indian or Alaska Native	2.73	2.17	2.69	2.07	0.84	1.09
Native Hawaiian or Other Pacific Islander	4.09	*	3.12	*	*	*
Chinese	2.07	2.67	1.19	1.99	*	*
Filipino	2.90	2.84	1.88	1.81	*	*
Japanese	3.19	*	0.93	*	0.64	*
Asian Indian	2.22	2.48	1.34	1.91	*	*
Korean	3.27	3.61	2.46	1.89	*	*
Vietnamese	2.69	*	*	*	*	*
HISPANIC OR LATINO¹	0.64	0.69	0.48	0.60	0.20	0.25
Mexican	0.73	0.81	0.55	0.71	0.29	0.30
Puerto Rican	1.69	1.89	1.19	1.26	0.27	0.55
Central or South American	2.13	2.23	1.66	2.10	0.63	0.72
Cuban	4.38	3.11	*	1.92	0.97	0.43

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.68P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.9274		0.3035		0.5422	
NOT HISPANIC OR LATINO¹	0.5136		0.7384		0.8460	
White	0.7334		0.8455		0.8735	
Black or African American	0.2204		0.9637		0.8781	
American Indian or Alaska Native	0.4271		0.4980		0.5808	
Native Hawaiian or Other Pacific Islander	*		*		*	
Chinese	0.1202		0.2228		*	
Filipino	0.4038		0.3804		*	
Japanese	*		*		*	
Asian Indian	0.7916		0.5321		*	
Korean	0.7882		0.6659		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.2903		0.1068		0.2992	
Mexican	0.2582		0.1451		0.5822	
Puerto Rican	0.2551		0.1490		0.2512	
Central or South American	0.8067		0.2586		0.9883	
Cuban	0.3225		*		0.1355	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.69C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	124	116	119	117	84	86
NOT HISPANIC OR LATINO¹	109	101	105	104	75	76
White	87	83	86	88	66	68
Black or African American	36	39	31	31	16	18
American Indian or Alaska Native	7	6	7	6	4	4
Native Hawaiian or Other Pacific Islander	8	7	8	6	6	4
Chinese	12	14	9	11	4	5
Filipino	10	9	9	9	7	7
Japanese	*	*	*	*	3	2
Asian Indian	11	11	8	7	5	4
Korean	10	9	10	8	*	6
Vietnamese	*	9	6	7	4	4
HISPANIC OR LATINO¹	47	48	48	47	29	31
Mexican	40	42	40	41	23	26
Puerto Rican	14	15	14	13	9	9
Central or South American	17	17	15	17	9	10
Cuban	5	6	6	7	4	5

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.69D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.40	0.36	0.38	0.36	0.27	0.27
NOT HISPANIC OR LATINO¹	0.42	0.39	0.41	0.40	0.29	0.29
White	0.45	0.42	0.44	0.44	0.34	0.34
Black or African American	0.85	0.90	0.74	0.71	0.39	0.42
American Indian or Alaska Native	3.51	3.49	3.39	3.38	1.86	1.98
Native Hawaiian or Other Pacific Islander	4.87	4.72	4.83	4.54	3.89	2.99
Chinese	4.11	4.07	3.09	3.20	1.25	1.51
Filipino	3.80	3.64	3.42	3.56	2.46	2.53
Japanese	*	*	*	*	3.71	3.61
Asian Indian	3.89	3.52	2.90	2.22	1.59	1.36
Korean	5.61	5.58	5.15	4.96	*	3.34
Vietnamese	*	5.64	3.87	4.22	2.52	2.63
HISPANIC OR LATINO¹	0.84	0.85	0.86	0.83	0.52	0.55
Mexican	1.02	1.03	1.02	1.02	0.59	0.65
Puerto Rican	2.50	2.86	2.63	2.60	1.58	1.74
Central or South American	2.28	2.27	2.10	2.23	1.20	1.28
Cuban	3.49	4.20	4.36	4.94	2.71	3.84

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.69P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.9921		0.6945		0.7692	
NOT HISPANIC OR LATINO¹	0.6358		0.6898		0.7109	
White	0.3003		0.2889		0.5208	
Black or African American	0.9046		0.7538		0.9927	
American Indian or Alaska Native	0.5355		0.8007		0.5744	
Native Hawaiian or Other Pacific Islander	0.3448		0.2106		0.6979	
Chinese	0.7405		0.4210		0.1908	
Filipino	0.6021		0.6489		0.8775	
Japanese	*		*		0.8855	
Asian Indian	0.5223		0.6521		0.5702	
Korean	0.4225		0.5083		*	
Vietnamese	*		0.7731		0.9789	
HISPANIC OR LATINO¹	0.3107		0.9260		0.9258	
Mexican	0.3609		0.8757		0.6925	
Puerto Rican	0.5396		0.5002		0.6363	
Central or South American	0.7294		0.3728		0.8756	
Cuban	0.9289		0.1787		0.5495	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.70C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	661	659	478	463	257	252
NOT HISPANIC OR LATINO¹	627	625	459	435	235	236
White	561	563	402	394	213	221
Black or African American	185	207	154	159	86	83
American Indian or Alaska Native	46	37	29	32	19	21
Native Hawaiian or Other Pacific Islander	*	*	17	*	8	8
Chinese	66	68	33	33	*	19
Filipino	77	96	39	43	13	17
Japanese	*	*	20	21	3	4
Asian Indian	58	60	37	34	13	6
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	228	207	177	163	90	83
Mexican	172	161	144	135	72	72
Puerto Rican	72	67	58	58	35	32
Central or South American	81	78	62	59	22	17
Cuban	42	43	43	39	*	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.70D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.37	0.36	0.27	0.25	0.14	0.14
NOT HISPANIC OR LATINO¹	0.39	0.39	0.29	0.27	0.15	0.15
White	0.43	0.43	0.31	0.30	0.16	0.17
Black or African American	0.96	1.06	0.80	0.81	0.45	0.42
American Indian or Alaska Native	4.64	4.02	2.91	3.46	1.91	2.23
Native Hawaiian or Other Pacific Islander	*	*	4.27	*	1.88	2.19
Chinese	4.12	3.93	2.05	1.90	*	1.08
Filipino	4.66	5.42	2.33	2.40	0.78	0.94
Japanese	*	*	2.83	2.74	0.48	0.57
Asian Indian	3.83	3.51	2.45	2.01	0.89	0.34
Korean	*	*	*	*	*	*
Vietnamese	*	*	*	*	*	*
HISPANIC OR LATINO¹	1.15	1.00	0.89	0.79	0.45	0.40
Mexican	1.35	1.23	1.13	1.04	0.57	0.55
Puerto Rican	3.34	3.20	2.70	2.78	1.63	1.53
Central or South American	2.81	2.44	2.15	1.83	0.76	0.52
Cuban	4.20	4.04	4.33	3.73	*	1.43

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.70P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.2310		0.5427		0.4932	
NOT HISPANIC OR LATINO¹	0.5207		0.7336		0.2328	
White	0.9485		0.6614		0.1588	
Black or African American	0.0641		0.0262		0.9342	
American Indian or Alaska Native	0.3066		0.8269		0.6744	
Native Hawaiian or Other Pacific Islander	*		*		0.2603	
Chinese	0.3174		0.4880		*	
Filipino	0.2598		0.8746		0.3087	
Japanese	*		0.8983		0.7242	
Asian Indian	0.7215		0.8121		0.6495	
Korean	*		*		*	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0808		0.4135		0.2984	
Mexican	0.0516		0.1470		0.4979	
Puerto Rican	0.7103		0.1783		0.8885	
Central or South American	0.2602		0.3017		0.4529	
Cuban	0.4574		0.7020		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.71C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	679	667	507	483	285	273
NOT HISPANIC OR LATINO¹	643	637	489	461	262	255
White	569	569	427	415	236	240
Black or African American	190	216	158	162	88	84
American Indian or Alaska Native	47	38	30	33	19	21
Native Hawaiian or Other Pacific Islander	28	25	20	22	10	9
Chinese	69	67	37	37	9	20
Filipino	75	96	39	43	14	18
Japanese	46	46	22	22	5	5
Asian Indian	59	62	37	35	14	8
Korean	39	36	27	25	22	12
Vietnamese	38	31	19	12	11	7
HISPANIC OR LATINO¹	231	211	182	169	94	89
Mexican	174	165	151	143	73	78
Puerto Rican	74	68	61	58	36	33
Central or South American	84	80	64	63	24	20
Cuban	43	43	42	40	*	14

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.71D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.32	0.31	0.24	0.23	0.13	0.13
NOT HISPANIC OR LATINO¹	0.35	0.34	0.26	0.25	0.14	0.14
White	0.38	0.38	0.28	0.27	0.16	0.16
Black or African American	0.81	0.90	0.67	0.68	0.38	0.35
American Indian or Alaska Native	3.95	3.46	2.56	3.03	1.64	1.90
Native Hawaiian or Other Pacific Islander	4.84	5.16	3.55	4.55	1.72	1.85
Chinese	3.62	3.20	1.94	1.79	0.49	0.93
Filipino	3.90	4.72	2.01	2.11	0.74	0.90
Japanese	5.73	5.60	2.82	2.70	0.60	0.66
Asian Indian	3.28	3.08	2.03	1.74	0.80	0.37
Korean	5.62	5.54	3.86	3.87	3.08	1.89
Vietnamese	5.31	5.01	2.70	2.02	1.61	1.09
HISPANIC OR LATINO¹	0.91	0.80	0.71	0.64	0.37	0.34
Mexican	1.05	0.97	0.91	0.84	0.44	0.46
Puerto Rican	2.74	2.60	2.25	2.22	1.33	1.28
Central or South American	2.33	2.03	1.78	1.58	0.65	0.49
Cuban	3.79	3.63	3.69	3.31	*	1.21

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.71P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Racial/Ethnic Subgroups:
P Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.2513		0.6884		0.4236	
NOT HISPANIC OR LATINO¹	0.6032		0.8977		0.1873	
White	0.8016		0.4436		0.1087	
Black or African American	0.0808		0.0281		0.9293	
American Indian or Alaska Native	0.2630		0.7644		0.7602	
Native Hawaiian or Other Pacific Islander	0.8368		0.5346		0.2609	
Chinese	0.3601		0.3436		0.1453	
Filipino	0.3119		0.9400		0.4001	
Japanese	0.1139		0.8837		0.6294	
Asian Indian	0.6452		0.7424		0.5295	
Korean	0.3985		0.8095		0.4578	
Vietnamese	0.8342		0.9228		0.7126	
HISPANIC OR LATINO¹	0.0504		0.4182		0.2904	
Mexican	0.0306		0.1448		0.4295	
Puerto Rican	0.8511		0.1470		0.9643	
Central or South American	0.2315		0.3968		0.4894	
Cuban	0.4989		0.8687		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.