

Table 6.36C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2003 and 2004

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	574	554	240	230	490	483
Marijuana and Hashish	457	450	179	179	396	395
Cocaine	232	209	87	96	213	179
Crack	123	116	60	63	103	91
Heroin	47	55	23	29	41	45
Hallucinogens	141	150	51	59	129	137
LSD	41	42	13	18	38	35
PCP	20	25	10	8	17	24
Ecstasy	102	109	39	42	94	99
Inhalants	67	78	29	52	62	63
Nonmedical Use of Psychotherapeutics ⁴	361	348	169	162	301	292
Pain Relievers	304	299	135	130	263	246
OxyContin [®]	--	72	--	40	--	61
Tranquilizers	195	203	112	113	157	154
Stimulants	124	140	61	72	106	115
Methamphetamine	93	107	43	60	82	84
Sedatives	83	70	64	39	54	51
ILLCIT DRUG OTHER THAN MARIJUANA³	425	421	188	184	366	348

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.36D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2003 and 2004

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	0.25	0.24	0.77	0.74	0.26	0.24
Marijuana and Hashish	0.21	0.20	0.60	0.61	0.21	0.20
Cocaine	0.11	0.09	0.30	0.34	0.11	0.09
Crack	0.06	0.05	0.21	0.23	0.06	0.05
Heroin	0.02	0.03	0.08	0.11	0.02	0.02
Hallucinogens	0.07	0.07	0.18	0.21	0.07	0.07
LSD	0.02	0.02	0.05	0.07	0.02	0.02
PCP	0.01	0.01	0.04	0.03	0.01	0.01
Ecstasy	0.05	0.05	0.14	0.15	0.05	0.05
Inhalants	0.03	0.04	0.10	0.19	0.03	0.03
Nonmedical Use of Psychotherapeutics ⁴	0.17	0.16	0.57	0.56	0.16	0.15
Pain Relievers	0.14	0.13	0.48	0.46	0.14	0.13
OxyContin [®]	--	0.03	--	0.15	--	0.03
Tranquilizers	0.09	0.09	0.39	0.40	0.08	0.08
Stimulants	0.06	0.07	0.22	0.26	0.06	0.06
Methamphetamine	0.04	0.05	0.15	0.22	0.04	0.05
Sedatives	0.04	0.03	0.23	0.14	0.03	0.03
ILLCIT DRUG OTHER THAN MARIJUANA³	0.19	0.19	0.62	0.62	0.20	0.18

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.36P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLICIT DRUG³	0.5923		0.8623		0.5363	
Marijuana and Hashish	0.9420		0.6276		0.9464	
Cocaine	0.3314		0.5106		0.1927	
Crack	0.6082		0.6912		0.4181	
Heroin	0.3394		0.3127		0.6059	
Hallucinogens	0.7185		0.4694		0.4977	
LSD	0.7309		0.6160		0.9425	
PCP	0.8107		0.4766		0.4980	
Ecstasy	0.1562		0.7377		0.1598	
Inhalants	0.2558		0.1689		0.6071	
Nonmedical Use of Psychotherapeutics ⁴	0.3762		0.6467		0.4617	
Pain Relievers	0.2465		0.7194		0.2734	
OxyContin [®]	--		--		--	
Tranquilizers	0.9542		0.2924		0.6176	
Stimulants	0.2507		0.3210		0.4162	
Methamphetamine	0.3971		0.1350		0.8941	
Sedatives	0.3354		0.2993		0.5213	
ILLICIT DRUG OTHER THAN MARIJUANA³	0.1745		0.7029		0.1931	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.37C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	1,042	1,015	355	365	913	908
Cigarettes	956	907	332	325	837	818
Smokeless Tobacco	314	270	104	107	291	263
Cigars	498	491	171	171	453	447
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	1,725	1,702	522	513	1,537	1,548
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	972	941	337	327	857	846
Cigarettes	901	858	322	301	796	777
Smokeless Tobacco	288	251	94	99	271	241
Cigars	353	342	125	103	316	321
Pipe Tobacco	130	146	52	65	122	131
Alcohol	1,507	1,527	438	445	1,362	1,398
Binge Alcohol Use ⁵	787	805	234	242	717	751
Heavy Alcohol Use ⁵	401	395	109	135	378	364

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.37D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	0.38	0.38	1.03	1.03	0.40	0.40
Cigarettes	0.38	0.36	0.99	0.97	0.39	0.39
Smokeless Tobacco	0.15	0.12	0.37	0.38	0.16	0.14
Cigars	0.21	0.21	0.58	0.59	0.23	0.22
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	0.42	0.41	1.01	1.07	0.44	0.43
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	0.37	0.37	0.97	0.97	0.39	0.39
Cigarettes	0.36	0.36	0.96	0.93	0.38	0.38
Smokeless Tobacco	0.13	0.11	0.33	0.35	0.15	0.12
Cigars	0.16	0.15	0.43	0.37	0.17	0.16
Pipe Tobacco	0.06	0.07	0.19	0.23	0.07	0.07
Alcohol	0.43	0.44	1.06	1.09	0.46	0.46
Binge Alcohol Use ⁵	0.32	0.32	0.73	0.79	0.34	0.34
Heavy Alcohol Use ⁵	0.18	0.18	0.37	0.47	0.20	0.19

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.37P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²	
			Received	Not Received
	2003	2004	2003	2004
PAST YEAR				
Tobacco Product ³	0.2632		0.7008	0.1790
Cigarettes	0.5391		0.6364	0.4285
Smokeless Tobacco	0.0670		0.7135	0.0401
Cigars	0.7294		0.7017	0.8670
Pipe Tobacco ⁴	--		--	--
Alcohol	0.7571		0.9755	0.6906
Binge Alcohol Use ⁵	--		--	--
Heavy Alcohol Use ⁵	--		--	--
PAST MONTH				
Tobacco Product ³	0.2030		0.9588	0.1878
Cigarettes	0.2428		0.8920	0.2503
Smokeless Tobacco	0.0557		0.6848	0.0335
Cigars	0.1640		0.6737	0.1928
Pipe Tobacco	0.3509		0.7381	0.2436
Alcohol	0.6725		0.1864	0.9948
Binge Alcohol Use ⁵	0.8457		0.0082	0.5186
Heavy Alcohol Use ⁵	0.5272		0.0322	0.9869

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.38C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	428	454	196	202	368	391
Illicit Drug ⁴	205	213	112	120	166	169
Alcohol	397	423	167	183	345	360
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	303	330	150	167	256	279
Illicit Drug ⁴	159	168	97	106	124	132
Alcohol	269	301	122	147	231	248

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.38D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	0.19	0.20	0.65	0.67	0.20	0.20
Illicit Drug ⁴	0.09	0.10	0.39	0.42	0.09	0.09
Alcohol	0.18	0.19	0.57	0.63	0.19	0.19
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	0.14	0.15	0.52	0.58	0.14	0.14
Illicit Drug ⁴	0.07	0.08	0.34	0.37	0.07	0.07
Alcohol	0.12	0.14	0.43	0.52	0.12	0.13

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.38P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling; P Values from Tests of Differences of Percentages, 2004 Versus 2003

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²	
			Received	Not Received
	2003	2004	2003	2004
DEPENDENCE OR ABUSE³				
Illicit Drug or Alcohol ⁴	0.2635		0.0108	0.9425
Illicit Drug ⁴	0.2373		0.1283	0.6320
Alcohol	0.2887		0.0162	0.9034
DEPENDENCE³				
Illicit Drug or Alcohol ⁴	0.0789		0.0162	0.4309
Illicit Drug ⁴	0.0870		0.1329	0.2548
Alcohol	0.0791		0.0104	0.4683

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.