

Table 7.121C No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drug or Alcohol ¹		Illicit Drug ¹		Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	89	88	67	67	82	81
RACE²						
American Indian or Alaska Native	6	6	6	7	6	6
Asian or Pacific Islander	16	18	14	13	14	16
Asian Only	16	18	13	13	14	16
Native Hawaiian or Other Pacific Islander Only	*	*	*	2	*	*
Black or African American	32	28	24	23	28	24
White	80	81	61	61	73	75
Two or More Races	11	11	11	8	10	10
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	36	42	28	30	34	41
Not Hispanic or Latino	83	76	61	59	76	71
Black or African American	31	27	23	22	28	24
White	71	67	53	53	65	62
GENDER						
Male	63	62	47	44	57	59
Female	62	66	45	50	60	59
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	36	39	28	32	31	34
100-199%	43	40	30	31	39	37
200% or More	72	75	53	53	66	70
REGION						
Northeast	36	41	29	34	34	36
Midwest	38	38	28	25	36	34
South	57	51	41	39	48	46
West	45	45	34	35	44	44

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.121D No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drug or Alcohol ¹		Illicit Drug ¹		Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	0.36	0.35	0.27	0.27	0.33	0.32
RACE²						
American Indian or Alaska Native	1.96	1.96	1.77	2.08	1.84	1.84
Asian or Pacific Islander	1.52	1.66	1.27	1.18	1.33	1.47
Asian Only	1.59	1.79	1.32	1.26	1.39	1.58
Native Hawaiian or Other Pacific Islander Only	*	*	*	2.64	*	*
Black or African American	0.81	0.70	0.61	0.57	0.72	0.61
White	0.41	0.42	0.32	0.32	0.38	0.39
Two or More Races	2.48	2.29	2.39	1.77	2.33	2.20
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.88	0.99	0.68	0.71	0.83	0.96
Not Hispanic or Latino	0.40	0.36	0.29	0.28	0.36	0.34
Black or African American	0.83	0.71	0.63	0.57	0.75	0.62
White	0.46	0.43	0.34	0.34	0.42	0.40
GENDER						
Male	0.49	0.48	0.37	0.34	0.45	0.45
Female	0.51	0.53	0.37	0.40	0.49	0.48
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.89	0.90	0.69	0.75	0.76	0.78
100-199%	0.76	0.72	0.53	0.55	0.69	0.67
200% or More	0.47	0.49	0.35	0.35	0.43	0.46
REGION						
Northeast	0.80	0.89	0.65	0.75	0.76	0.80
Midwest	0.66	0.67	0.49	0.43	0.62	0.60
South	0.64	0.56	0.46	0.43	0.54	0.52
West	0.76	0.74	0.57	0.58	0.75	0.73

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.121P No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drug or Alcohol ¹		Illicit Drug ¹		Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	0.7468		0.0803		0.8159	
RACE²						
American Indian or Alaska Native	0.3216		0.4288		0.9621	
Asian or Pacific Islander	0.8871		0.5622		0.8625	
Asian Only	0.9050		0.8091		0.8115	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.6054		0.8818		0.6153	
White	0.8090		0.1158		0.9326	
Two or More Races	0.9758		0.2447		0.9704	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.5160		0.4417		0.1324	
Not Hispanic or Latino	0.4994		0.1199		0.3559	
Black or African American	0.6496		0.7668		0.7070	
White	0.4206		0.1559		0.3028	
GENDER						
Male	0.7481		0.0724		0.8737	
Female	0.8773		0.4285		0.6300	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.8791		0.9329		0.7495	
100-199%	0.1479		0.2215		0.1588	
200% or More	0.7459		0.1386		0.7096	
REGION						
Northeast	0.9576		0.8362		0.7430	
Midwest	0.3875		0.5287		0.8976	
South	0.0467		0.0325		0.1438	
West	0.3485		0.5377		0.2276	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.122C Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	67	67	60	57	67	73
RACE³						
American Indian or Alaska Native	6	7	5	8	6	5
Asian or Pacific Islander	14	13	9	11	8	13
Asian Only	13	13	8	11	8	13
Native Hawaiian or Other Pacific Islander Only	*	2	*	2	*	*
Black or African American	24	23	20	19	20	18
White	61	61	53	53	62	68
Two or More Races	11	8	9	8	10	9
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	28	30	23	26	27	38
Not Hispanic or Latino	61	59	55	52	61	63
Black or African American	23	22	20	19	20	17
White	53	53	47	47	55	56
GENDER						
Male	47	44	43	39	46	52
Female	45	50	39	38	46	50
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	28	32	22	26	24	28
100-199%	30	31	27	27	30	32
200% or More	53	53	46	45	54	66
REGION						
Northeast	29	34	29	27	28	28
Midwest	28	25	24	23	29	29
South	41	39	37	31	38	38
West	34	35	29	31	37	48

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.122D Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	0.27	0.27	0.24	0.23	0.27	0.29
RACE³						
American Indian or Alaska Native	1.77	2.08	1.62	2.44	1.71	1.55
Asian or Pacific Islander	1.27	1.18	0.87	1.00	0.78	1.22
Asian Only	1.32	1.26	0.85	1.07	0.77	1.28
Native Hawaiian or Other Pacific Islander Only	*	2.64	*	2.18	*	*
Black or African American	0.61	0.57	0.52	0.49	0.52	0.45
White	0.32	0.32	0.28	0.27	0.32	0.35
Two or More Races	2.39	1.77	2.15	1.68	2.23	1.92
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.68	0.71	0.57	0.61	0.67	0.90
Not Hispanic or Latino	0.29	0.28	0.26	0.25	0.29	0.30
Black or African American	0.63	0.57	0.53	0.49	0.54	0.45
White	0.34	0.34	0.30	0.30	0.35	0.36
GENDER						
Male	0.37	0.34	0.34	0.30	0.36	0.40
Female	0.37	0.40	0.32	0.31	0.38	0.41
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.69	0.75	0.53	0.60	0.58	0.65
100-199%	0.53	0.55	0.48	0.49	0.54	0.58
200% or More	0.35	0.35	0.30	0.30	0.36	0.43
REGION						
Northeast	0.65	0.75	0.64	0.58	0.63	0.63
Midwest	0.49	0.43	0.43	0.41	0.51	0.52
South	0.46	0.43	0.41	0.35	0.42	0.42
West	0.57	0.58	0.50	0.52	0.64	0.80

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.122P Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0803		0.3316		0.2617	
RACE³						
American Indian or Alaska Native	0.4288		0.6534		0.8183	
Asian or Pacific Islander	0.5622		0.6509		0.4371	
Asian Only	0.8091		0.3684		0.3105	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.8818		0.4692		0.7908	
White	0.1158		0.2026		0.3039	
Two or More Races	0.2447		0.6717		0.9432	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.4417		0.8935		0.0074	
Not Hispanic or Latino	0.1199		0.3272		0.9448	
Black or African American	0.7668		0.4993		0.6944	
White	0.1559		0.1468		0.6427	
GENDER						
Male	0.0724		0.2279		0.3716	
Female	0.4285		0.7954		0.4607	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.9329		0.7318		0.3687	
100-199%	0.2215		0.5901		0.7155	
200% or More	0.1386		0.2631		0.2390	
REGION						
Northeast	0.8362		0.8380		0.8779	
Midwest	0.5287		0.2154		0.5416	
South	0.0325		0.1354		0.7925	
West	0.5377		0.3833		0.0931	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.123C Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	396	404	317	349	675	699
RACE³						
American Indian or Alaska Native	23	25	22	21	47	56
Asian or Pacific Islander	63	69	28	45	103	118
Asian Only	61	58	27	35	99	112
Native Hawaiian or Other Pacific Islander Only	13	*	9	*	22	*
Black or African American	140	132	115	112	232	231
White	350	374	292	316	612	638
Two or More Races	39	42	34	29	52	64
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	130	129	91	122	242	238
Not Hispanic or Latino	359	383	306	328	643	659
Black or African American	136	128	114	109	228	228
White	319	352	280	295	569	596
GENDER						
Male	280	298	241	267	517	511
Female	239	227	172	189	375	389
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	138	153	108	138	209	217
100-199%	167	155	136	139	269	302
200% or More	310	315	259	271	572	585
EDUCATION						
< High School	174	144	126	125	279	284
High School Graduate	207	216	171	180	366	400
Some College/College Graduate	275	290	224	256	483	495

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.123D Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	0.19	0.19	0.15	0.16	0.32	0.32
RACE³						
American Indian or Alaska Native	1.22	1.30	1.16	1.09	2.52	2.93
Asian or Pacific Islander	0.67	0.71	0.30	0.47	1.11	1.22
Asian Only	0.69	0.65	0.30	0.39	1.11	1.24
Native Hawaiian or Other Pacific Islander Only	2.98	*	1.97	*	4.91	*
Black or African American	0.57	0.53	0.47	0.45	0.95	0.93
White	0.20	0.21	0.17	0.18	0.35	0.36
Two or More Races	1.61	1.70	1.40	1.17	2.17	2.59
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.50	0.48	0.35	0.45	0.94	0.89
Not Hispanic or Latino	0.19	0.20	0.16	0.17	0.34	0.35
Black or African American	0.58	0.53	0.48	0.45	0.96	0.94
White	0.21	0.23	0.19	0.19	0.38	0.39
GENDER						
Male	0.27	0.29	0.24	0.26	0.51	0.49
Female	0.22	0.20	0.16	0.17	0.34	0.35
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.59	0.60	0.46	0.54	0.90	0.85
100-199%	0.42	0.40	0.34	0.35	0.67	0.77
200% or More	0.21	0.21	0.17	0.18	0.38	0.39
EDUCATION						
< High School	0.46	0.39	0.33	0.34	0.74	0.78
High School Graduate	0.31	0.32	0.26	0.26	0.55	0.59
Some College/College Graduate	0.26	0.26	0.21	0.23	0.45	0.45

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.123P Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	0.3915		0.7409		0.8457	
RACE³						
American Indian or Alaska Native	0.8902		0.5498		0.6441	
Asian or Pacific Islander	0.9843		0.3833		0.3636	
Asian Only	0.4548		0.8662		0.4266	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.8029		0.4659		0.5473	
White	0.3030		0.5265		0.7709	
Two or More Races	0.6667		0.7857		0.5541	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.2891		0.9201		0.5876	
Not Hispanic or Latino	0.5988		0.7162		0.6959	
Black or African American	0.9826		0.6645		0.5172	
White	0.5318		0.5421		0.5640	
GENDER						
Male	0.9804		0.8411		0.9016	
Female	0.1524		0.7318		0.8870	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.3124		0.0359		0.1570	
100-199%	0.0555		0.2035		0.2903	
200% or More	0.5216		0.4165		0.8193	
EDUCATION						
< High School	0.5092		0.7832		0.2445	
High School Graduate	0.3314		0.2865		0.8863	
Some College/College Graduate	0.9953		0.5699		0.3887	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.124C Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2003	2004
TOTAL	48	45
RACE¹		
American Indian or Alaska Native	3	4
Asian or Pacific Islander	13	15
Asian Only	12	15
Native Hawaiian or Other Pacific Islander Only	*	1
Black or African American	15	13
White	41	41
Two or More Races	5	6
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	22	22
Not Hispanic or Latino	43	41
Black or African American	15	12
White	36	37
GENDER		
Male	33	34
Female	31	34
FAMILY INCOME (% of Federal Poverty Threshold)		
Less than 100%	19	19
100-199%	19	20
200% or More	38	37
REGION		
Northeast	18	18
Midwest	17	17
South	29	30
West	29	23

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.124D Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2003	2004
TOTAL	0.19	0.18
RACE¹		
American Indian or Alaska Native	0.78	1.18
Asian or Pacific Islander	1.16	1.42
Asian Only	1.19	1.53
Native Hawaiian or Other Pacific Islander Only	*	0.95
Black or African American	0.38	0.31
White	0.22	0.21
Two or More Races	1.25	1.29
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	0.55	0.53
Not Hispanic or Latino	0.21	0.20
Black or African American	0.39	0.32
White	0.23	0.24
GENDER		
Male	0.26	0.26
Female	0.25	0.27
FAMILY INCOME (% of Federal Poverty Threshold)		
Less than 100%	0.46	0.43
100-199%	0.35	0.35
200% or More	0.25	0.24
REGION		
Northeast	0.40	0.39
Midwest	0.31	0.31
South	0.32	0.33
West	0.49	0.38

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.124P Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2003	2004
TOTAL	0.5597	
RACE¹		
American Indian or Alaska Native	0.1646	
Asian or Pacific Islander	0.6904	
Asian Only	0.4948	
Native Hawaiian or Other Pacific Islander Only	*	
Black or African American	0.4144	
White	0.9118	
Two or More Races	0.6593	
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	0.5019	
Not Hispanic or Latino	0.7761	
Black or African American	0.5979	
White	0.8698	
GENDER		
Male	0.5690	
Female	0.8074	
FAMILY INCOME (% of Federal Poverty Threshold)		
Less than 100%	0.6080	
100-199%	0.6210	
200% or More	0.4657	
REGION		
Northeast	0.1683	
Midwest	0.8921	
South	1.0000	
West	0.9641	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.125C Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2003	2004	2003	2004	2003	2004
TOTAL	24	24	57	55	43	40
RACE¹						
American Indian or Alaska Native	2	1	4	3	3	3
Asian or Pacific Islander	1	*	8	12	8	9
Asian Only	1	*	8	11	7	8
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	10	11	23	22	13	16
White	20	21	50	46	39	36
Two or More Races	3	3	6	10	4	5
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	11	13	29	27	19	16
Not Hispanic or Latino	20	20	48	47	37	37
Black or African American	10	11	22	21	13	16
White	15	17	40	38	33	32
GENDER						
Male	19	19	39	39	31	29
Female	13	15	37	37	27	27
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	12	14	27	25	20	20
100-199%	9	12	25	25	22	20
200% or More	16	15	39	39	28	29
REGION						
Northeast	7	9	21	27	14	13
Midwest	9	11	23	20	20	18
South	14	15	36	31	27	27
West	16	12	31	30	22	21

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.125D Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.28	0.28	0.61	0.58	0.47	0.46
RACE¹						
American Indian or Alaska Native	2.17	1.16	3.30	3.13	3.08	3.08
Asian or Pacific Islander	0.28	*	2.28	3.02	2.27	2.37
Asian Only	0.24	*	2.49	3.21	2.17	2.26
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	0.70	0.74	1.45	1.42	0.93	1.09
White	0.31	0.33	0.68	0.65	0.56	0.53
Two or More Races	1.80	1.54	3.39	4.53	2.57	2.79
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.78	0.87	1.66	1.65	1.27	1.05
Not Hispanic or Latino	0.28	0.29	0.62	0.61	0.49	0.52
Black or African American	0.73	0.78	1.48	1.43	0.96	1.12
White	0.29	0.33	0.70	0.68	0.58	0.61
GENDER						
Male	0.42	0.44	0.79	0.83	0.66	0.66
Female	0.31	0.35	0.81	0.79	0.62	0.62
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.82	0.89	1.57	1.39	1.24	1.16
100-199%	0.47	0.59	1.17	1.20	1.08	1.00
200% or More	0.34	0.31	0.74	0.71	0.54	0.57
REGION						
Northeast	0.44	0.62	1.22	1.60	0.88	0.84
Midwest	0.44	0.55	1.04	0.96	0.93	0.88
South	0.47	0.49	1.12	0.92	0.83	0.86
West	0.78	0.62	1.38	1.36	1.05	1.00

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.125P Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.4665		0.1211		0.3994	
RACE¹						
American Indian or Alaska Native	0.8480		0.6204		0.7173	
Asian or Pacific Islander	*		0.4511		0.9263	
Asian Only	*		0.4728		0.9751	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.8919		0.4229		0.6393	
White	0.4027		0.0923		0.2365	
Two or More Races	0.4822		0.9257		0.7825	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.6977		0.5287		0.2421	
Not Hispanic or Latino	0.5363		0.1384		0.7261	
Black or African American	0.8393		0.3294		0.6362	
White	0.5105		0.1280		0.4904	
GENDER						
Male	0.9269		0.2875		0.1996	
Female	0.2408		0.2097		0.8733	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.4581		0.2272		0.7129	
100-199%	0.6752		0.1190		0.0116	
200% or More	0.9493		0.6069		0.3284	
REGION						
Northeast	0.3883		0.6299		0.3094	
Midwest	0.1244		0.7966		0.6016	
South	0.8858		0.0498		0.1078	
West	0.7232		0.6360		0.9802	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.126C Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2003	2004	2003	2004	2003	2004
TOTAL	101	109	112	106	103	101
RACE²						
American Indian or Alaska Native	7	8	9	8	9	7
Asian or Pacific Islander	24	27	26	25	27	31
Asian Only	23	27	26	25	27	30
Native Hawaiian or Other Pacific Islander Only	*	*	5	*	5	*
Black or African American	40	40	44	49	44	43
White	86	97	94	94	85	90
Two or More Races	13	12	13	12	14	11
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	43	56	48	47	49	52
Not Hispanic or Latino	93	92	101	93	92	94
Black or African American	39	40	43	48	44	41
White	77	77	82	79	74	79
GENDER						
Male	70	73	74	72	72	78
Female	75	73	83	74	76	76
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	44	48	46	47	42	49
100-199%	48	53	47	52	53	51
200% or More	81	77	91	82	77	77
REGION						
Northeast	45	49	45	39	41	40
Midwest	41	43	46	47	39	43
South	61	61	65	62	67	65
West	52	64	63	62	54	50

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.126D Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2003	2004	2003	2004	2003	2004
TOTAL	0.41	0.44	0.46	0.43	0.42	0.40
RACE²						
American Indian or Alaska Native	2.35	2.59	2.76	2.66	2.74	2.39
Asian or Pacific Islander	2.26	2.54	2.44	2.36	2.56	2.86
Asian Only	2.43	2.72	2.65	2.51	2.77	3.04
Native Hawaiian or Other Pacific Islander Only	*	*	5.85	*	5.94	*
Black or African American	1.03	1.02	1.16	1.26	1.15	1.08
White	0.45	0.50	0.50	0.49	0.45	0.47
Two or More Races	2.93	2.56	3.08	2.55	3.10	2.45
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	1.08	1.34	1.20	1.13	1.22	1.25
Not Hispanic or Latino	0.45	0.44	0.49	0.45	0.44	0.45
Black or African American	1.06	1.06	1.18	1.29	1.19	1.09
White	0.50	0.50	0.54	0.52	0.48	0.52
GENDER						
Male	0.56	0.58	0.59	0.57	0.57	0.61
Female	0.62	0.60	0.69	0.61	0.63	0.62
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	1.09	1.11	1.16	1.11	1.06	1.15
100-199%	0.86	0.97	0.85	0.95	0.96	0.93
200% or More	0.53	0.51	0.60	0.54	0.51	0.51
REGION						
Northeast	1.01	1.09	1.03	0.86	0.93	0.89
Midwest	0.73	0.76	0.82	0.83	0.69	0.77
South	0.70	0.68	0.74	0.70	0.76	0.72
West	0.89	1.08	1.09	1.05	0.92	0.83

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.126P Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2003	2004	2003	2004	2003	2004
TOTAL	0.9528		0.0039		0.4992	
RACE²						
American Indian or Alaska Native	0.2060		0.0538		0.1120	
Asian or Pacific Islander	0.9158		0.8379		0.5843	
Asian Only	0.9755		0.8942		0.5565	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.6148		0.3018		0.5913	
White	0.7842		0.0017		0.4480	
Two or More Races	0.7323		0.7068		0.3384	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.4414		0.3545		0.7774	
Not Hispanic or Latino	0.7214		0.0063		0.3630	
Black or African American	0.7048		0.2812		0.7021	
White	0.5399		0.0030		0.2106	
GENDER						
Male	0.4521		0.0023		0.0834	
Female	0.4267		0.2404		0.4039	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.6541		0.1010		0.7535	
100-199%	0.3846		0.8804		0.7755	
200% or More	0.8206		0.0081		0.3436	
REGION						
Northeast	0.9845		0.2051		0.7528	
Midwest	0.2711		0.0937		0.1436	
South	0.3289		0.0529		0.6880	
West	0.9163		0.3945		0.9542	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.127C Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drug		Alcohol		Either Illicit Drug or Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	90	114	102	136	117	166
RACE¹						
American Indian or Alaska Native	11	3	13	*	14	*
Asian or Pacific Islander	6	*	7	*	8	*
Asian Only	6	*	*	*	7	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	51	63	58	59	60	74
White	71	81	80	120	96	139
Two or More Races	9	11	10	11	10	11
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	21	38	34	95	37	100
Not Hispanic or Latino	88	108	96	98	112	132
Black or African American	51	63	58	59	60	74
White	68	72	72	74	89	96
GENDER						
Male	74	88	86	131	98	144
Female	50	72	54	47	64	84
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	49	69	55	82	62	100
100-199%	45	57	50	49	58	65
200% or More	54	69	62	72	74	92
AGE GROUP						
12-17	14	16	12	15	17	18
18 or Older	88	113	102	135	116	164

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.127D Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drug		Alcohol		Either Illicit Drug or Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	0.39	0.46	0.44	0.55	0.50	0.65
RACE¹						
American Indian or Alaska Native	2.95	0.92	3.42	*	3.72	*
Asian or Pacific Islander	0.92	*	1.05	*	1.12	*
Asian Only	1.02	*	*	*	1.07	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	1.99	2.16	2.25	2.05	2.30	2.47
White	0.38	0.41	0.42	0.58	0.50	0.67
Two or More Races	2.68	2.86	2.77	2.80	2.86	3.00
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.70	1.14	1.10	2.74	1.18	2.84
Not Hispanic or Latino	0.44	0.51	0.48	0.47	0.55	0.61
Black or African American	2.06	2.24	2.32	2.13	2.37	2.56
White	0.42	0.43	0.45	0.43	0.54	0.56
GENDER						
Male	0.50	0.56	0.58	0.80	0.65	0.87
Female	0.62	0.86	0.67	0.57	0.78	0.97
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	1.33	1.53	1.47	1.73	1.60	2.01
100-199%	0.87	1.23	0.98	1.05	1.10	1.37
200% or More	0.39	0.45	0.44	0.47	0.53	0.59
AGE GROUP						
12-17	0.61	0.67	0.52	0.63	0.74	0.74
18 or Older	0.43	0.51	0.49	0.60	0.55	0.71

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.127P Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drug		Alcohol		Either Illicit Drug or Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0749		0.3360		0.0811	
RACE¹						
American Indian or Alaska Native	0.0401		*		*	
Asian or Pacific Islander	*		*		*	
Asian Only	*		*		*	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.1738		1.0000		0.2998	
White	0.3366		0.3027		0.2276	
Two or More Races	0.9706		0.9635		0.8723	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.3102		0.3902		0.2933	
Not Hispanic or Latino	0.1173		0.5661		0.1578	
Black or African American	0.1660		0.9185		0.2513	
White	0.6360		0.6106		0.5780	
GENDER						
Male	0.2708		0.2270		0.2110	
Female	0.1319		0.6683		0.2264	
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	0.2125		0.3839		0.0949	
100-199%	0.0518		0.4491		0.1100	
200% or More	0.9261		0.9447		0.9390	
AGE GROUP						
12-17	0.3254		0.1153		0.5346	
18 or Older	0.0967		0.4357		0.0945	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.