

**Table 1.19C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	909	969	558	535	404	415
12	30	27	20	21	13	14
13	35	38	29	28	19	19
14	37	40	34	36	22	27
15	43	43	38	39	30	32
16	46	43	42	39	35	32
17	46	46	43	44	36	34
18	54	54	55	51	46	42
19	51	51	52	52	47	42
20	50	51	51	51	41	42
21	48	49	44	55	38	47
22	53	47	48	50	36	44
23	49	48	45	43	34	35
24	44	52	45	42	38	36
25	45	48	41	43	34	33
26-29	180	181	157	160	138	124
30-34	210	208	159	143	120	124
35-39	228	237	166	164	131	110
40-44	247	235	163	188	130	135
45-49	225	228	183	158	138	130
50-54	328	342	160	199	126	138
55-59	259	279	105	114	64	77
60-64	217	228	85	72	47	54
65 or Older	296	272	78	83	74	46

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.19D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.38	0.40	0.23	0.22	0.17	0.17
12	0.72	0.68	0.48	0.53	0.33	0.36
13	0.81	0.86	0.67	0.65	0.44	0.44
14	0.90	0.92	0.82	0.81	0.53	0.62
15	1.03	1.01	0.91	0.91	0.72	0.74
16	1.10	1.04	1.00	0.95	0.84	0.77
17	1.12	1.13	1.04	1.08	0.89	0.83
18	1.17	1.17	1.19	1.11	1.00	0.92
19	1.32	1.30	1.34	1.31	1.21	1.07
20	1.24	1.27	1.26	1.27	1.01	1.05
21	1.22	1.21	1.12	1.35	0.96	1.15
22	1.33	1.18	1.18	1.24	0.90	1.10
23	1.22	1.23	1.13	1.08	0.86	0.88
24	1.22	1.29	1.23	1.03	1.05	0.90
25	1.24	1.34	1.14	1.18	0.93	0.93
26-29	1.22	1.18	1.07	1.05	0.93	0.81
30-34	1.04	1.06	0.79	0.73	0.60	0.63
35-39	1.09	1.13	0.79	0.78	0.62	0.52
40-44	1.09	1.04	0.72	0.83	0.57	0.60
45-49	1.05	1.06	0.86	0.73	0.65	0.60
50-54	1.64	1.76	0.80	1.02	0.63	0.71
55-59	1.74	1.70	0.71	0.69	0.43	0.47
60-64	1.84	1.82	0.73	0.58	0.40	0.43
65 or Older	0.86	0.79	0.23	0.24	0.22	0.13

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.19P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.2373		0.4285		0.2576	
12	0.2986		0.5077		0.8731	
13	0.8181		0.7434		0.7093	
14	0.4099		0.4136		0.4829	
15	0.7423		0.5971		0.5657	
16	0.4214		0.1214		0.0069	
17	0.9856		0.4152		0.6136	
18	0.9804		0.7460		0.3164	
19	0.3696		0.4850		0.6391	
20	0.0932		0.2081		0.0639	
21	0.6860		0.3743		0.5100	
22	0.5351		0.3301		0.4799	
23	0.0183		0.0110		0.0295	
24	0.1542		0.1088		0.4456	
25	0.8143		0.6479		0.6825	
26-29	0.2056		0.9434		0.8604	
30-34	0.1280		0.3927		0.4843	
35-39	0.1312		0.3982		0.1258	
40-44	0.7916		0.6845		0.5244	
45-49	0.7448		0.5224		0.9800	
50-54	0.0664		0.2113		0.3509	
55-59	0.9590		0.5174		0.3461	
60-64	0.8821		0.3431		0.9584	
65 or Older	0.1747		0.5773		0.3071	

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.20C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	868	912	455	451	327	360
12	10	11	8	9	7	5
13	20	20	17	18	9	12
14	30	31	27	28	19	19
15	37	37	33	33	25	28
16	45	41	40	37	34	29
17	42	45	39	41	32	31
18	52	53	54	49	44	39
19	55	53	50	51	45	40
20	52	52	49	48	38	40
21	51	51	43	50	35	44
22	53	52	44	48	34	41
23	49	50	40	40	31	31
24	46	50	42	40	35	34
25	46	48	39	40	32	32
26-29	188	190	141	143	120	113
30-34	208	212	140	121	100	107
35-39	224	234	135	139	109	95
40-44	255	241	134	153	108	115
45-49	230	229	136	137	95	114
50-54	318	330	127	178	93	132
55-59	252	276	83	87	62	62
60-64	200	210	33	30	30	16
65 or Older	238	240	21	42	18	30

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.20D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.37	0.38	0.19	0.19	0.14	0.15
12	0.25	0.29	0.20	0.23	0.16	0.13
13	0.47	0.46	0.39	0.41	0.21	0.28
14	0.72	0.70	0.64	0.65	0.45	0.44
15	0.90	0.86	0.81	0.76	0.61	0.64
16	1.06	1.00	0.94	0.90	0.80	0.70
17	1.03	1.11	0.96	1.00	0.79	0.77
18	1.14	1.16	1.17	1.06	0.94	0.84
19	1.41	1.34	1.30	1.29	1.15	1.02
20	1.28	1.31	1.21	1.20	0.95	1.00
21	1.29	1.26	1.10	1.23	0.90	1.08
22	1.32	1.28	1.09	1.19	0.84	1.01
23	1.22	1.26	1.00	1.01	0.77	0.79
24	1.27	1.24	1.16	0.98	0.95	0.84
25	1.26	1.34	1.06	1.12	0.88	0.88
26-29	1.28	1.24	0.95	0.93	0.81	0.74
30-34	1.03	1.08	0.70	0.62	0.49	0.55
35-39	1.07	1.12	0.65	0.67	0.52	0.45
40-44	1.12	1.06	0.59	0.67	0.48	0.51
45-49	1.07	1.06	0.63	0.64	0.44	0.53
50-54	1.59	1.70	0.64	0.92	0.46	0.68
55-59	1.69	1.68	0.56	0.53	0.42	0.38
60-64	1.70	1.67	0.28	0.24	0.26	0.12
65 or Older	0.69	0.69	0.06	0.12	0.05	0.09

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.20P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.4176		0.9024		0.6243	
12	0.6353		0.6656		0.5225	
13	0.6518		0.2788		0.2209	
14	0.2304		0.4075		0.6072	
15	0.7167		0.3188		0.6132	
16	0.4136		0.2564		0.0193	
17	0.6072		0.8932		0.8350	
18	0.4753		0.7725		0.0866	
19	0.9680		0.8569		0.3169	
20	0.1149		0.4065		0.3864	
21	0.8382		0.9126		0.8732	
22	0.3100		0.7379		0.6594	
23	0.0124		0.0636		0.1028	
24	0.7036		0.5491		0.7113	
25	0.4300		0.7989		0.8174	
26-29	0.3597		0.6014		0.7722	
30-34	0.1453		0.0855		0.5156	
35-39	0.0607		0.7590		0.2176	
40-44	0.7898		0.7226		0.7188	
45-49	0.8163		0.4367		0.1871	
50-54	0.0280		0.0531		0.0960	
55-59	0.8527		0.6485		0.9129	
60-64	0.7483		0.8188		0.4275	
65 or Older	0.3894		0.3358		0.7494	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.21C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	565	636	229	205	147	119
12	5	2	5	2	2	*
13	7	5	3	5	3	3
14	11	11	9	6	6	3
15	14	13	12	10	7	7
16	20	17	16	15	9	6
17	23	21	17	18	10	11
18	31	30	24	24	16	14
19	32	35	24	27	15	14
20	33	31	23	24	12	13
21	34	39	25	31	13	18
22	37	39	26	26	15	14
23	38	36	26	23	14	14
24	35	36	25	22	14	14
25	34	36	20	19	12	11
26-29	149	138	97	83	50	43
30-34	155	154	74	74	52	48
35-39	196	207	78	65	55	45
40-44	228	226	76	86	49	47
45-49	202	222	73	56	37	38
50-54	237	244	73	54	66	28
55-59	144	149	28	24	*	23
60-64	99	78	*	23	*	22
65 or Older	81	94	14	16	14	12

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.21D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.24	0.26	0.10	0.09	0.06	0.05
12	0.12	0.06	0.11	0.05	0.04	*
13	0.15	0.12	0.08	0.11	0.07	0.08
14	0.26	0.24	0.21	0.14	0.15	0.07
15	0.33	0.30	0.30	0.24	0.18	0.17
16	0.47	0.42	0.38	0.36	0.21	0.13
17	0.55	0.53	0.42	0.44	0.24	0.27
18	0.67	0.65	0.52	0.52	0.35	0.31
19	0.83	0.90	0.63	0.69	0.38	0.35
20	0.83	0.77	0.58	0.61	0.29	0.34
21	0.87	0.95	0.63	0.76	0.34	0.44
22	0.93	0.97	0.66	0.66	0.38	0.34
23	0.94	0.92	0.65	0.58	0.35	0.36
24	0.97	0.89	0.70	0.54	0.40	0.34
25	0.95	1.00	0.56	0.52	0.33	0.30
26-29	1.01	0.90	0.66	0.54	0.34	0.28
30-34	0.77	0.78	0.37	0.38	0.26	0.24
35-39	0.94	0.99	0.37	0.31	0.26	0.22
40-44	1.01	1.00	0.34	0.38	0.22	0.21
45-49	0.94	1.03	0.34	0.26	0.17	0.18
50-54	1.18	1.26	0.37	0.28	0.33	0.15
55-59	0.97	0.91	0.19	0.15	*	0.14
60-64	0.84	0.62	*	0.18	*	0.18
65 or Older	0.24	0.27	0.04	0.05	0.04	0.03

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 1.21P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.1760		0.2923		0.1297	
12	0.4189		0.3410		*	
13	0.5967		0.5034		0.6160	
14	0.4366		0.0813		0.2482	
15	1.0000		0.5833		0.7713	
16	0.8013		0.8883		0.1496	
17	0.6830		0.8926		0.8283	
18	0.4372		0.4436		0.1100	
19	0.8443		0.4689		0.7809	
20	0.7512		0.2121		0.1438	
21	0.1938		0.0456		0.0625	
22	0.1237		0.2330		0.7566	
23	0.4331		0.2539		1.0000	
24	0.6077		0.3238		0.5428	
25	0.5220		0.2220		0.1804	
26-29	0.4105		0.8790		0.5236	
30-34	0.0508		0.8446		0.9619	
35-39	0.0921		0.1666		0.0976	
40-44	0.8344		0.5177		0.6755	
45-49	0.9368		0.2820		0.8254	
50-54	0.3646		0.3305		0.1293	
55-59	0.7786		0.8211		*	
60-64	0.3566		*		*	
65 or Older	0.4114		0.6860		0.9042	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.22C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	264	280	124	117	91	66
12	*	1	*	1	*	*
13	2	3	2	3	2	3
14	5	5	4	4	2	3
15	7	6	6	3	4	1
16	9	9	7	7	5	1
17	11	8	7	5	3	3
18	16	14	9	10	7	3
19	18	17	10	9	3	5
20	17	14	9	5	5	4
21	17	19	8	12	3	7
22	18	22	7	7	1	3
23	21	18	7	8	3	5
24	21	17	10	7	2	4
25	20	19	8	4	3	3
26-29	83	76	41	40	27	18
30-34	101	87	36	38	26	25
35-39	114	121	44	42	35	24
40-44	114	128	48	43	34	26
45-49	101	105	47	28	22	22
50-54	93	105	51	50	49	24
55-59	44	62	11	14	*	14
60-64	5	22	*	7	*	*
65 or Older	23	54	*	16	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.22D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.11	0.12	0.05	0.05	0.04	0.03
12	*	0.03	*	0.03	*	*
13	0.05	0.07	0.05	0.07	0.05	0.07
14	0.13	0.11	0.10	0.08	0.04	0.06
15	0.18	0.14	0.14	0.08	0.09	0.03
16	0.21	0.22	0.18	0.17	0.13	0.03
17	0.26	0.21	0.17	0.12	0.06	0.07
18	0.34	0.31	0.20	0.23	0.15	0.06
19	0.45	0.44	0.25	0.23	0.08	0.14
20	0.43	0.34	0.22	0.13	0.14	0.11
21	0.44	0.47	0.21	0.29	0.07	0.18
22	0.44	0.55	0.17	0.18	0.02	0.06
23	0.51	0.45	0.17	0.21	0.07	0.13
24	0.59	0.43	0.27	0.18	0.06	0.09
25	0.55	0.54	0.23	0.12	0.07	0.09
26-29	0.56	0.49	0.28	0.26	0.18	0.11
30-34	0.50	0.44	0.18	0.19	0.13	0.13
35-39	0.55	0.58	0.21	0.20	0.17	0.11
40-44	0.50	0.56	0.21	0.19	0.15	0.11
45-49	0.47	0.49	0.22	0.13	0.10	0.10
50-54	0.47	0.54	0.25	0.26	0.24	0.13
55-59	0.29	0.38	0.07	0.09	*	0.09
60-64	0.04	0.17	*	0.05	*	*
65 or Older	0.07	0.16	*	0.05	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.22P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.5877		0.4826		0.1928	
12	*		*		*	
13	0.7518		0.7518		0.7518	
14	0.5605		0.5234		0.6705	
15	0.6553		0.2393		0.1672	
16	0.1549		0.2286		0.0282	
17	0.5753		0.1773		0.6776	
18	0.7994		0.7135		0.1187	
19	0.9522		0.8013		0.4248	
20	0.0927		0.0547		0.6903	
21	0.3421		0.0466		0.0271	
22	0.5178		0.9914		0.1549	
23	0.3146		0.7582		0.4168	
24	0.1587		0.2377		0.9523	
25	0.6438		0.1039		0.5207	
26-29	0.2809		0.9700		0.5880	
30-34	0.1520		0.5525		0.7102	
35-39	0.3330		0.8961		0.1828	
40-44	1.0000		0.9584		0.4513	
45-49	0.2645		0.2661		0.6596	
50-54	0.4339		0.9198		0.3199	
55-59	0.2906		0.9224		*	
60-64	0.0808		*		*	
65 or Older	0.2035		*		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.23C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	589	618	147	153	72	58
12	9	6	6	4	3	3
13	10	12	7	10	4	5
14	13	16	12	13	7	7
15	21	18	16	15	9	5
16	23	22	18	19	11	11
17	28	26	22	20	12	12
18	36	36	28	26	16	13
19	43	36	28	25	14	12
20	43	38	30	24	14	12
21	43	45	24	28	12	13
22	45	46	22	23	13	10
23	41	41	20	21	10	11
24	41	41	23	19	11	9
25	42	42	16	18	8	11
26-29	169	167	71	68	41	30
30-34	163	158	54	53	19	20
35-39	185	190	27	56	18	20
40-44	209	202	27	39	22	14
45-49	185	195	22	24	*	6
50-54	225	264	37	5	*	*
55-59	142	147	*	18	*	*
60-64	79	71	*	*	*	*
65 or Older	49	85	*	24	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.23D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.25	0.26	0.06	0.06	0.03	0.02
12	0.21	0.15	0.14	0.11	0.08	0.08
13	0.24	0.26	0.17	0.23	0.08	0.12
14	0.32	0.36	0.29	0.31	0.18	0.16
15	0.51	0.42	0.39	0.36	0.22	0.13
16	0.55	0.53	0.44	0.47	0.26	0.26
17	0.67	0.65	0.55	0.50	0.28	0.30
18	0.77	0.79	0.61	0.57	0.34	0.28
19	1.10	0.90	0.71	0.63	0.35	0.31
20	1.08	0.94	0.73	0.60	0.35	0.31
21	1.10	1.11	0.62	0.70	0.30	0.33
22	1.13	1.15	0.55	0.57	0.32	0.24
23	1.02	1.06	0.51	0.55	0.26	0.27
24	1.13	1.01	0.64	0.46	0.31	0.23
25	1.16	1.18	0.45	0.49	0.22	0.29
26-29	1.14	1.09	0.48	0.44	0.28	0.20
30-34	0.81	0.81	0.27	0.27	0.09	0.10
35-39	0.89	0.91	0.13	0.27	0.08	0.10
40-44	0.92	0.89	0.12	0.17	0.10	0.06
45-49	0.86	0.91	0.10	0.11	*	0.03
50-54	1.12	1.36	0.19	0.03	*	*
55-59	0.95	0.90	*	0.11	*	*
60-64	0.67	0.56	*	*	*	*
65 or Older	0.14	0.25	*	0.07	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.23P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.6086		0.6126		0.1840	
12	0.0718		0.3662		0.6921	
13	0.3538		0.1695		0.3711	
14	0.3250		0.2576		0.0648	
15	0.4802		0.6418		0.0383	
16	0.3470		0.9767		0.5012	
17	0.5687		0.3023		0.7038	
18	0.1057		0.2513		0.2018	
19	0.0025		0.3924		0.3861	
20	0.0006		0.0028		0.3070	
21	0.7340		0.6289		0.3595	
22	0.8980		0.3337		0.3343	
23	0.2953		0.5993		0.9440	
24	0.1005		0.1715		0.6139	
25	0.8980		0.8551		0.7523	
26-29	0.5273		0.9750		0.8786	
30-34	0.9097		0.6667		0.7439	
35-39	0.1467		0.2243		0.9703	
40-44	0.0975		0.2024		0.7954	
45-49	0.7317		0.6374		*	
50-54	0.0065		0.0821		*	
55-59	0.7272		*		*	
60-64	0.6835		*		*	
65 or Older	0.1378		*		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.24C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	441	480	83	90	45	64
12	22	23	12	15	8	9
13	28	28	21	19	11	10
14	27	31	22	23	12	10
15	27	31	18	21	10	13
16	28	28	19	16	9	7
17	26	30	17	21	7	10
18	34	33	20	19	8	9
19	33	31	14	16	5	5
20	35	33	14	16	9	5
21	35	38	11	17	2	4
22	34	39	12	14	5	10
23	35	34	15	10	8	7
24	34	37	12	11	7	4
25	32	33	9	8	5	5
26-29	135	124	25	19	4	10
30-34	138	134	13	29	8	24
35-39	150	160	16	25	12	18
40-44	154	179	25	25	24	13
45-49	169	143	25	16	12	14
50-54	171	194	9	21	9	*
55-59	101	116	*	*	*	*
60-64	53	69	*	40	*	40
65 or Older	76	83	17	*	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 1.24D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.19	0.20	0.03	0.04	0.02	0.03
12	0.53	0.57	0.30	0.38	0.20	0.23
13	0.64	0.64	0.47	0.42	0.26	0.22
14	0.65	0.72	0.54	0.52	0.28	0.23
15	0.65	0.71	0.44	0.49	0.24	0.31
16	0.68	0.68	0.47	0.38	0.22	0.16
17	0.64	0.73	0.42	0.53	0.18	0.23
18	0.73	0.71	0.43	0.42	0.18	0.20
19	0.85	0.80	0.36	0.40	0.14	0.13
20	0.88	0.82	0.35	0.39	0.21	0.13
21	0.90	0.92	0.28	0.41	0.05	0.11
22	0.85	0.98	0.29	0.35	0.11	0.24
23	0.87	0.88	0.37	0.25	0.20	0.17
24	0.94	0.90	0.32	0.28	0.20	0.09
25	0.89	0.91	0.24	0.21	0.14	0.14
26-29	0.92	0.81	0.17	0.12	0.03	0.07
30-34	0.69	0.68	0.06	0.15	0.04	0.12
35-39	0.72	0.77	0.07	0.12	0.06	0.08
40-44	0.68	0.79	0.11	0.11	0.11	0.06
45-49	0.79	0.67	0.12	0.08	0.06	0.06
50-54	0.85	1.00	0.05	0.11	0.04	*
55-59	0.68	0.70	*	*	*	*
60-64	0.45	0.55	*	0.32	*	0.32
65 or Older	0.22	0.24	0.05	*	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.24P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.4574		0.2265		0.4365	
12	0.7253		0.0272		0.3600	
13	0.8625		0.1523		0.0473	
14	0.5001		0.9134		0.8123	
15	0.2781		0.1705		0.5127	
16	0.8938		0.6740		0.4353	
17	0.5033		0.8629		0.8854	
18	0.1424		0.7321		0.4094	
19	0.2475		0.7949		0.6203	
20	0.0171		0.9914		0.1926	
21	0.8701		0.0374		0.2048	
22	0.2848		0.6514		0.4900	
23	0.5909		0.1080		0.7959	
24	0.3323		0.6533		0.4448	
25	0.4557		0.8091		0.9246	
26-29	0.9433		0.9815		0.3195	
30-34	0.2909		0.0100		0.0698	
35-39	0.8623		0.2843		0.3538	
40-44	0.9584		0.8141		0.2084	
45-49	0.2612		0.1709		0.6193	
50-54	0.6329		0.5476		*	
55-59	0.4998		*		*	
60-64	0.6552		*		*	
65 or Older	0.7726		*		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.25C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	641	729	354	353	250	210
12	21	17	15	14	9	9
13	23	26	18	22	12	14
14	25	28	23	24	15	19
15	30	33	28	25	20	17
16	35	33	29	28	20	18
17	37	37	33	32	23	20
18	44	48	36	39	25	26
19	43	44	34	38	25	27
20	44	48	38	40	26	22
21	45	50	33	41	22	28
22	44	47	31	37	20	25
23	45	45	31	33	20	21
24	42	41	34	30	24	20
25	42	40	28	28	19	17
26-29	164	166	108	112	78	75
30-34	181	176	119	109	84	76
35-39	201	199	110	114	74	65
40-44	219	224	107	120	67	70
45-49	210	211	134	102	109	61
50-54	266	293	110	101	79	52
55-59	189	188	62	82	18	51
60-64	140	136	79	53	37	25
65 or Older	161	162	75	68	72	33

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.25D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.27	0.30	0.15	0.15	0.11	0.09
12	0.50	0.44	0.36	0.36	0.22	0.24
13	0.54	0.59	0.43	0.49	0.28	0.32
14	0.61	0.65	0.56	0.55	0.37	0.42
15	0.72	0.76	0.67	0.59	0.47	0.40
16	0.84	0.81	0.70	0.67	0.48	0.44
17	0.91	0.92	0.81	0.79	0.55	0.48
18	0.95	1.05	0.78	0.85	0.54	0.57
19	1.11	1.13	0.87	0.96	0.65	0.67
20	1.08	1.21	0.93	1.00	0.65	0.55
21	1.14	1.23	0.84	1.00	0.57	0.69
22	1.10	1.17	0.77	0.92	0.51	0.63
23	1.13	1.13	0.78	0.84	0.50	0.54
24	1.15	1.02	0.93	0.73	0.65	0.50
25	1.15	1.10	0.76	0.78	0.53	0.47
26-29	1.11	1.08	0.73	0.73	0.53	0.49
30-34	0.90	0.90	0.59	0.56	0.41	0.38
35-39	0.96	0.95	0.53	0.54	0.35	0.31
40-44	0.97	0.99	0.47	0.53	0.30	0.31
45-49	0.98	0.98	0.63	0.47	0.51	0.29
50-54	1.33	1.51	0.55	0.52	0.40	0.27
55-59	1.27	1.15	0.41	0.50	0.12	0.31
60-64	1.19	1.09	0.68	0.42	0.31	0.20
65 or Older	0.47	0.47	0.22	0.20	0.21	0.10

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.25P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.6481		0.2977		0.2207	
12	0.1308		0.5089		0.3470	
13	0.3652		0.1905		0.7325	
14	0.6559		0.8300		0.2880	
15	0.4066		0.5923		0.3127	
16	0.7960		0.6513		0.0330	
17	0.8830		0.1926		0.3076	
18	0.5468		0.1455		0.3374	
19	0.1997		0.8112		0.4175	
20	0.7817		0.1730		0.0561	
21	0.1265		0.0375		0.0724	
22	0.0113		0.0030		0.2834	
23	0.4888		0.3875		0.4128	
24	0.1504		0.0249		0.3271	
25	0.4901		0.6664		0.3604	
26-29	0.6097		0.9807		0.9490	
30-34	0.1087		0.9822		0.6517	
35-39	0.1850		0.2247		0.6310	
40-44	0.3517		0.5981		0.7793	
45-49	0.8219		0.2831		0.1907	
50-54	0.0350		0.3572		0.3599	
55-59	0.7972		0.3447		0.0474	
60-64	0.4750		0.1632		0.5179	
65 or Older	0.8871		0.9686		0.1577	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.26C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	517	564	299	300	207	170
12	19	16	14	13	8	9
13	22	24	18	20	11	13
14	25	27	22	23	14	17
15	28	29	26	23	18	16
16	31	31	27	26	18	16
17	34	35	30	29	19	17
18	42	46	35	36	22	24
19	40	41	32	34	23	23
20	42	45	35	34	24	20
21	42	49	31	38	19	25
22	43	45	28	33	19	23
23	42	42	30	28	18	19
24	39	40	32	28	19	17
25	36	35	24	22	16	15
26-29	154	150	94	95	65	54
30-34	161	151	108	94	80	63
35-39	162	166	99	93	65	50
40-44	173	179	94	104	56	62
45-49	160	173	114	90	92	55
50-54	185	235	89	73	64	45
55-59	137	140	48	77	12	28
60-64	119	91	72	22	31	18
65 or Older	129	123	66	42	63	33

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.26D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.22	0.23	0.13	0.12	0.09	0.07
12	0.46	0.41	0.34	0.33	0.19	0.23
13	0.51	0.54	0.41	0.46	0.26	0.31
14	0.60	0.61	0.53	0.54	0.34	0.40
15	0.68	0.68	0.62	0.53	0.44	0.36
16	0.75	0.76	0.63	0.64	0.42	0.38
17	0.83	0.87	0.73	0.70	0.47	0.42
18	0.92	1.01	0.75	0.77	0.47	0.52
19	1.02	1.04	0.82	0.86	0.58	0.59
20	1.05	1.12	0.87	0.85	0.60	0.49
21	1.06	1.20	0.79	0.93	0.48	0.61
22	1.06	1.11	0.70	0.83	0.47	0.58
23	1.06	1.06	0.74	0.72	0.46	0.49
24	1.07	0.99	0.87	0.68	0.52	0.43
25	1.00	0.97	0.67	0.62	0.44	0.41
26-29	1.04	0.98	0.64	0.62	0.44	0.35
30-34	0.80	0.77	0.53	0.48	0.39	0.32
35-39	0.78	0.79	0.47	0.45	0.31	0.24
40-44	0.76	0.79	0.42	0.46	0.25	0.27
45-49	0.75	0.81	0.53	0.42	0.43	0.25
50-54	0.93	1.21	0.44	0.38	0.32	0.23
55-59	0.92	0.85	0.32	0.47	0.08	0.17
60-64	1.01	0.72	0.61	0.17	0.27	0.15
65 or Older	0.37	0.35	0.19	0.12	0.18	0.10

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.26P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.8008		0.1920		0.2092	
12	0.2178		0.6651		0.5269	
13	0.9752		0.6049		0.6930	
14	0.8486		0.8244		0.1498	
15	0.9055		0.3099		0.3183	
16	0.2200		0.8480		0.0577	
17	0.7005		0.3395		0.3399	
18	0.1210		0.4271		0.5652	
19	0.4031		0.9220		0.8039	
20	0.8209		0.1091		0.0210	
21	0.0295		0.0555		0.0628	
22	0.0151		0.0244		0.3217	
23	0.7530		0.1093		0.5163	
24	0.1447		0.1235		0.7600	
25	0.3845		0.1914		0.4164	
26-29	0.8843		0.8437		0.9738	
30-34	0.5840		0.9474		0.7022	
35-39	0.3332		0.1700		0.4224	
40-44	0.8410		0.6298		0.6782	
45-49	0.7625		0.2138		0.2649	
50-54	0.1292		0.5545		0.7859	
55-59	0.6444		0.1225		0.1398	
60-64	0.0987		0.0497		0.4595	
65 or Older	0.8759		0.7144		0.3510	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 1.27C Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	756	827	420	410	289	255
12	29	26	19	20	12	13
13	33	35	27	26	17	17
14	33	38	29	32	18	22
15	36	38	32	31	23	21
16	41	38	34	30	22	21
17	43	42	36	37	25	24
18	49	52	43	42	30	30
19	49	50	41	42	29	29
20	50	52	42	44	29	26
21	49	54	38	49	25	31
22	52	49	38	41	26	30
23	48	47	37	36	25	24
24	46	48	39	32	27	25
25	46	46	32	32	21	20
26-29	175	188	123	131	83	86
30-34	206	197	129	127	91	91
35-39	229	229	127	134	89	78
40-44	237	251	127	143	90	87
45-49	235	231	144	113	115	70
50-54	321	316	127	114	101	58
55-59	216	223	68	83	18	56
60-64	164	149	79	70	37	52
65 or Older	195	187	76	73	73	35

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.27D Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.32	0.34	0.18	0.17	0.12	0.11
12	0.70	0.65	0.47	0.50	0.29	0.33
13	0.77	0.80	0.61	0.60	0.40	0.39
14	0.79	0.88	0.70	0.73	0.45	0.50
15	0.88	0.88	0.77	0.72	0.55	0.49
16	0.98	0.92	0.82	0.74	0.52	0.51
17	1.04	1.03	0.88	0.91	0.62	0.60
18	1.06	1.12	0.94	0.92	0.65	0.65
19	1.26	1.26	1.06	1.07	0.75	0.74
20	1.23	1.30	1.04	1.11	0.72	0.66
21	1.24	1.32	0.96	1.19	0.63	0.76
22	1.29	1.22	0.96	1.01	0.65	0.75
23	1.19	1.19	0.92	0.92	0.62	0.62
24	1.25	1.18	1.07	0.80	0.76	0.63
25	1.27	1.28	0.88	0.90	0.57	0.57
26-29	1.19	1.22	0.83	0.85	0.56	0.56
30-34	1.02	1.00	0.64	0.65	0.45	0.46
35-39	1.09	1.09	0.61	0.64	0.42	0.37
40-44	1.04	1.11	0.56	0.63	0.40	0.38
45-49	1.10	1.07	0.67	0.53	0.54	0.32
50-54	1.60	1.63	0.63	0.59	0.50	0.30
55-59	1.45	1.36	0.45	0.51	0.12	0.34
60-64	1.40	1.19	0.68	0.55	0.31	0.41
65 or Older	0.57	0.54	0.22	0.21	0.21	0.10

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.27P Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.2179		0.1310		0.0716	
12	0.2742		0.5694		0.5978	
13	0.8957		0.9165		0.3310	
14	0.8291		0.4305		0.4013	
15	0.1634		0.6926		0.2812	
16	0.7679		0.3714		0.0812	
17	0.7223		0.2133		0.4533	
18	0.7432		0.4346		0.7618	
19	0.1297		0.3989		0.6541	
20	0.0708		0.0153		0.0270	
21	0.4676		0.0387		0.0396	
22	0.3108		0.0547		0.4627	
23	0.1154		0.0504		0.1316	
24	0.0790		0.0034		0.4032	
25	0.6022		0.9669		0.3418	
26-29	0.3448		0.5699		0.6307	
30-34	0.0679		0.8466		0.3455	
35-39	0.0827		0.5766		0.3500	
40-44	0.9025		0.5822		0.4549	
45-49	0.6133		0.1267		0.1266	
50-54	0.1107		0.2250		0.1045	
55-59	0.8740		0.5132		0.0158	
60-64	0.2948		0.5187		0.6188	
65 or Older	0.9155		0.8784		0.1602	

\*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.