

Table 7.70C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
ILLCIT DRUG²	138	131	13	15	136	129
Marijuana and Hashish	110	107	11	13	109	106
Cocaine	41	42	3	3	41	42
Crack	21	22	0	*	21	22
Heroin	8	9	1	1	8	8
Hallucinogens	27	24	3	2	26	23
LSD	7	5	0	1	7	5
PCP	4	4	1	1	4	3
Ecstasy	22	21	2	2	22	21
Inhalants	16	15	1	1	16	15
Nonmedical Use of Psychotherapeutics ³	83	82	7	8	83	82
Pain Relievers	70	70	7	8	70	70
OxyContin [®]	--	12	--	1	--	12
Tranquilizers	44	42	2	1	44	42
Stimulants	37	39	3	3	37	39
Methamphetamine	27	27	1	1	27	27
Sedatives	22	19	1	2	22	19
ILLCIT DRUG OTHER THAN MARIJUANA²	94	90	8	9	93	90

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.70D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
ILLCIT DRUG²	0.22	0.21	0.54	0.60	0.23	0.22
Marijuana and Hashish	0.18	0.17	0.45	0.51	0.19	0.18
Cocaine	0.07	0.07	0.13	0.11	0.07	0.07
Crack	0.03	0.04	0.02	*	0.03	0.04
Heroin	0.01	0.01	0.03	0.06	0.01	0.01
Hallucinogens	0.04	0.04	0.11	0.09	0.04	0.04
LSD	0.01	0.01	0.02	0.05	0.01	0.01
PCP	0.01	0.01	0.06	0.06	0.01	0.01
Ecstasy	0.04	0.03	0.08	0.07	0.04	0.03
Inhalants	0.03	0.02	0.03	0.05	0.03	0.02
Nonmedical Use of Psychotherapeutics ³	0.14	0.13	0.31	0.32	0.14	0.14
Pain Relievers	0.11	0.11	0.28	0.31	0.12	0.12
OxyContin [®]	--	0.04	--	0.08	--	0.04
Tranquilizers	0.07	0.07	0.08	0.05	0.07	0.07
Stimulants	0.06	0.06	0.13	0.13	0.06	0.07
Methamphetamine	0.04	0.04	0.04	0.04	0.05	0.05
Sedatives	0.04	0.03	0.06	0.07	0.04	0.03
ILLCIT DRUG OTHER THAN MARIJUANA²	0.15	0.15	0.31	0.34	0.16	0.15

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.70P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
ILLCIT DRUG²	0.3433		0.5599		0.3505	
Marijuana and Hashish	0.5673		0.8319		0.6115	
Cocaine	0.5022		0.8034		0.4677	
Crack	0.9295		*		0.9049	
Heroin	0.9793		0.3244		0.8707	
Hallucinogens	0.0066		0.6327		0.0088	
LSD	0.3376		0.3246		0.2480	
PCP	0.2568		0.2048		0.1898	
Ecstasy	0.0035		0.8452		0.0046	
Inhalants	0.8672		0.2530		0.7767	
Nonmedical Use of Psychotherapeutics ³	0.9655		0.6017		0.9552	
Pain Relievers	0.4268		0.1574		0.4848	
OxyContin [®]	--		--		--	
Tranquilizers	0.7936		0.0820		0.8618	
Stimulants	0.5738		0.6201		0.5710	
Methamphetamine	0.6840		0.3374		0.6539	
Sedatives	0.1400		0.3150		0.1237	
ILLCIT DRUG OTHER THAN MARIJUANA²	0.5005		0.4696		0.4979	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.71C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	138	131	13	15	136	129
AGE						
15-17	27	28	3	4	26	28
18-25	50	51	8	8	49	51
26-44	119	113	9	11	118	110
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	129	123	12	13	127	122
White	114	109	10	11	112	107
Black or African American	43	44	6	7	42	44
American Indian or Alaska Native	9	10	*	*	9	10
Native Hawaiian or Other Pacific Islander	7	*	*	*	7	*
Asian	22	17	*	*	22	17
Two or More Races	14	18	*	*	13	18
Hispanic or Latino	43	39	5	8	43	39
TRIMESTER²						
First	N/A	N/A	9	10	N/A	N/A
Second	N/A	N/A	6	9	N/A	N/A
Third	N/A	N/A	6	7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.71D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.22	0.21	0.54	0.60	0.23	0.22
AGE						
15-17	0.44	0.46	3.39	3.98	0.45	0.47
18-25	0.32	0.32	0.86	0.93	0.34	0.34
26-44	0.30	0.28	0.64	0.74	0.31	0.29
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.25	0.24	0.63	0.62	0.25	0.24
White	0.29	0.28	0.70	0.68	0.29	0.28
Black or African American	0.52	0.54	2.09	2.26	0.54	0.56
American Indian or Alaska Native	2.34	2.60	*	*	2.44	2.73
Native Hawaiian or Other Pacific Islander	3.89	*	*	*	4.01	*
Asian	0.72	0.54	*	*	0.76	0.57
Two or More Races	2.07	2.52	*	*	2.17	2.62
Hispanic or Latino	0.46	0.41	0.92	1.75	0.49	0.43
TRIMESTER²						
First	N/A	N/A	1.28	1.24	N/A	N/A
Second	N/A	N/A	0.61	0.99	N/A	N/A
Third	N/A	N/A	0.82	0.76	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.71P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.3433		0.5599		0.3505	
AGE						
15-17	0.3212		0.3004		0.2362	
18-25	0.1405		0.7376		0.1324	
26-44	0.6800		0.4301		0.6823	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4869		0.7636		0.5688	
White	0.3823		0.7947		0.4506	
Black or African American	0.8816		0.9061		0.9664	
American Indian or Alaska Native	0.4160		*		0.4612	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1746		*		0.1715	
Two or More Races	0.0306		*		0.0399	
Hispanic or Latino	0.4755		0.1921		0.3045	
TRIMESTER²						
First	N/A		0.7492		N/A	
Second	N/A		0.5164		N/A	
Third	N/A		0.8861		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.72C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	110	107	11	13	109	106
AGE						
15-17	24	24	3	3	24	24
18-25	46	45	7	7	45	44
26-44	91	91	7	10	91	89
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	105	104	10	10	104	104
White	92	92	9	9	91	91
Black or African American	38	40	4	4	37	40
American Indian or Alaska Native	8	9	*	*	8	9
Native Hawaiian or Other Pacific Islander	3	3	*	*	3	3
Asian	14	10	*	*	13	10
Two or More Races	12	14	*	*	12	13
Hispanic or Latino	31	30	4	8	31	29
TRIMESTER²						
First	N/A	N/A	7	7	N/A	N/A
Second	N/A	N/A	5	8	N/A	N/A
Third	N/A	N/A	6	6	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.72D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.18	0.17	0.45	0.51	0.19	0.18
AGE						
15-17	0.40	0.40	2.91	3.43	0.40	0.41
18-25	0.30	0.28	0.76	0.80	0.31	0.30
26-44	0.23	0.23	0.52	0.66	0.24	0.23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.20	0.52	0.50	0.21	0.21
White	0.23	0.23	0.63	0.60	0.24	0.24
Black or African American	0.46	0.49	1.38	1.24	0.47	0.51
American Indian or Alaska Native	2.02	2.48	*	*	2.15	2.60
Native Hawaiian or Other Pacific Islander	1.45	1.84	*	*	1.50	1.88
Asian	0.45	0.33	*	*	0.47	0.34
Two or More Races	1.82	1.88	*	*	1.92	1.95
Hispanic or Latino	0.34	0.31	0.81	1.63	0.35	0.32
TRIMESTER²						
First	N/A	N/A	1.05	0.95	N/A	N/A
Second	N/A	N/A	0.53	0.94	N/A	N/A
Third	N/A	N/A	0.81	0.73	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.72P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.5673		0.8319		0.6115	
AGE						
15-17	0.7234		0.3847		0.6739	
18-25	0.0057		0.7750		0.0065	
26-44	0.5707		0.4768		0.5860	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6934		0.5234		0.8218	
White	0.5658		0.6741		0.6573	
Black or African American	0.8254		0.5084		0.6687	
American Indian or Alaska Native	0.2081		*		0.2580	
Native Hawaiian or Other Pacific Islander	0.8157		*		0.8868	
Asian	0.4010		*		0.4100	
Two or More Races	0.2534		*		0.3094	
Hispanic or Latino	0.6568		0.3049		0.4504	
TRIMESTER²						
First	N/A		0.6090		N/A	
Second	N/A		0.4426		N/A	
Third	N/A		0.8567		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.73C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOBACCO PRODUCT²	232	232	29	32	227	228
Cigarettes	229	231	29	31	225	226
Smokeless Tobacco	22	18	3	1	22	18
Cigars	66	65	7	7	64	63
Pipe Tobacco	19	20	4	4	19	20
ALCOHOL	237	238	23	28	231	232
Binge Alcohol Use ³	192	188	14	16	191	185
Heavy Alcohol Use ³	94	90	5	4	94	90

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.73D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOBACCO PRODUCT²	0.38	0.38	1.21	1.24	0.39	0.39
Cigarettes	0.37	0.37	1.18	1.22	0.38	0.38
Smokeless Tobacco	0.04	0.03	0.14	0.04	0.04	0.03
Cigars	0.11	0.11	0.28	0.27	0.11	0.11
Pipe Tobacco	0.03	0.03	0.16	0.15	0.03	0.03
ALCOHOL	0.38	0.38	0.94	1.12	0.39	0.39
Binge Alcohol Use ³	0.31	0.30	0.58	0.63	0.32	0.31
Heavy Alcohol Use ³	0.15	0.15	0.22	0.15	0.16	0.15

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.73P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOBACCO PRODUCT²	0.0466		0.8749		0.0554	
Cigarettes	0.0335		0.9652		0.0372	
Smokeless Tobacco	0.2844		0.2291		0.3695	
Cigars	0.4283		0.7048		0.4151	
Pipe Tobacco	0.9629		0.8019		0.9688	
ALCOHOL	0.5332		0.1936		0.6571	
Binge Alcohol Use ³	0.9282		0.4165		0.8469	
Heavy Alcohol Use ³	0.1729		0.3598		0.1359	

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.74C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	229	231	29	31	225	226
AGE						
15-17	30	31	4	4	29	30
18-25	71	69	15	15	69	67
26-44	209	206	22	25	205	203
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	213	212	27	30	209	209
White	189	187	26	27	185	183
Black or African American	78	78	5	7	78	78
American Indian or Alaska Native	15	15	*	*	14	15
Native Hawaiian or Other Pacific Islander	10	*	*	*	10	*
Asian	43	26	*	*	43	25
Two or More Races	20	21	*	*	20	21
Hispanic or Latino	75	73	7	9	74	73
TRIMESTER²						
First	N/A	N/A	15	17	N/A	N/A
Second	N/A	N/A	17	13	N/A	N/A
Third	N/A	N/A	17	19	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.74D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.37	0.37	1.18	1.22	0.38	0.38
AGE						
15-17	0.49	0.51	4.11	4.44	0.49	0.51
18-25	0.45	0.43	1.58	1.66	0.47	0.45
26-44	0.52	0.52	1.57	1.63	0.53	0.53
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.41	1.42	1.43	0.42	0.42
White	0.47	0.47	1.80	1.76	0.48	0.48
Black or African American	0.94	0.95	1.67	2.49	0.98	0.99
American Indian or Alaska Native	3.83	4.12	*	*	3.97	4.18
Native Hawaiian or Other Pacific Islander	5.41	*	*	*	5.55	*
Asian	1.41	0.82	*	*	1.50	0.85
Two or More Races	3.10	2.92	*	*	3.21	3.01
Hispanic or Latino	0.80	0.77	1.31	1.87	0.85	0.81
TRIMESTER²						
First	N/A	N/A	2.15	2.26	N/A	N/A
Second	N/A	N/A	1.75	1.46	N/A	N/A
Third	N/A	N/A	2.23	2.22	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.74P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.0335		0.9652		0.0372	
AGE						
15-17	0.0340		0.9862		0.0302	
18-25	0.0690		0.8942		0.0492	
26-44	0.1522		0.5651		0.1762	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0321		0.6845		0.0422	
White	0.0767		0.4308		0.1112	
Black or African American	0.9927		0.0507		0.9442	
American Indian or Alaska Native	0.8756		*		0.9523	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0081		*		0.0090	
Two or More Races	0.2022		*		0.1985	
Hispanic or Latino	0.9818		0.4766		0.9043	
TRIMESTER²						
First	N/A		0.7573		N/A	
Second	N/A		0.1846		N/A	
Third	N/A		0.7167		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.75C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	237	238	23	28	231	232
AGE						
15-17	35	36	3	3	35	35
18-25	81	80	10	10	79	77
26-44	216	217	20	27	211	212
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	221	218	21	24	214	212
White	183	184	18	21	176	180
Black or African American	83	81	4	6	83	80
American Indian or Alaska Native	14	14	*	*	14	14
Native Hawaiian or Other Pacific Islander	11	*	*	*	11	*
Asian	66	61	*	*	65	59
Two or More Races	22	21	*	*	22	21
Hispanic or Latino	98	88	10	15	98	85
TRIMESTER²						
First	N/A	N/A	17	22	N/A	N/A
Second	N/A	N/A	12	13	N/A	N/A
Third	N/A	N/A	10	11	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.75D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.38	0.38	0.94	1.12	0.39	0.39
AGE						
15-17	0.58	0.58	3.53	3.68	0.59	0.59
18-25	0.52	0.50	1.05	1.07	0.54	0.52
26-44	0.54	0.55	1.43	1.72	0.55	0.56
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.42	1.07	1.15	0.43	0.42
White	0.46	0.47	1.26	1.34	0.46	0.47
Black or African American	1.01	0.98	1.39	2.06	1.05	1.01
American Indian or Alaska Native	3.64	3.79	*	*	3.81	3.90
Native Hawaiian or Other Pacific Islander	5.63	*	*	*	5.81	*
Asian	2.17	1.92	*	*	2.27	1.98
Two or More Races	3.37	2.94	*	*	3.50	3.04
Hispanic or Latino	1.05	0.93	1.88	3.21	1.12	0.95
TRIMESTER²						
First	N/A	N/A	2.36	2.83	N/A	N/A
Second	N/A	N/A	1.25	1.44	N/A	N/A
Third	N/A	N/A	1.39	1.29	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.75P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.5332		0.1936		0.6571	
AGE						
15-17	0.8762		0.8971		0.7980	
18-25	0.7206		0.9290		0.9371	
26-44	0.4558		0.1575		0.6467	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6909		0.3332		0.9629	
White	0.4172		0.4935		0.2261	
Black or African American	0.1574		0.0940		0.1518	
American Indian or Alaska Native	0.9500		*		0.8094	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1234		*		0.2013	
Two or More Races	0.1634		*		0.2363	
Hispanic or Latino	0.7512		0.3797		0.5616	
TRIMESTER²						
First	N/A		0.2664		N/A	
Second	N/A		0.5111		N/A	
Third	N/A		0.8783		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.76C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	192	188	14	16	191	185
AGE						
15-17	30	31	2	3	30	31
18-25	73	70	6	6	72	68
26-44	168	171	12	15	167	169
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	182	179	13	14	181	176
White	153	154	10	11	152	151
Black or African American	71	63	2	5	69	62
American Indian or Alaska Native	15	14	*	*	14	13
Native Hawaiian or Other Pacific Islander	*	7	*	*	*	7
Asian	33	33	*	*	32	31
Two or More Races	16	19	*	*	16	19
Hispanic or Latino	70	68	3	7	70	68
TRIMESTER²						
First	N/A	N/A	12	13	N/A	N/A
Second	N/A	N/A	5	5	N/A	N/A
Third	N/A	N/A	4	5	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.76D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.31	0.30	0.58	0.63	0.32	0.31
AGE						
15-17	0.49	0.51	2.32	2.86	0.50	0.51
18-25	0.47	0.44	0.66	0.70	0.49	0.46
26-44	0.42	0.43	0.88	0.94	0.43	0.44
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.35	0.34	0.70	0.70	0.36	0.35
White	0.39	0.39	0.74	0.69	0.40	0.40
Black or African American	0.86	0.76	0.81	1.75	0.88	0.79
American Indian or Alaska Native	3.75	3.70	*	*	3.96	3.80
Native Hawaiian or Other Pacific Islander	*	3.90	*	*	*	4.00
Asian	1.09	1.03	*	*	1.11	1.02
Two or More Races	2.47	2.64	*	*	2.60	2.75
Hispanic or Latino	0.75	0.71	0.66	1.46	0.80	0.76
TRIMESTER²						
First	N/A	N/A	1.75	1.74	N/A	N/A
Second	N/A	N/A	0.50	0.58	N/A	N/A
Third	N/A	N/A	0.49	0.59	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.76P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.9282		0.4165		0.8469	
AGE						
15-17	0.1557		0.4320		0.2027	
18-25	0.4978		0.6823		0.6150	
26-44	0.5496		0.4829		0.6625	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9930		0.6105		0.8638	
White	0.5011		0.9634		0.3498	
Black or African American	0.1638		0.1092		0.1376	
American Indian or Alaska Native	0.4372		*		0.5056	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9615		*		0.9122	
Two or More Races	0.2759		*		0.3286	
Hispanic or Latino	0.7071		0.5581		0.8059	
TRIMESTER²						
First	N/A		0.8319		N/A	
Second	N/A		0.4091		N/A	
Third	N/A		0.3409		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.77C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	94	90	5	4	94	90
AGE						
15-17	13	15	0	*	13	15
18-25	42	43	2	2	41	43
26-44	78	75	5	3	78	75
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	90	89	5	4	89	89
White	80	79	5	3	79	78
Black or African American	31	29	1	*	31	29
American Indian or Alaska Native	8	8	*	*	7	7
Native Hawaiian or Other Pacific Islander	4	3	*	*	4	3
Asian	22	15	*	*	22	15
Two or More Races	8	8	*	*	8	8
Hispanic or Latino	24	19	1	1	23	19
TRIMESTER²						
First	N/A	N/A	4	4	N/A	N/A
Second	N/A	N/A	*	*	N/A	N/A
Third	N/A	N/A	1	1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.77D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.15	0.15	0.22	0.15	0.16	0.15
AGE						
15-17	0.22	0.24	0.09	*	0.22	0.25
18-25	0.27	0.27	0.26	0.27	0.28	0.29
26-44	0.20	0.19	0.35	0.19	0.20	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.17	0.17	0.28	0.19	0.18	0.18
White	0.20	0.20	0.34	0.22	0.21	0.21
Black or African American	0.37	0.35	0.26	*	0.39	0.37
American Indian or Alaska Native	2.05	2.09	*	*	1.95	1.99
Native Hawaiian or Other Pacific Islander	1.90	1.58	*	*	1.93	1.58
Asian	0.71	0.49	*	*	0.75	0.52
Two or More Races	1.16	1.16	*	*	1.22	1.21
Hispanic or Latino	0.25	0.20	0.19	0.21	0.27	0.21
TRIMESTER²						
First	N/A	N/A	0.54	0.51	N/A	N/A
Second	N/A	N/A	*	*	N/A	N/A
Third	N/A	N/A	0.08	0.07	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 7.77P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL	0.1729		0.3598		0.1359	
AGE						
15-17	0.2957		*		0.3135	
18-25	0.7460		0.9967		0.7944	
26-44	0.2806		0.2336		0.2248	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1138		0.3301		0.0820	
White	0.0414		0.6199		0.0291	
Black or African American	0.7238		*		0.8271	
American Indian or Alaska Native	0.4767		*		0.5355	
Native Hawaiian or Other Pacific Islander	0.2582		*		0.2419	
Asian	0.7150		*		0.7192	
Two or More Races	0.7545		*		0.7360	
Hispanic or Latino	0.6846		0.5270		0.6677	
TRIMESTER²						
First	N/A		0.7795		N/A	
Second	N/A		*		N/A	
Third	N/A		0.7859		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.