

**Table 6.10C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Numbers in Thousands, 2003 and 2004**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2003	2004	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	481	922	99	156	459	891	112	200	458	888
Marijuana	481	922	61	95	471	910	76	137	466	904
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	481	922	78	135	467	900	89	159	467	901
<b>ALCOHOL</b>	481	922	130	219	452	902	173	254	440	893
<b>ILLCIT DRUG OR ALCOHOL<sup>2</sup></b>	481	922	156	257	435	885	189	298	427	877
<b>BOTH ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	481	922	45	*	475	909	82	137	470	906

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.10D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Percentages, 2003 and 2004**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2003	2004	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLICIT DRUG<sup>2</sup></b>	0.20	0.41	2.16	3.30	0.20	0.40	1.66	2.82	0.20	0.40
Marijuana	0.20	0.41	2.41	4.01	0.20	0.40	1.93	3.39	0.20	0.40
<b>ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.20	0.41	3.15	4.68	0.20	0.40	2.46	4.00	0.20	0.40
<b>ALCOHOL</b>	0.20	0.41	1.57	2.48	0.20	0.41	0.91	1.41	0.20	0.43
<b>ILLICIT DRUG OR ALCOHOL<sup>2</sup></b>	0.20	0.41	1.33	2.10	0.20	0.41	0.83	1.37	0.20	0.43
<b>BOTH ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	0.20	0.41	3.90	*	0.20	0.40	2.50	4.08	0.20	0.40

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.10P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2003	2004	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	0.1052		0.5767		0.1325		0.3394		0.1566	
Marijuana	0.1052		0.4218		0.1059		0.2999		0.1215	
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.1052		0.7786		0.1419		0.7570		0.1325	
<b>ALCOHOL</b>	0.1052		0.2619		0.2001		0.7469		0.1296	
<b>ILLCIT DRUG OR ALCOHOL<sup>2</sup></b>	0.1052		0.3699		0.1770		0.4219		0.1703	
<b>BOTH ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	0.1052		*		0.1518		0.8789		0.1196	

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.11C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2003 and 2004**

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	205	356	112	200	169	296
Marijuana	138	210	76	137	114	162
Illicit Drug Other Than Marijuana <sup>3</sup>	161	295	89	159	131	247
Alcohol	397	562	173	254	344	530
Illicit Drug or Alcohol <sup>3</sup>	428	615	189	298	368	564
Both Illicit Drug and Alcohol <sup>3</sup>	134	255	82	137	103	223
<b>DEPENDENCE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	159	281	99	156	128	234
Marijuana	101	151	61	95	80	118
Illicit Drug Other Than Marijuana <sup>3</sup>	131	240	78	135	103	199
Alcohol	269	383	130	219	230	334
Illicit Drug or Alcohol <sup>3</sup>	303	448	156	257	254	379
Both Illicit Drug and Alcohol <sup>3</sup>	74	157	45	96	58	128

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.11D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2003 and 2004**

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	0.09	0.16	0.55	0.91	0.09	0.15
Marijuana	0.06	0.10	0.37	0.63	0.06	0.08
Illicit Drug Other Than Marijuana <sup>3</sup>	0.08	0.13	0.44	0.73	0.07	0.13
Alcohol	0.18	0.26	0.81	1.22	0.17	0.27
Illicit Drug or Alcohol <sup>3</sup>	0.19	0.28	0.86	1.41	0.18	0.29
Both Illicit Drug and Alcohol <sup>3</sup>	0.06	0.12	0.41	0.63	0.05	0.11
<b>DEPENDENCE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	0.07	0.13	0.48	0.71	0.07	0.12
Marijuana	0.05	0.07	0.31	0.44	0.04	0.06
Illicit Drug Other Than Marijuana <sup>3</sup>	0.06	0.11	0.38	0.61	0.05	0.10
Alcohol	0.12	0.18	0.62	1.03	0.12	0.17
Illicit Drug or Alcohol <sup>3</sup>	0.14	0.20	0.71	1.20	0.13	0.19
Both Illicit Drug and Alcohol <sup>3</sup>	0.03	0.07	0.23	0.44	0.03	0.07

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.11P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	0.4787		0.5749		0.8827	
Marijuana	0.5839		0.9410		0.2927	
Illicit Drug Other Than Marijuana <sup>3</sup>	0.2966		0.6388		0.4801	
Alcohol	0.4996		0.7930		0.5433	
Illicit Drug or Alcohol <sup>3</sup>	0.6522		0.9771		0.8587	
Both Illicit Drug and Alcohol <sup>3</sup>	0.1631		0.7509		0.2070	
<b>DEPENDENCE<sup>2</sup></b>						
Illicit Drug <sup>3</sup>	0.4132		0.6901		0.6912	
Marijuana	0.2826		0.6866		0.1792	
Illicit Drug Other Than Marijuana <sup>3</sup>	0.1048		0.3163		0.2799	
Alcohol	0.2460		0.3628		0.6546	
Illicit Drug or Alcohol <sup>3</sup>	0.4448		0.7074		0.8059	
Both Illicit Drug and Alcohol <sup>3</sup>	0.0243		0.0570		0.2237	

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>3</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.12C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004**

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	481	922	112	200	458	888
18-25	117	181	45	83	104	160
26-49	308	502	95	180	294	461
50 or Older	320	744	*	*	317	742
<b>MALE</b>	282	639	72	179	273	615
18-25	68	113	29	60	59	94
26-49	184	346	65	*	179	313
50 or Older	202	525	*	*	202	523
<b>FEMALE</b>	374	646	79	90	357	638
18-25	85	145	30	55	80	134
26-49	246	354	67	*	236	349
50 or Older	247	530	*	*	245	530

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.12D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2003 and 2004**

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.20	0.41	1.66	2.82	0.20	0.40
18-25	0.32	0.51	1.46	2.48	0.31	0.50
26-49	0.28	0.48	2.76	5.06	0.28	0.46
50 or Older	0.38	0.86	*	*	0.38	0.86
<b>MALE</b>	0.26	0.60	1.82	3.64	0.26	0.60
18-25	0.40	0.65	1.60	2.97	0.39	0.62
26-49	0.36	0.68	3.15	*	0.36	0.64
50 or Older	0.52	1.35	*	*	0.53	1.35
<b>FEMALE</b>	0.31	0.54	2.78	4.31	0.30	0.54
18-25	0.45	0.78	2.40	4.12	0.45	0.78
26-49	0.44	0.66	4.68	*	0.43	0.66
50 or Older	0.54	1.11	*	*	0.54	1.11

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).



**Table 6.12P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>	
			Yes	No
	2003	2004	2003	2004
<b>TOTAL</b>	0.1052		0.3394	0.1566
18-25	0.7137		0.2358	0.3464
26-49	0.9705		0.6608	0.8198
50 or Older	0.0279		*	0.0318
<b>MALE</b>	0.1422		0.0463	0.3530
18-25	0.5455		0.6223	0.2620
26-49	0.1952		*	0.5180
50 or Older	0.2804		*	0.3439
<b>FEMALE</b>	0.4026		0.4088	0.2952
18-25	0.9670		0.0944	0.7062
26-49	0.2171		*	0.3696
50 or Older	0.0440		*	0.0381

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.13C Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	205	356	112	200	169	296
18-25	79	139	45	83	64	107
26-49	173	312	95	180	142	258
50 or Older	61	113	28	44	55	106
<b>MALE</b>	166	316	72	179	147	257
18-25	64	113	29	60	55	95
26-49	141	284	65	165	122	231
50 or Older	55	54	*	*	54	35
<b>FEMALE</b>	116	184	79	90	83	157
18-25	48	71	30	55	36	47
26-49	101	139	67	72	74	116
50 or Older	26	101	24	16	11	100

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.13D Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Percentages, 2003 and 2004**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.09	0.16	0.55	0.91	0.09	0.15
18-25	0.23	0.40	0.89	1.67	0.22	0.36
26-49	0.17	0.30	0.87	1.60	0.16	0.28
50 or Older	0.08	0.14	0.57	0.68	0.07	0.14
<b>MALE</b>	0.16	0.29	1.04	2.24	0.15	0.26
18-25	0.37	0.63	1.48	3.06	0.37	0.60
26-49	0.28	0.56	1.82	3.79	0.27	0.49
50 or Older	0.15	0.14	*	*	0.15	0.10
<b>FEMALE</b>	0.10	0.17	0.59	0.68	0.08	0.16
18-25	0.29	0.44	1.04	1.84	0.26	0.36
26-49	0.20	0.27	0.92	1.09	0.17	0.26
50 or Older	0.06	0.22	0.80	0.39	0.03	0.24

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.13P Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>	
			Yes	No
	2003	2004	2003	2004
<b>TOTAL</b>	0.4787		0.5749	0.8827
18-25	0.5997		0.1447	0.7882
26-49	0.7066		0.5470	0.9646
50 or Older	0.4513		0.6178	0.6710
<b>MALE</b>	0.2103		0.1213	0.8498
18-25	0.3130		0.1916	0.5488
26-49	0.2273		0.1581	0.6976
50 or Older	0.5750		*	0.0816
<b>FEMALE</b>	0.4319		0.1051	0.9758
18-25	0.6020		0.3870	0.0995
26-49	0.1089		0.1356	0.5128
50 or Older	0.1828		0.3141	0.1064

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.14C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	481	922	173	254	440	893
18-25	117	181	58	112	96	139
26-49	308	502	132	224	286	451
50 or Older	320	744	95	*	305	738
<b>MALE</b>	282	639	126	215	253	597
18-25	68	113	39	72	56	85
26-49	184	346	90	185	161	287
50 or Older	202	525	80	*	186	520
<b>FEMALE</b>	374	646	121	156	344	631
18-25	85	145	40	84	72	119
26-49	246	354	98	116	225	340
50 or Older	247	530	*	*	241	526

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.14D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2003 and 2004**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.20	0.41	0.91	1.41	0.20	0.43
18-25	0.32	0.51	0.94	1.59	0.32	0.49
26-49	0.28	0.48	1.31	2.36	0.29	0.47
50 or Older	0.38	0.86	3.75	*	0.38	0.88
<b>MALE</b>	0.26	0.60	1.02	1.74	0.26	0.63
18-25	0.40	0.65	1.03	1.71	0.42	0.66
26-49	0.36	0.68	1.33	2.77	0.36	0.64
50 or Older	0.52	1.35	4.31	*	0.51	1.40
<b>FEMALE</b>	0.31	0.54	1.84	2.53	0.30	0.55
18-25	0.45	0.78	1.60	3.07	0.46	0.78
26-49	0.44	0.66	2.92	3.98	0.43	0.67
50 or Older	0.54	1.11	*	*	0.54	1.11

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.14P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>	
			Yes	No
	2003	2004	2003	2004
<b>TOTAL</b>	0.1052		0.7469	0.1296
18-25	0.7137		0.7362	0.3965
26-49	0.9705		0.7765	0.8168
50 or Older	0.0279		*	0.0247
<b>MALE</b>	0.1422		0.7646	0.1706
18-25	0.5455		0.7334	0.4607
26-49	0.1952		0.4946	0.3297
50 or Older	0.2804		*	0.2362
<b>FEMALE</b>	0.4026		0.6394	0.4152
18-25	0.9670		0.2790	0.5919
26-49	0.2171		0.9294	0.2465
50 or Older	0.0440		*	0.0456

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.15C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	397	562	173	254	344	530
18-25	132	228	58	112	116	187
26-49	297	445	132	224	254	419
50 or Older	209	322	95	98	188	306
<b>MALE</b>	328	496	126	215	294	481
18-25	99	173	39	72	91	155
26-49	251	392	90	185	222	383
50 or Older	178	273	80	*	161	264
<b>FEMALE</b>	205	289	121	156	162	239
18-25	77	126	40	84	61	90
26-49	152	217	98	116	120	180
50 or Older	98	164	51	68	83	150

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).



**Table 6.15D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Standard Errors of Percentages, 2003 and 2004**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>			
			Yes		No	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.18	0.26	0.81	1.22	0.17	0.27
18-25	0.34	0.59	1.08	1.99	0.36	0.57
26-49	0.28	0.43	1.17	1.93	0.27	0.45
50 or Older	0.25	0.38	1.92	1.54	0.24	0.39
<b>MALE</b>	0.31	0.45	1.66	2.70	0.30	0.48
18-25	0.55	0.91	1.90	3.33	0.56	0.92
26-49	0.47	0.76	2.30	3.91	0.46	0.79
50 or Older	0.46	0.70	4.20	*	0.45	0.72
<b>FEMALE</b>	0.18	0.26	0.88	1.18	0.17	0.25
18-25	0.41	0.68	1.25	2.54	0.41	0.62
26-49	0.30	0.42	1.28	1.73	0.27	0.40
50 or Older	0.22	0.36	1.68	1.62	0.20	0.37

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.15P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS <sup>1</sup>	
			Yes	No
	2003	2004	2003	2004
<b>TOTAL</b>	0.4996		0.7930	0.5433
18-25	0.2366		0.2536	0.4151
26-49	0.6158		0.5945	0.7768
50 or Older	0.9567		0.3318	0.8989
<b>MALE</b>	0.2608		0.7612	0.2939
18-25	0.1363		0.4741	0.1569
26-49	0.2840		0.6499	0.4397
50 or Older	0.8227		*	0.9876
<b>FEMALE</b>	0.5942		0.7560	0.5228
18-25	0.9821		0.3339	0.4658
26-49	0.3902		0.8856	0.4466
50 or Older	0.7989		0.8660	0.8562

\*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).