

Table 2.103C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	160	154	165	154	154	148
GENDER						
Male	101	108	109	108	102	106
Female	109	100	113	101	105	94
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	146	136	150	138	143	134
White	121	115	124	121	116	116
Black or African American	55	59	53	55	46	45
American Indian or Alaska Native	10	10	9	9	7	7
Native Hawaiian or Other Pacific Islander	8	*	*	*	*	*
Asian	39	33	34	31	29	24
Two or More Races	15	15	15	16	14	13
Hispanic or Latino	61	65	62	62	52	61
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	73	81	76	82	78	81
Female, White, Not Hispanic	82	76	83	79	78	74
Male, Black, Not Hispanic	38	43	37	40	32	31
Female, Black, Not Hispanic	40	36	36	35	29	30
Male, Hispanic	48	50	49	48	37	46
Female, Hispanic	44	44	48	43	38	41

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.103D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.43	0.41	0.44	0.41	0.41	0.39
GENDER						
Male	0.53	0.56	0.57	0.56	0.53	0.55
Female	0.59	0.55	0.62	0.55	0.57	0.51
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.47	0.43	0.48	0.44	0.46	0.43
White	0.52	0.49	0.53	0.52	0.50	0.50
Black or African American	1.00	1.06	0.96	0.98	0.84	0.81
American Indian or Alaska Native	3.99	4.44	3.77	3.81	2.94	2.92
Native Hawaiian or Other Pacific Islander	5.81	*	*	*	*	*
Asian	2.53	2.15	2.24	2.01	1.93	1.57
Two or More Races	2.80	2.59	2.79	2.75	2.55	2.28
Hispanic or Latino	0.98	1.02	1.01	0.98	0.84	0.97
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.61	0.68	0.64	0.68	0.66	0.68
Female, White, Not Hispanic	0.71	0.67	0.73	0.70	0.68	0.65
Male, Black, Not Hispanic	1.35	1.50	1.34	1.38	1.14	1.10
Female, Black, Not Hispanic	1.45	1.33	1.32	1.28	1.05	1.10
Male, Hispanic	1.50	1.52	1.51	1.48	1.16	1.41
Female, Hispanic	1.48	1.44	1.62	1.40	1.29	1.35

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.103P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.1120		0.6875		0.6311	
GENDER						
Male	0.9325		0.3781		0.7321	
Female	0.0180		0.1362		0.6987	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2461		0.9972		0.4246	
White	0.6987		0.7660		0.3736	
Black or African American	0.0963		0.6876		0.4690	
American Indian or Alaska Native	0.5083		0.6187		0.6685	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.5152		0.9600		0.5010	
Two or More Races	0.3893		0.7972		0.7079	
Hispanic or Latino	0.1712		0.3170		0.4452	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.8405		0.2436		0.7970	
Female, White, Not Hispanic	0.4536		0.5202		0.2583	
Male, Black, Not Hispanic	0.4764		0.6193		0.5940	
Female, Black, Not Hispanic	0.0877		0.2622		0.6092	
Male, Hispanic	0.4921		0.2617		0.6062	
Female, Hispanic	0.2290		0.9249		0.0933	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.104C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	569	679	874	875	899	943
GENDER						
Male	342	406	550	557	582	612
Female	461	494	632	630	589	659
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	530	627	846	838	852	888
White	417	481	721	714	759	777
Black or African American	236	254	282	279	274	300
American Indian or Alaska Native	*	33	48	*	46	63
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	177	256	221	267	223	229
Two or More Races	69	30	73	58	72	65
Hispanic or Latino	243	280	306	318	307	284
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	232	290	461	440	500	490
Female, White, Not Hispanic	350	364	495	505	478	555
Male, Black, Not Hispanic	115	163	197	187	202	190
Female, Black, Not Hispanic	178	193	202	219	170	206
Male, Hispanic	161	163	221	224	230	215
Female, Hispanic	188	192	195	200	182	193

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.104D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.28	0.33	0.44	0.43	0.45	0.46
GENDER						
Male	0.36	0.42	0.57	0.57	0.61	0.63
Female	0.44	0.47	0.61	0.60	0.56	0.62
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.30	0.35	0.48	0.47	0.48	0.50
White	0.29	0.33	0.50	0.50	0.53	0.54
Black or African American	1.07	1.13	1.28	1.24	1.25	1.34
American Indian or Alaska Native	*	3.13	4.79	*	4.61	5.89
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.15	3.05	2.68	3.18	2.70	2.73
Two or More Races	3.37	1.54	3.60	2.94	3.54	3.32
Hispanic or Latino	1.03	1.14	1.29	1.29	1.29	1.15
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.34	0.42	0.67	0.64	0.73	0.71
Female, White, Not Hispanic	0.47	0.49	0.67	0.68	0.64	0.74
Male, Black, Not Hispanic	1.20	1.66	2.05	1.90	2.10	1.94
Female, Black, Not Hispanic	1.45	1.53	1.64	1.74	1.38	1.64
Male, Hispanic	1.33	1.29	1.83	1.78	1.90	1.70
Female, Hispanic	1.62	1.59	1.68	1.66	1.56	1.60

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.104P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.1183		0.7763		0.6426	
GENDER						
Male	0.2687		0.7813		0.6086	
Female	0.2274		0.8922		0.2240	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2481		0.4605		0.3536	
White	0.9647		0.1407		0.1386	
Black or African American	0.0248		0.2467		0.4956	
American Indian or Alaska Native	*		*		0.3465	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4895		0.8919		0.5418	
Two or More Races	0.0819		0.0504		0.0221	
Hispanic or Latino	0.3221		0.3004		0.2851	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4798		0.1046		0.5834	
Female, White, Not Hispanic	0.6126		0.6198		0.0980	
Male, Black, Not Hispanic	0.0621		0.3629		0.2179	
Female, Black, Not Hispanic	0.1260		0.4733		0.7831	
Male, Hispanic	0.7707		0.0184		0.1507	
Female, Hispanic	0.2459		0.4691		0.9293	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.105C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	154	148	135	138	75	92
GENDER						
Male	102	106	93	100	56	66
Female	105	94	82	82	42	48
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	143	134	122	123	71	86
White	116	116	106	109	65	76
Black or African American	46	45	35	32	15	17
American Indian or Alaska Native	7	7	7	5	3	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	4	1
Asian	29	24	20	21	12	9
Two or More Races	14	13	11	12	4	9
Hispanic or Latino	52	61	49	56	25	31
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	78	81	71	77	49	54
Female, White, Not Hispanic	78	74	66	66	37	42
Male, Black, Not Hispanic	32	31	26	27	14	16
Female, Black, Not Hispanic	29	30	18	19	5	4
Male, Hispanic	37	46	37	43	22	26
Female, Hispanic	38	41	30	36	12	18

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.105D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.41	0.39	0.36	0.36	0.20	0.24
GENDER						
Male	0.53	0.55	0.49	0.52	0.29	0.34
Female	0.57	0.51	0.44	0.44	0.23	0.26
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.46	0.43	0.39	0.39	0.23	0.27
White	0.50	0.50	0.45	0.47	0.28	0.33
Black or African American	0.84	0.81	0.63	0.58	0.27	0.30
American Indian or Alaska Native	2.94	2.92	2.85	2.31	1.19	1.75
Native Hawaiian or Other Pacific Islander	*	*	*	*	2.75	0.65
Asian	1.93	1.57	1.30	1.35	0.79	0.60
Two or More Races	2.55	2.28	2.04	2.00	0.76	1.49
Hispanic or Latino	0.84	0.97	0.80	0.88	0.41	0.49
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.66	0.68	0.60	0.64	0.41	0.45
Female, White, Not Hispanic	0.68	0.65	0.57	0.58	0.32	0.37
Male, Black, Not Hispanic	1.14	1.10	0.95	0.94	0.49	0.55
Female, Black, Not Hispanic	1.05	1.10	0.66	0.71	0.18	0.16
Male, Hispanic	1.16	1.41	1.15	1.30	0.69	0.78
Female, Hispanic	1.29	1.35	1.01	1.16	0.42	0.58

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.105P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.6311		0.3790		0.5649	
GENDER						
Male	0.7321		0.6162		0.4605	
Female	0.6987		0.4014		0.9935	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4246		0.9306		0.7830	
White	0.3736		0.9184		0.8331	
Black or African American	0.4690		0.3167		0.3197	
American Indian or Alaska Native	0.6685		0.6053		0.7491	
Native Hawaiian or Other Pacific Islander	*		*		0.1094	
Asian	0.5010		0.4184		0.1548	
Two or More Races	0.7079		0.5940		0.0450	
Hispanic or Latino	0.4452		0.0324		0.3122	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.7970		0.7644		0.7257	
Female, White, Not Hispanic	0.2583		0.7970		0.8642	
Male, Black, Not Hispanic	0.5940		0.6148		0.2341	
Female, Black, Not Hispanic	0.6092		0.3194		0.6078	
Male, Hispanic	0.6062		0.6613		0.6023	
Female, Hispanic	0.0933		0.0042		0.2963	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.106C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	899	943	661	688	366	371
GENDER						
Male	582	612	508	504	316	313
Female	589	659	365	384	163	168
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	852	888	629	647	342	350
White	759	777	556	584	309	335
Black or African American	274	300	228	227	133	107
American Indian or Alaska Native	46	63	42	53	25	33
Native Hawaiian or Other Pacific Islander	*	*	*	*	13	12
Asian	223	229	100	111	38	52
Two or More Races	72	65	51	59	28	28
Hispanic or Latino	307	284	236	239	121	118
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	500	490	420	434	272	277
Female, White, Not Hispanic	478	555	316	323	140	163
Male, Black, Not Hispanic	202	190	182	160	111	94
Female, Black, Not Hispanic	170	206	113	140	59	51
Male, Hispanic	230	215	203	200	117	105
Female, Hispanic	182	193	117	125	25	43

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.106D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.45	0.46	0.33	0.34	0.18	0.18
GENDER						
Male	0.61	0.63	0.53	0.52	0.33	0.32
Female	0.56	0.62	0.35	0.36	0.16	0.16
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.50	0.36	0.36	0.19	0.20
White	0.53	0.54	0.39	0.41	0.22	0.23
Black or African American	1.25	1.34	1.04	1.01	0.60	0.48
American Indian or Alaska Native	4.61	5.89	4.13	4.93	2.53	3.13
Native Hawaiian or Other Pacific Islander	*	*	*	*	3.71	2.26
Asian	2.70	2.73	1.21	1.32	0.46	0.62
Two or More Races	3.54	3.32	2.50	2.98	1.36	1.43
Hispanic or Latino	1.29	1.15	1.00	0.97	0.51	0.48
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.73	0.71	0.61	0.63	0.40	0.40
Female, White, Not Hispanic	0.64	0.74	0.43	0.43	0.19	0.22
Male, Black, Not Hispanic	2.10	1.94	1.89	1.63	1.16	0.95
Female, Black, Not Hispanic	1.38	1.64	0.92	1.11	0.48	0.41
Male, Hispanic	1.90	1.70	1.68	1.58	0.97	0.83
Female, Hispanic	1.56	1.60	1.01	1.04	0.21	0.35

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.106P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.6426		0.8846		0.5786	
GENDER						
Male	0.6086		0.9011		0.6408	
Female	0.2240		0.9434		0.7539	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.3536		0.6982		0.5053	
White	0.1386		0.5820		0.3946	
Black or African American	0.4956		0.4460		0.7432	
American Indian or Alaska Native	0.3465		0.4974		0.4338	
Native Hawaiian or Other Pacific Islander	*		*		0.1174	
Asian	0.5418		0.2974		0.3395	
Two or More Races	0.0221		0.6151		0.9152	
Hispanic or Latino	0.2851		0.5005		0.8669	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.5834		0.4266		0.4192	
Female, White, Not Hispanic	0.0980		0.9168		0.7392	
Male, Black, Not Hispanic	0.2179		0.1523		0.6575	
Female, Black, Not Hispanic	0.7831		0.5619		0.9185	
Male, Hispanic	0.1507		0.5263		0.7207	
Female, Hispanic	0.9293		0.7202		0.6361	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.107C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	115	113	120	116	114	108
NOT HISPANIC OR LATINO¹	106	102	108	104	105	99
White	87	86	87	89	87	84
Black or African American	42	44	39	41	31	33
American Indian or Alaska Native	8	7	8	7	7	5
Native Hawaiian or Other Pacific Islander	7	6	7	6	6	6
Chinese	13	12	11	10	7	8
Filipino	8	9	8	10	6	6
Japanese	*	*	*	*	3	*
Asian Indian	10	12	9	12	9	8
Korean	7	9	8	9	7	8
Vietnamese	9	10	10	9	8	7
HISPANIC OR LATINO¹	46	43	47	45	39	42
Mexican	40	37	41	37	33	36
Puerto Rican	15	15	16	15	13	12
Central or South American	16	14	16	16	15	16
Cuban	6	8	7	7	8	7

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.107D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.31	0.30	0.32	0.31	0.31	0.29
NOT HISPANIC OR LATINO¹	0.34	0.33	0.35	0.33	0.34	0.32
White	0.37	0.37	0.37	0.38	0.37	0.36
Black or African American	0.77	0.79	0.72	0.73	0.58	0.60
American Indian or Alaska Native	3.01	2.97	3.01	2.80	2.84	2.16
Native Hawaiian or Other Pacific Islander	4.71	4.75	5.26	4.84	4.38	4.83
Chinese	4.12	3.65	3.54	3.16	2.15	2.57
Filipino	3.00	3.43	2.99	3.64	2.23	2.39
Japanese	*	*	*	*	4.01	*
Asian Indian	3.30	3.62	3.12	3.56	2.86	2.59
Korean	4.76	5.58	4.82	5.52	4.23	4.99
Vietnamese	4.87	5.78	5.12	5.43	4.10	4.25
HISPANIC OR LATINO¹	0.75	0.69	0.78	0.72	0.64	0.67
Mexican	0.92	0.84	0.95	0.85	0.76	0.81
Puerto Rican	2.13	2.09	2.23	2.17	1.77	1.79
Central or South American	2.30	2.06	2.32	2.29	2.19	2.33
Cuban	3.58	3.76	3.83	3.70	4.36	3.44

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.107P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.0197		0.4111		0.8867	
NOT HISPANIC OR LATINO¹	0.0194		0.2574		0.5265	
White	0.1605		0.6220		0.7704	
Black or African American	0.0393		0.3040		0.8791	
American Indian or Alaska Native	0.1089		0.3735		0.1686	
Native Hawaiian or Other Pacific Islander	0.8308		0.4383		0.8937	
Chinese	0.0064		0.0615		0.0384	
Filipino	0.1071		0.2091		0.1428	
Japanese	*		*		*	
Asian Indian	0.8828		0.9636		0.4526	
Korean	0.1074		0.1342		0.3751	
Vietnamese	0.3938		0.5515		0.5010	
HISPANIC OR LATINO¹	0.5463		0.5801		0.2628	
Mexican	0.7962		0.3452		0.2335	
Puerto Rican	0.0764		0.2433		0.0921	
Central or South American	0.8239		0.7716		0.5772	
Cuban	0.0702		0.1962		0.3007	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.108C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	438	443	644	636	673	665
NOT HISPANIC OR LATINO¹	399	415	606	618	637	632
White	303	320	522	548	568	568
Black or African American	161	177	211	201	189	213
American Indian or Alaska Native	37	27	40	49	47	38
Native Hawaiian or Other Pacific Islander	*	*	29	*	28	25
Chinese	54	70	72	76	68	67
Filipino	64	78	80	104	75	93
Japanese	*	28	*	42	46	45
Asian Indian	60	65	64	64	59	62
Korean	*	*	*	*	38	35
Vietnamese	*	*	*	*	*	30
HISPANIC OR LATINO¹	175	181	239	226	232	209
Mexican	143	141	186	184	175	163
Puerto Rican	56	58	75	70	74	67
Central or South American	69	77	87	92	85	81
Cuban	28	35	38	47	42	42

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.108D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.22	0.22	0.32	0.32	0.34	0.33
NOT HISPANIC OR LATINO¹	0.23	0.23	0.35	0.35	0.36	0.36
White	0.21	0.22	0.37	0.38	0.40	0.40
Black or African American	0.74	0.80	0.97	0.91	0.87	0.96
American Indian or Alaska Native	3.34	2.62	3.57	4.75	4.22	3.70
Native Hawaiian or Other Pacific Islander	*	*	5.67	*	5.52	5.81
Chinese	3.02	3.55	4.00	3.87	3.78	3.42
Filipino	3.51	4.06	4.40	5.42	4.08	4.83
Japanese	*	3.46	*	5.30	6.03	5.67
Asian Indian	3.54	3.39	3.73	3.30	3.46	3.19
Korean	*	*	*	*	5.99	6.05
Vietnamese	*	*	*	*	*	5.46
HISPANIC OR LATINO¹	0.75	0.75	1.02	0.94	0.99	0.87
Mexican	0.94	0.91	1.23	1.18	1.16	1.05
Puerto Rican	2.28	2.43	3.03	2.92	2.99	2.78
Central or South American	2.04	2.07	2.57	2.48	2.53	2.17
Cuban	2.58	3.09	3.48	4.19	3.82	3.74

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.108P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.1544		0.0814		0.2408	
NOT HISPANIC OR LATINO¹	0.6052		0.4177		0.6167	
White	0.9267		0.9933		0.8065	
Black or African American	0.2474		0.1119		0.0676	
American Indian or Alaska Native	0.5721		0.2600		0.3211	
Native Hawaiian or Other Pacific Islander	*		*		0.8416	
Chinese	0.4874		0.6532		0.3565	
Filipino	0.3598		0.9651		0.2617	
Japanese	*		*		0.1840	
Asian Indian	0.7327		0.3952		0.6927	
Korean	*		*		0.3574	
Vietnamese	*		*		*	
HISPANIC OR LATINO¹	0.0542		0.0124		0.0367	
Mexican	0.2155		0.0181		0.0230	
Puerto Rican	0.1844		0.9916		0.8112	
Central or South American	0.3699		0.3270		0.2108	
Cuban	0.4099		0.7032		0.4802	

*Low precision; no estimate reported.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

**Table 2.109C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004**

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	114	108	102	99	55	63
NOT HISPANIC OR LATINO¹	105	99	92	89	52	59
White	87	84	79	78	48	53
Black or African American	31	33	24	25	10	11
American Indian or Alaska Native	7	5	6	5	2	3
Native Hawaiian or Other Pacific Islander	6	6	6	6	2	2
Chinese	7	8	6	7	3	4
Filipino	6	6	5	5	2	2
Japanese	3	*	3	*	*	2
Asian Indian	9	8	5	6	2	2
Korean	7	8	5	5	*	*
Vietnamese	8	7	4	3	*	0
HISPANIC OR LATINO¹	39	42	36	39	17	21
Mexican	33	36	30	32	16	18
Puerto Rican	13	12	11	11	5	5
Central or South American	15	16	13	15	5	5
Cuban	8	7	8	6	3	4

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.109D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.31	0.29	0.27	0.26	0.15	0.17
NOT HISPANIC OR LATINO¹	0.34	0.32	0.29	0.28	0.17	0.19
White	0.37	0.36	0.34	0.34	0.21	0.23
Black or African American	0.58	0.60	0.44	0.44	0.18	0.20
American Indian or Alaska Native	2.84	2.16	2.19	1.90	0.84	1.09
Native Hawaiian or Other Pacific Islander	4.38	4.83	4.04	4.56	1.50	1.54
Chinese	2.15	2.57	1.87	2.10	1.09	1.17
Filipino	2.23	2.39	2.00	1.88	0.88	0.91
Japanese	4.01	*	3.62	*	*	2.67
Asian Indian	2.86	2.59	1.78	1.72	0.76	0.57
Korean	4.23	4.99	3.23	3.34	*	*
Vietnamese	4.10	4.25	2.27	1.71	*	0.20
HISPANIC OR LATINO¹	0.64	0.67	0.59	0.62	0.29	0.33
Mexican	0.76	0.81	0.69	0.73	0.37	0.41
Puerto Rican	1.77	1.79	1.48	1.51	0.70	0.73
Central or South American	2.19	2.33	1.89	2.19	0.67	0.75
Cuban	4.36	3.44	4.34	3.02	1.73	1.98

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.109P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.8867		0.5327		0.7247	
NOT HISPANIC OR LATINO¹	0.5265		0.8495		0.8441	
White	0.7704		0.8658		0.6493	
Black or African American	0.8791		0.8336		0.8469	
American Indian or Alaska Native	0.1686		0.3632		0.3604	
Native Hawaiian or Other Pacific Islander	0.8937		0.8489		0.9915	
Chinese	0.0384		0.4493		0.5625	
Filipino	0.1428		0.7146		0.8099	
Japanese	*		*		*	
Asian Indian	0.4526		0.9152		0.4804	
Korean	0.3751		0.7278		*	
Vietnamese	0.5010		0.2369		*	
HISPANIC OR LATINO¹	0.2628		0.0331		0.5213	
Mexican	0.2335		0.0399		0.9007	
Puerto Rican	0.0921		0.1842		0.7845	
Central or South American	0.5772		0.0709		0.7344	
Cuban	0.3007		0.0445		0.4741	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.110C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	673	665	500	479	276	265
NOT HISPANIC OR LATINO¹	637	632	480	455	253	247
White	568	568	418	409	228	232
Black or African American	189	213	158	161	88	84
American Indian or Alaska Native	47	38	30	33	19	21
Native Hawaiian or Other Pacific Islander	28	25	20	22	10	9
Chinese	68	67	36	36	8	19
Filipino	75	93	39	43	14	18
Japanese	46	45	22	22	4	5
Asian Indian	59	62	36	35	14	7
Korean	38	35	26	25	*	12
Vietnamese	*	30	19	12	11	7
HISPANIC OR LATINO¹	232	209	182	168	93	87
Mexican	175	163	151	141	72	76
Puerto Rican	74	67	60	58	36	33
Central or South American	85	81	65	62	23	19
Cuban	42	42	41	39	*	15

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.110D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2003-2004

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.34	0.33	0.25	0.24	0.14	0.13
NOT HISPANIC OR LATINO¹	0.36	0.36	0.27	0.26	0.14	0.14
White	0.40	0.40	0.29	0.29	0.16	0.16
Black or African American	0.87	0.96	0.73	0.73	0.40	0.38
American Indian or Alaska Native	4.22	3.70	2.69	3.21	1.72	2.04
Native Hawaiian or Other Pacific Islander	5.52	5.81	3.91	5.09	1.94	2.14
Chinese	3.78	3.42	1.98	1.84	0.46	0.98
Filipino	4.08	4.83	2.12	2.23	0.78	0.95
Japanese	6.03	5.67	2.86	2.73	0.58	0.63
Asian Indian	3.46	3.19	2.14	1.82	0.84	0.38
Korean	5.99	6.05	4.16	4.25	*	2.08
Vietnamese	*	5.46	2.97	2.20	1.78	1.22
HISPANIC OR LATINO¹	0.99	0.87	0.78	0.69	0.40	0.36
Mexican	1.16	1.05	0.99	0.90	0.48	0.49
Puerto Rican	2.99	2.78	2.43	2.44	1.45	1.37
Central or South American	2.53	2.17	1.92	1.67	0.69	0.51
Cuban	3.82	3.74	3.75	3.43	*	1.30

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.110P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Racial/Ethnic Subgroups: *P* Values from Tests of Differences of Percentages, 2003-2004 Versus 2002-2003

Racial/Ethnic Subgroup	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002-2003	2003-2004	2002-2003	2003-2004	2002-2003	2003-2004
TOTAL¹	0.2408		0.6725		0.4162	
NOT HISPANIC OR LATINO¹	0.6167		0.9580		0.1737	
White	0.8065		0.4290		0.1065	
Black or African American	0.0676		0.0257		0.9548	
American Indian or Alaska Native	0.3211		0.7924		0.7209	
Native Hawaiian or Other Pacific Islander	0.8416		0.5258		0.2816	
Chinese	0.3565		0.3109		0.1353	
Filipino	0.2617		0.9862		0.4180	
Japanese	0.1840		0.9198		0.4863	
Asian Indian	0.6927		0.7941		0.5731	
Korean	0.3574		0.8670		*	
Vietnamese	*		0.9920		0.8061	
HISPANIC OR LATINO¹	0.0367		0.2949		0.2684	
Mexican	0.0230		0.0872		0.4386	
Puerto Rican	0.8112		0.1474		0.9743	
Central or South American	0.2108		0.3103		0.4410	
Cuban	0.4802		0.7951		*	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

¹ Totals include data from respondents reporting racial/ethnic subgroups not shown, as well as respondents reporting more than one subgroup.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, and 2004.

Table 2.111C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	160	154	165	154	154	148
GEOGRAPHIC DIVISION						
Northeast	55	71	65	70	58	68
New England	20	25	31	31	28	40
Middle Atlantic	51	66	56	63	51	55
Midwest	66	71	69	72	63	73
East North Central	49	59	51	59	50	57
West North Central	44	39	46	42	39	45
South	101	83	99	83	97	76
South Atlantic	76	60	75	57	77	52
East South Central	35	34	34	37	29	31
West South Central	54	46	54	49	49	45
West	91	79	91	77	81	75
Mountain	40	35	36	35	29	30
Pacific	81	72	84	69	76	68
COUNTY TYPE						
Large Metro	122	124	120	122	110	112
Small Metro	80	86	89	90	87	91
250K – 1 Mil. Pop.	66	67	74	71	73	72
< 250K Pop.	44	51	47	54	46	54
Nonmetro	58	63	62	64	62	53
Urbanized	40	39	41	39	41	39
Less Urbanized	43	47	44	45	44	37
Completely Rural	15	17	15	16	13	13

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.111D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.43	0.41	0.44	0.41	0.41	0.39
GEOGRAPHIC DIVISION						
Northeast	0.83	1.04	0.97	1.04	0.87	1.00
New England	1.16	1.45	1.80	1.77	1.65	2.27
Middle Atlantic	1.04	1.32	1.14	1.25	1.02	1.08
Midwest	0.76	0.83	0.80	0.84	0.73	0.85
East North Central	0.81	0.99	0.85	0.98	0.83	0.95
West North Central	1.68	1.54	1.76	1.66	1.49	1.79
South	0.75	0.62	0.74	0.62	0.72	0.57
South Atlantic	1.13	0.90	1.11	0.85	1.15	0.79
East South Central	1.55	1.57	1.52	1.69	1.30	1.42
West South Central	1.23	1.02	1.22	1.08	1.11	1.00
West	1.03	0.88	1.03	0.86	0.92	0.83
Mountain	1.57	1.34	1.41	1.37	1.16	1.18
Pacific	1.29	1.11	1.33	1.07	1.20	1.06
COUNTY TYPE						
Large Metro	0.62	0.63	0.61	0.62	0.56	0.57
Small Metro	0.71	0.72	0.79	0.76	0.77	0.76
250K – 1 Mil. Pop.	0.88	0.86	0.98	0.91	0.97	0.93
< 250K Pop.	1.15	1.24	1.23	1.32	1.20	1.33
Nonmetro	0.90	1.01	0.95	1.03	0.96	0.85
Urbanized	1.36	1.47	1.41	1.47	1.41	1.46
Less Urbanized	1.47	1.53	1.48	1.48	1.49	1.19
Completely Rural	2.49	3.49	2.49	3.31	2.22	2.60

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.111P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.1120		0.6875		0.6311	
GEOGRAPHIC DIVISION						
Northeast	0.9522		0.6162		0.7762	
New England	0.7765		0.5668		0.8464	
Middle Atlantic	0.8717		0.7422		0.8322	
Midwest	1.0000		0.9199		0.7857	
East North Central	0.6980		0.7387		0.7277	
West North Central	0.6242		0.8438		0.9582	
South	0.0128		0.1466		0.5770	
South Atlantic	0.0507		0.2321		0.3543	
East South Central	0.6980		0.6506		0.2601	
West South Central	0.0277		0.1513		0.6086	
West	0.9329		0.6773		0.7797	
Mountain	0.9090		0.9253		0.7424	
Pacific	0.8841		0.6817		0.8587	
COUNTY TYPE						
Large Metro	0.2615		0.9113		0.4460	
Small Metro	0.9389		0.7386		0.1454	
250K – 1 Mil. Pop.	0.6911		0.8105		0.3506	
< 250K Pop.	0.7011		0.3707		0.2304	
Nonmetro	0.0682		0.2014		0.0505	
Urbanized	0.1726		0.1910		0.1243	
Less Urbanized	0.2937		0.4080		0.1131	
Completely Rural	0.5379		0.7390		0.3779	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.112C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	569	679	874	875	899	943
GEOGRAPHIC DIVISION						
Northeast	236	269	354	376	365	412
New England	103	139	186	142	195	163
Middle Atlantic	213	231	300	347	307	375
Midwest	200	199	352	342	364	371
East North Central	176	154	270	291	280	296
West North Central	95	125	225	178	233	221
South	378	452	557	575	576	611
South Atlantic	263	366	446	443	446	481
East South Central	177	157	223	182	244	189
West South Central	199	211	254	314	275	314
West	288	379	439	419	456	458
Mountain	161	156	220	196	216	213
Pacific	239	346	382	371	402	406
COUNTY TYPE						
Large Metro	421	500	641	637	680	675
Small Metro	293	337	485	477	458	514
250K – 1 Mil. Pop.	206	272	385	358	375	386
< 250K Pop.	216	190	307	289	264	312
Nonmetro	252	295	348	360	351	369
Urbanized	155	139	224	214	241	221
Less Urbanized	190	239	260	260	226	250
Completely Rural	76	68	123	98	102	116

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.112D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.28	0.33	0.44	0.43	0.45	0.46
GEOGRAPHIC DIVISION						
Northeast	0.61	0.70	0.92	0.97	0.94	1.07
New England	1.01	1.37	1.82	1.40	1.92	1.60
Middle Atlantic	0.75	0.81	1.06	1.22	1.08	1.32
Midwest	0.44	0.44	0.78	0.75	0.81	0.81
East North Central	0.56	0.48	0.85	0.91	0.88	0.93
West North Central	0.71	0.91	1.67	1.30	1.73	1.61
South	0.53	0.62	0.78	0.79	0.81	0.84
South Atlantic	0.70	0.95	1.18	1.15	1.18	1.25
East South Central	1.48	1.29	1.87	1.49	2.04	1.55
West South Central	0.92	0.96	1.17	1.42	1.27	1.43
West	0.64	0.83	0.97	0.92	1.01	1.00
Mountain	1.23	1.16	1.68	1.46	1.65	1.59
Pacific	0.75	1.07	1.20	1.15	1.26	1.26
COUNTY TYPE						
Large Metro	0.40	0.46	0.60	0.59	0.64	0.62
Small Metro	0.49	0.56	0.81	0.80	0.76	0.86
250K – 1 Mil. Pop.	0.52	0.69	0.98	0.91	0.95	0.98
< 250K Pop.	1.05	0.93	1.49	1.42	1.28	1.53
Nonmetro	0.75	0.86	1.03	1.05	1.04	1.07
Urbanized	1.04	0.99	1.50	1.53	1.61	1.58
Less Urbanized	1.24	1.39	1.70	1.51	1.48	1.45
Completely Rural	2.17	2.21	3.53	3.20	2.92	3.79

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.112P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.1183		0.7763		0.6426	
GEOGRAPHIC DIVISION						
Northeast	0.8892		0.6137		0.3195	
New England	0.3720		0.2095		0.6266	
Middle Atlantic	0.7744		0.9972		0.3799	
Midwest	0.1976		0.8418		0.3333	
East North Central	0.7172		0.6316		0.1738	
West North Central	0.0798		0.7176		0.8564	
South	0.3865		0.6968		0.8303	
South Atlantic	0.0867		0.9764		0.1965	
East South Central	0.1776		0.2874		0.3292	
West South Central	0.9488		0.9720		0.4149	
West	0.2537		0.6892		0.7383	
Mountain	0.9226		0.6297		0.7231	
Pacific	0.1619		0.4224		0.5379	
COUNTY TYPE						
Large Metro	0.2763		0.3088		0.4517	
Small Metro	0.2975		0.4287		0.4376	
250K – 1 Mil. Pop.	0.1544		0.4157		0.3841	
< 250K Pop.	0.9469		0.8059		0.8822	
Nonmetro	0.6238		0.1551		0.1336	
Urbanized	0.7715		0.5665		0.9080	
Less Urbanized	0.9910		0.0141		0.0293	
Completely Rural	0.4405		0.5049		0.1056	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.113C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	154	148	135	138	75	92
GEOGRAPHIC DIVISION						
Northeast	58	68	53	63	31	52
New England	28	40	26	38	14	37
Middle Atlantic	51	55	46	50	27	36
Midwest	63	73	59	70	34	41
East North Central	50	57	50	50	28	27
West North Central	39	45	31	49	19	31
South	97	76	79	70	44	49
South Atlantic	77	52	64	43	34	32
East South Central	29	31	27	29	15	17
West South Central	49	45	38	47	23	32
West	81	75	74	69	40	41
Mountain	29	30	26	28	19	15
Pacific	76	68	70	63	35	38
COUNTY TYPE						
Large Metro	110	112	99	100	50	58
Small Metro	87	91	75	83	44	63
250K – 1 Mil. Pop.	73	72	63	66	34	48
< 250K Pop.	46	54	42	53	29	40
Nonmetro	62	53	55	47	35	28
Urbanized	41	39	36	31	28	19
Less Urbanized	44	37	39	34	19	17
Completely Rural	13	13	12	12	5	8

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.113D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.41	0.39	0.36	0.36	0.20	0.24
GEOGRAPHIC DIVISION						
Northeast	0.87	1.00	0.80	0.93	0.46	0.77
New England	1.65	2.27	1.55	2.15	0.83	2.10
Middle Atlantic	1.02	1.08	0.92	0.99	0.54	0.72
Midwest	0.73	0.85	0.68	0.81	0.39	0.48
East North Central	0.83	0.95	0.83	0.83	0.46	0.45
West North Central	1.49	1.79	1.19	1.93	0.72	1.21
South	0.72	0.57	0.59	0.52	0.33	0.36
South Atlantic	1.15	0.79	0.96	0.65	0.51	0.48
East South Central	1.30	1.42	1.19	1.31	0.69	0.78
West South Central	1.11	1.00	0.86	1.03	0.52	0.71
West	0.92	0.83	0.84	0.77	0.45	0.46
Mountain	1.16	1.18	1.01	1.10	0.74	0.60
Pacific	1.20	1.06	1.11	0.98	0.56	0.59
COUNTY TYPE						
Large Metro	0.56	0.57	0.50	0.51	0.25	0.30
Small Metro	0.77	0.76	0.67	0.70	0.39	0.53
250K – 1 Mil. Pop.	0.97	0.93	0.84	0.84	0.46	0.62
< 250K Pop.	1.20	1.33	1.10	1.29	0.76	0.97
Nonmetro	0.96	0.85	0.84	0.75	0.54	0.44
Urbanized	1.41	1.46	1.25	1.18	0.95	0.73
Less Urbanized	1.49	1.19	1.32	1.11	0.64	0.55
Completely Rural	2.22	2.60	1.97	2.38	0.86	1.64

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 2.113P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2004 Versus 2003**

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.6311		0.3790		0.5649	
GEOGRAPHIC DIVISION						
Northeast	0.7762		0.4216		0.2102	
New England	0.8464		0.4089		0.7556	
Middle Atlantic	0.8322		0.6375		0.1849	
Midwest	0.7857		0.6645		0.7699	
East North Central	0.7277		0.9478		0.3899	
West North Central	0.9582		0.4178		0.6623	
South	0.5770		0.7479		0.7343	
South Atlantic	0.3543		0.8796		0.9155	
East South Central	0.2601		0.7716		0.4438	
West South Central	0.6086		0.5819		0.7675	
West	0.7797		0.7140		0.7801	
Mountain	0.7424		0.6778		0.0410	
Pacific	0.8587		0.8093		0.4685	
COUNTY TYPE						
Large Metro	0.4460		0.7431		0.9650	
Small Metro	0.1454		0.1701		0.3539	
250K – 1 Mil. Pop.	0.3506		0.2573		0.8822	
< 250K Pop.	0.2304		0.4968		0.1965	
Nonmetro	0.0505		0.6713		0.7839	
Urbanized	0.1243		0.2033		0.4964	
Less Urbanized	0.1131		0.9816		0.8956	
Completely Rural	0.3779		0.1654		0.1343	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.114C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	899	943	661	688	366	371
GEOGRAPHIC DIVISION						
Northeast	365	412	297	324	146	162
New England	195	163	165	156	70	76
Middle Atlantic	307	375	249	284	128	143
Midwest	364	371	279	296	168	190
East North Central	280	296	227	257	143	170
West North Central	233	221	163	147	89	85
South	576	611	408	421	228	215
South Atlantic	446	481	304	322	173	146
East South Central	244	189	163	127	78	73
West South Central	275	314	216	237	123	140
West	456	458	319	322	181	168
Mountain	216	213	147	131	86	65
Pacific	402	406	282	294	159	155
COUNTY TYPE						
Large Metro	680	675	503	512	281	265
Small Metro	458	514	390	391	194	233
250K – 1 Mil. Pop.	375	386	315	301	160	184
< 250K Pop.	264	312	204	235	100	133
Nonmetro	351	369	229	265	135	131
Urbanized	241	221	156	170	88	81
Less Urbanized	226	250	163	180	91	98
Completely Rural	102	116	65	67	44	52

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.114D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.45	0.46	0.33	0.34	0.18	0.18
GEOGRAPHIC DIVISION						
Northeast	0.94	1.07	0.77	0.84	0.38	0.42
New England	1.92	1.60	1.62	1.53	0.69	0.75
Middle Atlantic	1.08	1.32	0.88	1.00	0.45	0.50
Midwest	0.81	0.81	0.62	0.65	0.37	0.42
East North Central	0.88	0.93	0.72	0.81	0.45	0.53
West North Central	1.73	1.61	1.21	1.07	0.66	0.62
South	0.81	0.84	0.57	0.58	0.32	0.30
South Atlantic	1.18	1.25	0.81	0.84	0.46	0.38
East South Central	2.04	1.55	1.36	1.04	0.66	0.60
West South Central	1.27	1.43	0.99	1.08	0.57	0.63
West	1.01	1.00	0.71	0.71	0.40	0.37
Mountain	1.65	1.59	1.12	0.98	0.65	0.48
Pacific	1.26	1.26	0.88	0.91	0.50	0.48
COUNTY TYPE						
Large Metro	0.64	0.62	0.47	0.47	0.26	0.24
Small Metro	0.76	0.86	0.65	0.65	0.32	0.39
250K – 1 Mil. Pop.	0.95	0.98	0.80	0.76	0.41	0.47
< 250K Pop.	1.28	1.53	0.99	1.15	0.48	0.65
Nonmetro	1.04	1.07	0.68	0.77	0.40	0.38
Urbanized	1.61	1.58	1.04	1.21	0.59	0.57
Less Urbanized	1.48	1.45	1.07	1.05	0.59	0.57
Completely Rural	2.92	3.79	1.86	2.17	1.27	1.69

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 2.114P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2003	2004	2003	2004	2003	2004
TOTAL	0.6426		0.8846		0.5786	
GEOGRAPHIC DIVISION						
Northeast	0.3195		0.4493		0.5346	
New England	0.6266		0.6203		0.5740	
Middle Atlantic	0.3799		0.5614		0.6798	
Midwest	0.3333		0.3716		0.0989	
East North Central	0.1738		0.4325		0.0923	
West North Central	0.8564		0.6686		0.7206	
South	0.8303		0.8114		0.5225	
South Atlantic	0.1965		0.6994		0.9228	
East South Central	0.3292		0.8238		0.3311	
West South Central	0.4149		0.8224		0.5612	
West	0.7383		0.1097		0.3170	
Mountain	0.7231		0.6118		0.5892	
Pacific	0.5379		0.1178		0.3959	
COUNTY TYPE						
Large Metro	0.4517		0.9401		0.8591	
Small Metro	0.4376		0.6580		0.1840	
250K – 1 Mil. Pop.	0.3841		0.6904		0.3000	
< 250K Pop.	0.8822		0.8284		0.3991	
Nonmetro	0.1336		0.2537		0.8655	
Urbanized	0.9080		0.4055		0.7294	
Less Urbanized	0.0293		0.6930		0.6545	
Completely Rural	0.1056		0.0731		0.2501	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.