

Table 6.10A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Numbers in Thousands, 2003 and 2004

Dependence/Abuse	Total		DEPENDENCE ¹				DEPENDENCE OR ABUSE ¹			
			Yes		No		Yes		No	
	2003	2004	2003	2004	2003	2004	2003	2004	2003	2004
ILLCIT DRUG²	19,588	21,417	1,298	1,492	18,290	19,925	1,694	1,977	17,895	19,440
Marijuana	19,588	21,417	649	664	18,939	20,753	927	1,024	18,662	20,392
ILLCIT DRUG OTHER THAN MARIJUANA²	19,588	21,417	775	1,000	18,813	20,416	1,035	1,216	18,553	20,201
ALCOHOL	19,588	21,417	1,941	2,367	17,648	19,050	3,334	3,561	16,254	17,856
ILLCIT DRUG OR ALCOHOL²	19,588	21,417	2,875	3,258	16,713	18,159	4,179	4,559	15,409	16,858
BOTH ILLICIT DRUG AND ALCOHOL²	19,588	21,417	364	*	19,224	20,816	848	978	18,740	20,438

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.10B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Percentages, 2003 and 2004

Dependence/Abuse	Total		DEPENDENCE ¹				DEPENDENCE OR ABUSE ¹			
			Yes		No		Yes		No	
	2003	2004	2003	2004	2003	2004	2003	2004	2003	2004
ILLCIT DRUG²	9.2	9.9	35.3	37.4	8.8	9.4	30.5	33.5	8.6	9.3
Marijuana	9.2	9.9	32.2	35.9	9.0	9.7	28.6	32.5	8.9	9.6
ILLCIT DRUG OTHER THAN MARIJUANA²	9.2	9.9	39.6	41.2	8.9	9.6	34.9	36.3	8.8	9.5
ALCOHOL	9.2	9.9	27.5	30.8	8.6	9.2	20.4	21.0	8.3	9.0
ILLCIT DRUG OR ALCOHOL²	9.2	9.9	29.2	31.5	8.2	8.9	21.6	22.9	8.0	8.6
BOTH ILLICIT DRUG AND ALCOHOL²	9.2	9.9	40.5	*	9.1	9.7	33.8	33.1	8.9	9.6

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.11A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Numbers in Thousands, 2003 and 2004

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE²						
Illicit Drug ³	5,548	5,894	1,694	1,977	3,854	3,918
Marijuana	3,243	3,148	927	1,024	2,316	2,123
Illicit Drug Other Than Marijuana ³	2,969	3,351	1,035	1,216	1,933	2,136
Alcohol	16,334	16,989	3,334	3,561	13,000	13,428
Illicit Drug or Alcohol ³	19,372	19,929	4,179	4,559	15,193	15,370
Both Illicit Drug and Alcohol ³	2,509	2,954	848	978	1,661	1,976
DEPENDENCE²						
Illicit Drug ³	3,680	3,987	1,298	1,492	2,382	2,495
Marijuana	2,018	1,847	649	664	1,368	1,184
Illicit Drug Other Than Marijuana ³	1,956	2,429	775	1,000	1,181	1,429
Alcohol	7,051	7,672	1,941	2,367	5,110	5,306
Illicit Drug or Alcohol ³	9,832	10,351	2,875	3,258	6,957	7,093
Both Illicit Drug and Alcohol ³	899 ^a	1,309	364 ^a	601	535	708

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.11B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Percentages, 2003 and 2004

Dependence/Abuse	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE²						
Illicit Drug ³	2.6	2.7	8.6	9.2	2.0	2.0
Marijuana	1.5	1.5	4.7	4.8	1.2	1.1
Illicit Drug Other Than Marijuana ³	1.4	1.6	5.3	5.7	1.0	1.1
Alcohol	7.7	7.9	17.0	16.6	6.7	6.9
Illicit Drug or Alcohol ³	9.1	9.3	21.3	21.3	7.9	7.9
Both Illicit Drug and Alcohol ³	1.2	1.4	4.3	4.6	0.9	1.0
DEPENDENCE²						
Illicit Drug ³	1.7	1.9	6.6	7.0	1.2	1.3
Marijuana	0.9	0.9	3.3	3.1	0.7	0.6
Illicit Drug Other Than Marijuana ³	0.9	1.1	4.0	4.7	0.6	0.7
Alcohol	3.3	3.6	9.9	11.1	2.6	2.7
Illicit Drug or Alcohol ³	4.6	4.8	14.7	15.2	3.6	3.7
Both Illicit Drug and Alcohol ³	0.4 ^a	0.6	1.9	2.8	0.3	0.4

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.12A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2003 and 2004

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE ^{1,2}			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	19,588	21,417	1,694	1,977	17,895	19,440
18-25	4,420	4,413	734	852	3,687	3,561
26-49	10,418	10,408	906	1,020	9,512	9,389
50 or Older	4,750 ^a	6,596	*	*	4,696 ^a	6,491
MALE	6,887	7,976	786 ^a	1,212	6,100	6,763
18-25	1,690	1,640	355	417	1,335	1,223
26-49	3,438	3,941	418	*	3,020	3,230
50 or Older	1,758	2,395	*	*	1,744	2,310
FEMALE	12,702	13,441	907	764	11,794	12,677
18-25	2,730	2,773	378	435	2,351	2,337
26-49	6,980	6,467	488	*	6,491	6,158
50 or Older	2,992 ^a	4,201	*	*	2,952 ^a	4,181

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.12B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Dependence or Abuse, Gender, and Age Group: Percentages, 2003 and 2004

Gender/Age	Total		ILLCIT DRUG DEPENDENCE OR ABUSE ^{1,2}			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	9.2	9.9	30.5	33.5	8.6	9.3
18-25	13.9	13.7	29.5	32.7	12.6	12.0
26-49	10.4	10.4	32.1	34.6	9.8	9.7
50 or Older	5.9 ^a	7.9	*	*	5.8 ^a	7.9
MALE	6.7	7.7	22.3 ^a	30.2	6.2	6.8
18-25	10.6	10.1	22.8	24.4	9.3	8.5
26-49	7.0	8.0	23.5	*	6.4	6.8
50 or Older	4.7	6.3	*	*	4.7	6.1
FEMALE	11.5	12.0	44.7	40.5	10.9	11.5
18-25	17.3	17.3	40.8	48.6	15.8	15.5
26-49	13.8	12.7	46.8	*	13.1	12.3
50 or Older	6.8 ^a	9.3	*	*	6.7 ^a	9.3

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.13A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Numbers in Thousands, 2003 and 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	5,548	5,894	1,694	1,977	3,854	3,918
18-25	2,486	2,603	734	852	1,752	1,751
26-49	2,820	2,948	906	1,020	1,914	1,928
50 or Older	242	344	54	105	188	239
MALE	3,519	4,009	786 ^a	1,212	2,733	2,796
18-25	1,558	1,707	355	417	1,203	1,290
26-49	1,777	2,156	418	711	1,360	1,446
50 or Older	184	145	*	*	170	60
FEMALE	2,028	1,885	907	764	1,121	1,121
18-25	928	896	378	435	550	460
26-49	1,042	791	488	309	554	482
50 or Older	58	198	41	20	17	179

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.13B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Percentages, 2003 and 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	2.6	2.7	8.6	9.2	2.0	2.0
18-25	7.8	8.1	16.6	19.3	6.4	6.3
26-49	2.8	2.9	8.7	9.8	2.1	2.1
50 or Older	0.3	0.4	1.1	1.6	0.2	0.3
MALE	3.4	3.9	11.4	15.2	2.9	2.9
18-25	9.8	10.6	21.0	25.4	8.5	8.9
26-49	3.6	4.4	12.1	18.0	3.0	3.2
50 or Older	0.5	0.4	*	*	0.5	0.2
FEMALE	1.8	1.7	7.1	5.7	1.1	1.1
18-25	5.9	5.6	13.9	15.7	4.2	3.5
26-49	2.1	1.6	7.0	4.8	1.3	1.1
50 or Older	0.1	0.4	1.4	0.5	0.0	0.4

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.14A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2003 and 2004

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	19,588	21,417	3,334	3,561	16,254	17,856
18-25	4,420	4,413	1,200	1,313	3,220	3,100
26-49	10,418	10,408	1,756	1,882	8,661	8,527
50 or Older	4,750 ^a	6,596	377	*	4,373 ^a	6,229
MALE	6,887	7,976	1,715	1,909	5,172	6,066
18-25	1,690	1,640	596	623	1,095	1,017
26-49	3,438	3,941	865	1,076	2,573	2,865
50 or Older	1,758	2,395	254	*	1,504	2,185
FEMALE	12,702	13,441	1,619	1,651	11,083	11,789
18-25	2,730	2,773	605	689	2,125	2,083
26-49	6,980	6,467	891	805	6,089	5,662
50 or Older	2,992 ^a	4,201	*	*	2,869 ^a	4,044

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.14B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2003 and 2004

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	9.2	9.9	20.4	21.0	8.3	9.0
18-25	13.9	13.7	22.0	22.6	12.3	11.7
26-49	10.4	10.4	20.5	21.2	9.5	9.3
50 or Older	5.9 ^a	7.9	16.5	*	5.6 ^a	7.7
MALE	6.7	7.7	15.4	16.1	5.7	6.6
18-25	10.6	10.1	17.2	16.5	8.8	8.2
26-49	7.0	8.0	14.6	16.7	5.9	6.7
50 or Older	4.7	6.3	14.9	*	4.3	6.0
FEMALE	11.5	12.0	31.0	32.4	10.5	11.1
18-25	17.3	17.3	30.3	34.1	15.4	14.9
26-49	13.8	12.7	33.6	33.2	12.7	11.7
50 or Older	6.8 ^a	9.3	*	*	6.6 ^a	9.1

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.15A Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Numbers in Thousands, 2003 and 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	16,334	16,989	3,334	3,561	13,000	13,428
18-25	5,462	5,807	1,200	1,313	4,262	4,494
26-49	8,584	8,857	1,756	1,882	6,828	6,976
50 or Older	2,287	2,325	377	366	1,910	1,959
MALE	11,105	11,892	1,715	1,909	9,390	9,982
18-25	3,469	3,784	596	623	2,873	3,160
26-49	5,935	6,431	865	1,076	5,070	5,355
50 or Older	1,701	1,677	254	*	1,447	1,467
FEMALE	5,229	5,097	1,619	1,651	3,610	3,446
18-25	1,993	2,023	605	689	1,389	1,334
26-49	2,649	2,426	891	805	1,758	1,621
50 or Older	586	648	123	157	463	491

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.15B Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress, Gender, and Age Group: Percentages, 2003 and 2004

Gender/Age	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	7.7	7.9	17.0	16.6	6.7	6.9
18-25	17.2	18.0	27.2	29.8	15.6	16.2
26-49	8.6	8.8	16.9	18.1	7.6	7.8
50 or Older	2.8	2.8	7.9	5.6	2.5	2.6
MALE	10.9	11.5	24.9	23.9	9.8	10.4
18-25	21.8	23.4	35.2	38.0	20.2	21.7
26-49	12.1	13.0	25.2	27.3	11.1	11.8
50 or Older	4.6	4.4	14.4	*	4.1	4.1
FEMALE	4.7	4.6	12.7	12.3	3.7	3.5
18-25	12.6	12.6	22.2	24.9	10.6	10.1
26-49	5.2	4.8	12.8	12.5	4.0	3.7
50 or Older	1.3	1.4	4.1	3.7	1.1	1.2

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).