

Table 1.19A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	110,205	110,057	34,993	34,807	19,470	19,071
12	503	443	255	265	112	112
13	811	810	516	509	211	204
14	1,086	1,104	772	777	349	395
15	1,414	1,489	1,043	1,054	551	547
16	1,835	1,750	1,391	1,277	780 ^b	638
17	1,981	1,970	1,478	1,419	807	778
18	2,464	2,456	1,761	1,782	1,042	974
19	2,265	2,229	1,551	1,520	914	897
20	2,497	2,354	1,623	1,520	967	851
21	2,415	2,541	1,372	1,492	812	885
22	2,570	2,529	1,347	1,412	786	826
23	2,538	2,335	1,287 ^a	1,111	719 ^a	605
24	2,265	2,421	1,095	1,117	626	655
25	2,181	2,178	939	959	569	546
26-29	8,555	9,210	3,483	3,603	1,982	2,028
30-34	11,463	10,693	3,345	3,080	1,771	1,837
35-39	12,903	12,434	3,141	2,953	1,757	1,500
40-44	14,821	14,723	3,175	3,272	1,831	1,705
45-49	13,349	13,295	2,691	2,543	1,463	1,465
50-54	10,389	10,917	1,474	1,743	784	935
55-59	5,716	6,264	663	830	298	427
60-64	2,798	3,032	336	248	127	140
65 or Older	3,387	2,879	254	321	209	122

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.19B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	46.4	45.8	14.7	14.5	8.2	7.9
12	12.2	11.2	6.2	6.7	2.7	2.8
13	18.7	18.4	11.9	11.6	4.9	4.6
14	26.3	25.2	18.7	17.8	8.5	9.0
15	34.2	34.7	25.2	24.6	13.3	12.7
16	43.8	42.5	33.2	31.0	18.6 ^b	15.5
17	48.4	48.4	36.1	34.9	19.7	19.1
18	53.5	53.4	38.2	38.8	22.6	21.2
19	58.3	56.6	39.9	38.6	23.5	22.8
20	62.0	59.0	40.3	38.1	24.0	21.3
21	61.6	62.3	35.0	36.6	20.7	21.7
22	64.0	62.9	33.5	35.1	19.6	20.5
23	63.4 ^a	59.5	32.2 ^a	28.3	18.0 ^a	15.4
24	62.3	59.8	30.1	27.6	17.2	16.2
25	60.1	60.5	25.9	26.7	15.7	15.2
26-29	57.9	60.0	23.6	23.5	13.4	13.2
30-34	56.8	54.5	16.6	15.7	8.8	9.4
35-39	61.7	59.4	15.0	14.1	8.4	7.2
40-44	65.3	64.9	14.0	14.4	8.1	7.5
45-49	62.3	61.8	12.6	11.8	6.8	6.8
50-54	52.0	56.3	7.4	9.0	3.9	4.8
55-59	38.3	38.2	4.4	5.1	2.0	2.6
60-64	23.8	24.2	2.9	2.0	1.1	1.1
65 or Older	9.9	8.3	0.7	0.9	0.6	0.4

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.20A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	96,611	96,772	25,231	25,451	14,638	14,576
12	59	64	38	42	21	15
13	225	241	159	188	62	81
14	578	560	459	453	218	217
15	947	964	759	740	378	412
16	1,434	1,358	1,114	1,032	638 ^a	525
17	1,645	1,602	1,216	1,201	654	659
18	2,109	2,047	1,459	1,433	897	790
19	1,953	1,975	1,290	1,294	795	748
20	2,255	2,116	1,385	1,315	799	744
21	2,200	2,303	1,168	1,207	699	736
22	2,342	2,270	1,135	1,156	668	692
23	2,276 ^a	2,065	1,001	881	574	494
24	2,015 ^a	2,216	857	918	506	544
25	1,954	1,990	749	758	462	448
26-29	7,547	8,093	2,556	2,764	1,543	1,651
30-34	10,287	9,573	2,488 ^a	2,114	1,396	1,265
35-39	11,840	11,274	2,113	2,058	1,256	1,083
40-44	13,751	13,650	2,227	2,152	1,357	1,297
45-49	12,349	12,325	1,609	1,769	884	1,093
50-54	9,488	10,194	910	1,299	489	737
55-59	4,896	5,456	418	515	258	273
60-64	2,225	2,468	74	68	51	26
65 or Older	2,236	1,969	48	94	33	45

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.20B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	40.6	40.2	10.6	10.6	6.2	6.1
12	1.4	1.6	0.9	1.1	0.5	0.4
13	5.2	5.5	3.7	4.3	1.4	1.8
14	14.0	12.8	11.1	10.4	5.3	5.0
15	22.9	22.4	18.4	17.2	9.2	9.6
16	34.2	33.0	26.6	25.1	15.2 ^a	12.8
17	40.2	39.4	29.7	29.5	16.0	16.2
18	45.8	44.5	31.7	31.2	19.5	17.2
19	50.2	50.2	33.2	32.9	20.5	19.0
20	56.0	53.0	34.4	33.0	19.8	18.6
21	56.1	56.5	29.8	29.6	17.8	18.0
22	58.3	56.5	28.3	28.8	16.6	17.2
23	56.8 ^a	52.6	25.0	22.4	14.3	12.6
24	55.4	54.8	23.6	22.7	13.9	13.4
25	53.8	55.3	20.6	21.1	12.7	12.4
26-29	51.1	52.7	17.3	18.0	10.4	10.8
30-34	50.9	48.8	12.3	10.8	6.9	6.4
35-39	56.6	53.8	10.1	9.8	6.0	5.2
40-44	60.6	60.2	9.8	9.5	6.0	5.7
45-49	57.7	57.3	7.5	8.2	4.1	5.1
50-54	47.5 ^a	52.6	4.5	6.7	2.4	3.8
55-59	32.8	33.3	2.8	3.1	1.7	1.7
60-64	18.9	19.7	0.6	0.5	0.4	0.2
65 or Older	6.5	5.7	0.1	0.3	0.1	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.21A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	34,891	34,153	5,908	5,658	2,281	2,021
12	8	4	7	3	2	*
13	22	17	9	13	7	5
14	62	54	41	24	14	7
15	99	103	74	68	28	26
16	187	177	129	130	42	26
17	270	256	178	173	60	63
18	445	409	285	258	115	80
19	468	465	283	260	92	87
20	567	548	278	317	67	92
21	610	705	271 ^a	364	81 ^a	125
22	678	763	273	316	104	97
23	728	673	272	227	82	81
24	660	707	233	224	78	74
25	598	624	203	167	77	55
26-29	2,590	2,519	769	780	263	230
30-34	3,722 ^a	3,178	650	612	256	252
35-39	5,454	4,989	627	486	323	205
40-44	6,781	6,709	591	659	264	235
45-49	5,440	5,487	419	319	144	157
50-54	3,614	3,809	255	163	167	56
55-59	1,208	1,268	47	43	*	34
60-64	410	317	*	29	*	22
65 or Older	269	374	14	23	14	12

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.21B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	14.7	14.2	2.5	2.4	1.0	0.8
12	0.2	0.1	0.2	0.1	0.1	*
13	0.5	0.4	0.2	0.3	0.2	0.1
14	1.5	1.2	1.0	0.6	0.3	0.2
15	2.4	2.4	1.8	1.6	0.7	0.6
16	4.5	4.3	3.1	3.1	1.0	0.6
17	6.6	6.3	4.3	4.3	1.5	1.5
18	9.7	8.9	6.2	5.6	2.5	1.7
19	12.0	11.8	7.3	6.6	2.4	2.2
20	14.1	13.7	6.9	8.0	1.7	2.3
21	15.5	17.3	6.9 ^a	8.9	2.1	3.1
22	16.9	19.0	6.8	7.9	2.6	2.4
23	18.2	17.1	6.8	5.8	2.1	2.1
24	18.2	17.5	6.4	5.5	2.1	1.8
25	16.5	17.4	5.6	4.6	2.1	1.5
26-29	17.5	16.4	5.2	5.1	1.8	1.5
30-34	18.4	16.2	3.2	3.1	1.3	1.3
35-39	26.1	23.8	3.0	2.3	1.5	1.0
40-44	29.9	29.6	2.6	2.9	1.2	1.0
45-49	25.4	25.5	2.0	1.5	0.7	0.7
50-54	18.1	19.7	1.3	0.8	0.8	0.3
55-59	8.1	7.7	0.3	0.3	*	0.2
60-64	3.5	2.5	*	0.2	*	0.2
65 or Older	0.8	1.1	0.0	0.1	0.0	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.22A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	7,949	7,840	1,406	1,304	604	467
12	*	1	*	1	*	*
13	2	3	2	3	2	3
14	18	14	11	8	2	4
15	28	24	18	11	7	2
16	53	34	33	21	14 ^a	2
17	57	48	31	19	7	6
18	91	86	40	45	18	6
19	114	117	44	41	7	12
20	148	108	40	20	8	11
21	128	159	28 ^a	58	6 ^a	23
22	170	189	29	29	1	5
23	187	156	29	31	6	11
24	184	163	37	25	5	6
25	171	157	29	13	4	6
26-29	776	681	151	159	60	45
30-34	1,152	918	165	129	82	66
35-39	1,547	1,389	200	192	113	56
40-44	1,515	1,514	224	227	122	90
45-49	990	1,163	172	111	50	65
50-54	457	544	113	117	90	34
55-59	124	219	11	14	*	14
60-64	5	44	*	7	*	*
65 or Older	32	108	*	23	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.22B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	3.3	3.3	0.6	0.5	0.3	0.2
12	*	0.0	*	0.0	*	*
13	0.0	0.1	0.0	0.1	0.0	0.1
14	0.4	0.3	0.3	0.2	0.1	0.1
15	0.7	0.6	0.4	0.3	0.2	0.0
16	1.3	0.8	0.8	0.5	0.3 ^a	0.0
17	1.4	1.2	0.8	0.5	0.2	0.1
18	2.0	1.9	0.9	1.0	0.4	0.1
19	2.9	3.0	1.1	1.0	0.2	0.3
20	3.7	2.7	1.0	0.5	0.2	0.3
21	3.3	3.9	0.7 ^a	1.4	0.1 ^a	0.6
22	4.2	4.7	0.7	0.7	0.0	0.1
23	4.7	4.0	0.7	0.8	0.2	0.3
24	5.1	4.0	1.0	0.6	0.1	0.1
25	4.7	4.4	0.8	0.4	0.1	0.2
26-29	5.3	4.4	1.0	1.0	0.4	0.3
30-34	5.7	4.7	0.8	0.7	0.4	0.3
35-39	7.4	6.6	1.0	0.9	0.5	0.3
40-44	6.7	6.7	1.0	1.0	0.5	0.4
45-49	4.6	5.4	0.8	0.5	0.2	0.3
50-54	2.3	2.8	0.6	0.6	0.4	0.2
55-59	0.8	1.3	0.1	0.1	*	0.1
60-64	0.0	0.4	*	0.1	*	*
65 or Older	0.1	0.3	*	0.1	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.23A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	34,363	34,333	3,936	3,878	1,042	929
12	44	24	21	14	7	8
13	63	78	41	59	12	18
14	140	127	101	86	39	22
15	217	205	127	142	47	26
16	321	284	207	204	70	59
17	465	441	282	248	84	77
18	667	579	366	320	108	81
19	780 ^b	622	319	292	87	71
20	963 ^b	753	414 ^b	299	95	75
21	949	964	279	308	55	74
22	1,062	1,054	220	251	68	53
23	1,050	970	207	187	55	53
24	998	1,004	183	161	46	43
25	910	910	123	126	30	26
26-29	3,406	3,695	475	491	125	122
30-34	3,679	3,601	273	296	40	46
35-39	4,305	3,936	102	178	41	42
40-44	5,300	4,799	69	130	33	26
45-49	4,486	4,598	56	40	*	6
50-54	3,158 ^a	3,990	70	5	*	*
55-59	1,006	1,180	*	18	*	*
60-64	287	262	*	*	*	*
65 or Older	109	257	*	24	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.23B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	14.5	14.3	1.7	1.6	0.4	0.4
12	1.1	0.6	0.5	0.3	0.2	0.2
13	1.5	1.8	0.9	1.3	0.3	0.4
14	3.4	2.9	2.4	2.0	0.9	0.5
15	5.2	4.8	3.1	3.3	1.1 ^a	0.6
16	7.7	6.9	4.9	4.9	1.7	1.4
17	11.4	10.9	6.9	6.1	2.1	1.9
18	14.5	12.6	7.9	7.0	2.3	1.8
19	20.1 ^b	15.8	8.2	7.4	2.2	1.8
20	23.9 ^b	18.9	10.3 ^b	7.5	2.4	1.9
21	24.2	23.6	7.1	7.5	1.4	1.8
22	26.4	26.2	5.5	6.2	1.7	1.3
23	26.2	24.7	5.2	4.8	1.4	1.4
24	27.5	24.8	5.0	4.0	1.3	1.1
25	25.1	25.3	3.4	3.5	0.8	0.7
26-29	23.1	24.1	3.2	3.2	0.8	0.8
30-34	18.2	18.3	1.4	1.5	0.2	0.2
35-39	20.6	18.8	0.5	0.8	0.2	0.2
40-44	23.4	21.2	0.3	0.6	0.1	0.1
45-49	20.9	21.4	0.3	0.2	*	0.0
50-54	15.8 ^b	20.6	0.4	0.0	*	*
55-59	6.7	7.2	*	0.1	*	*
60-64	2.4	2.1	*	*	*	*
65 or Older	0.3	0.7	*	0.1	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.24A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	22,995	22,798	2,075	2,255	570	638
12	289	291	107	145	44	53
13	461	461	231	195	76	47
14	473	531	235	246	57	64
15	489	554	193	241	60	73
16	485	481	186	172	48	38
17	473	444	166	170	35	37
18	577	503	171	161	35	45
19	524	476	109	116	16	20
20	632 ^a	513	100	99	28	15
21	584	616	60 ^a	104	4	10
22	571	628	63	71	12	19
23	640	601	79	48	17	20
24	609	625	50	48	14	9
25	592	553	31	28	7	7
26-29	2,052	2,146	65	67	4	16
30-34	2,707	2,432	25 ^a	107	8	54
35-39	2,677	2,717	34	65	18	38
40-44	2,988	2,974	73	81	54	20
45-49	2,619	2,377	69	28	23	14
50-54	1,606	1,681	12	25	9	*
55-59	522	680	*	*	*	*
60-64	189	241	*	40	*	40
65 or Older	238	273	17	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.24B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	9.7	9.5	0.9	0.9	0.2	0.3
12	7.0	7.3	2.6 ^a	3.6	1.1	1.3
13	10.6	10.5	5.3	4.4	1.8 ^a	1.1
14	11.5	12.1	5.7	5.6	1.4	1.5
15	11.8	12.9	4.7	5.6	1.4	1.7
16	11.6	11.7	4.4	4.2	1.1	0.9
17	11.6	10.9	4.1	4.2	0.9	0.9
18	12.5	10.9	3.7	3.5	0.8	1.0
19	13.5	12.1	2.8	2.9	0.4	0.5
20	15.7 ^a	12.9	2.5	2.5	0.7	0.4
21	14.9	15.1	1.5 ^a	2.5	0.1	0.2
22	14.2	15.6	1.6	1.8	0.3	0.5
23	16.0	15.3	2.0	1.2	0.4	0.5
24	16.8	15.4	1.4	1.2	0.4	0.2
25	16.3	15.4	0.8	0.8	0.2	0.2
26-29	13.9	14.0	0.4	0.4	0.0	0.1
30-34	13.4	12.4	0.1 ^a	0.5	0.0	0.3
35-39	12.8	13.0	0.2	0.3	0.1	0.2
40-44	13.2	13.1	0.3	0.4	0.2	0.1
45-49	12.2	11.1	0.3	0.1	0.1	0.1
50-54	8.0	8.7	0.1	0.1	0.0	*
55-59	3.5	4.1	*	*	*	*
60-64	1.6	1.9	*	0.3	*	0.3
65 or Older	0.7	0.8	0.1	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.25A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	47,882	48,013	14,986	14,643	6,336	6,007
12	237	189	132	115	60	46
13	330	366	211	251	90	98
14	452	461	310	321	134	169
15	612	672	442	438	205	186
16	808	806	555	526	240 ^a	179
17	913	899	640	574	268	236
18	1,180	1,217	706	784	285	319
19	1,161	1,098	675	672	266	300
20	1,181	1,152	701	622	317	248
21	1,153	1,306	554 ^a	689	233 ^a	309
22	1,176 ^a	1,339	489 ^b	629	215	249
23	1,245	1,179	566	517	225	198
24	1,097	1,133	516	467	198	188
25	1,017	969	392	371	174	149
26-29	3,824	4,096	1,491	1,546	597	627
30-34	4,795	4,265	1,454	1,417	595	629
35-39	5,518	5,145	1,377	1,186	544	498
40-44	6,591	6,300	1,187	1,269	508	480
45-49	5,873	5,967	1,288	1,106	609	447
50-54	4,320	5,013	616	461	280	183
55-59	2,120	2,261	216	337	40 ^a	154
60-64	1,056	979	263	141	76	51
65 or Older	1,221	1,202	206	205	176	65

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.25B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	20.1	20.0	6.3	6.1	2.7	2.5
12	5.7	4.8	3.2	2.9	1.5	1.2
13	7.6	8.3	4.9	5.7	2.1	2.2
14	11.0	10.5	7.5	7.3	3.3	3.9
15	14.8	15.7	10.7	10.2	5.0	4.3
16	19.3	19.6	13.2	12.8	5.7 ^a	4.3
17	22.3	22.1	15.6	14.1	6.5	5.8
18	25.6	26.5	15.3	17.1	6.2	6.9
19	29.9	27.9	17.4	17.1	6.9	7.6
20	29.3	28.9	17.4	15.6	7.9	6.2
21	29.4	32.0	14.1 ^a	16.9	5.9	7.6
22	29.3 ^a	33.3	12.2 ^b	15.7	5.3	6.2
23	31.1	30.0	14.1	13.2	5.6	5.0
24	30.2	28.0	14.2 ^a	11.5	5.4	4.6
25	28.0	26.9	10.8	10.3	4.8	4.1
26-29	25.9	26.7	10.1	10.1	4.0	4.1
30-34	23.7	21.7	7.2	7.2	2.9	3.2
35-39	26.4	24.6	6.6	5.7	2.6	2.4
40-44	29.0	27.8	5.2	5.6	2.2	2.1
45-49	27.4	27.7	6.0	5.1	2.8	2.1
50-54	21.6 ^a	25.9	3.1	2.4	1.4	0.9
55-59	14.2	13.8	1.4	2.1	0.3 ^a	0.9
60-64	9.0	7.8	2.2	1.1	0.6	0.4
65 or Older	3.6	3.5	0.6	0.6	0.5	0.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.26A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	31,207	31,768	11,671	11,256	4,693	4,404
12	199	163	110	98	48	39
13	286	291	186	203	76	84
14	369	398	260	282	111	153
15	521	547	371	349	170	153
16	654	698	458	442	191 ^a	142
17	768	783	533	490	204	179
18	987	1,083	618	657	226	244
19	960	925	543	546	208	219
20	1,014	990	583	499	270 ^a	194
21	929 ^a	1,114	450 ^a	564	170 ^a	236
22	944 ^a	1,094	393 ^a	490	167	196
23	1,009	972	484	412	174	154
24	895	909	417	396	146	154
25	775	725	315	269	125	107
26-29	2,947	3,095	1,155	1,173	389	401
30-34	3,342	3,134	1,095	1,073	491	440
35-39	3,360	3,135	1,033	850	388	322
40-44	3,615	3,663	866	932	316	351
45-49	3,230	3,318	949	771	449	331
50-54	2,079	2,455	394	314	175	149
55-59	970	1,160	120	277	17	64
60-64	648	433	189	46	51	26
65 or Older	704	683	149	122	131	65

*Low precision; no estimate reported.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.26B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	13.1	13.2	4.9	4.7	2.0	1.8
12	4.8	4.1	2.7	2.5	1.2	1.0
13	6.6	6.6	4.3	4.6	1.8	1.9
14	8.9	9.1	6.3	6.5	2.7	3.5
15	12.6	12.7	9.0	8.1	4.1	3.6
16	15.6	17.0	10.9	10.7	4.6	3.5
17	18.8	19.2	13.0	12.1	5.0	4.4
18	21.4	23.6	13.4	14.3	4.9	5.3
19	24.7	23.5	14.0	13.9	5.3	5.6
20	25.2	24.8	14.5	12.5	6.7 ^a	4.9
21	23.7 ^a	27.3	11.5	13.8	4.3	5.8
22	23.5 ^a	27.2	9.8 ^a	12.2	4.2	4.9
23	25.2	24.7	12.1	10.5	4.3	3.9
24	24.6	22.5	11.5	9.8	4.0	3.8
25	21.4	20.1	8.7	7.5	3.5	3.0
26-29	20.0	20.2	7.8	7.6	2.6	2.6
30-34	16.5	16.0	5.4	5.5	2.4	2.2
35-39	16.1	15.0	4.9	4.1	1.9	1.5
40-44	15.9	16.2	3.8	4.1	1.4	1.5
45-49	15.1	15.4	4.4	3.6	2.1	1.5
50-54	10.4	12.7	2.0	1.6	0.9	0.8
55-59	6.5	7.1	0.8	1.7	0.1	0.4
60-64	5.5	3.5	1.6 ^a	0.4	0.4	0.2
65 or Older	2.1	2.0	0.4	0.4	0.4	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.27A Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	71,128	70,657	20,305	19,658	8,849	8,247
12	471	411	229	236	95	101
13	695	711	416	417	168	146
14	805	841	519	514	202	239
15	948	1,058	600	641	285	261
16	1,143	1,106	744	689	309	250
17	1,258	1,229	844	772	358	329
18	1,552	1,571	968	1,015	427	412
19	1,490	1,405	878	840	368	392
20	1,630	1,486	966 ^a	818	427 ^a	335
21	1,601	1,721	762 ^a	926	296 ^a	391
22	1,700	1,774	718	824	325	353
23	1,731	1,599	780 ^a	666	318	260
24	1,574	1,628	689	613	278	277
25	1,492	1,445	505	499	227	198
26-29	5,592	6,055	2,062	2,038	859	835
30-34	7,589 ^a	6,856	1,933	1,912	773	870
35-39	8,810	8,257	1,779	1,675	807	694
40-44	9,817	9,771	1,715	1,815	821	726
45-49	8,738	8,601	1,615	1,329	768	565
50-54	6,527	7,032	839	610	434	233
55-59	2,990	3,238	263	362	40 ^a	188
60-64	1,404	1,244	263	209	76	113
65 or Older	1,573	1,618	220	239	190	77

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^aDifference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.27B Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2003 and 2004

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	29.9	29.4	8.5	8.2	3.7	3.4
12	11.4	10.3	5.6	5.9	2.3	2.6
13	16.0	16.2	9.6	9.5	3.9	3.3
14	19.5	19.2	12.6	11.8	4.9	5.5
15	22.9	24.6	14.5	14.9	6.9	6.1
16	27.3	26.9	17.8	16.7	7.4	6.1
17	30.7	30.2	20.6	19.0	8.7	8.1
18	33.7	34.2	21.0	22.1	9.3	9.0
19	38.3	35.7	22.6	21.3	9.5	10.0
20	40.5	37.2	24.0 ^a	20.5	10.6 ^a	8.4
21	40.8	42.2	19.4 ^a	22.7	7.5 ^a	9.6
22	42.3	44.1	17.9	20.5	8.1	8.8
23	43.2	40.7	19.5	17.0	7.9	6.6
24	43.3	40.2	19.0 ^b	15.2	7.7	6.8
25	41.1	40.2	13.9	13.9	6.3	5.5
26-29	37.9	39.5	14.0	13.3	5.8	5.4
30-34	37.6	34.9	9.6	9.7	3.8	4.4
35-39	42.1	39.4	8.5	8.0	3.9	3.3
40-44	43.3	43.1	7.6	8.0	3.6	3.2
45-49	40.8	40.0	7.5	6.2	3.6	2.6
50-54	32.6	36.3	4.2	3.1	2.2	1.2
55-59	20.0	19.7	1.8	2.2	0.3 ^a	1.1
60-64	11.9	9.9	2.2	1.7	0.6	0.9
65 or Older	4.6	4.7	0.6	0.7	0.6	0.2

*Low precision; no estimate reported.

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.