

Table 6.1C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004

Age Category	Total		GENDER			
			Male		Female	
	2003	2004	2003	2004	2003	2004
TOTAL	432	872	264	618	345	601
18	37	52	27	32	25	40
19	37	53	22	37	28	37
20	34	64	23	46	25	47
21	33	80	22	53	24	57
22	31	54	20	33	23	43
23	31	43	20	27	25	34
24	30	51	23	26	24	44
25	30	50	16	27	24	42
26-29	120	178	74	135	94	114
30-34	116	218	69	141	93	160
35-39	126	211	73	138	109	153
40-44	139	257	80	184	107	178
45-49	146	209	96	155	111	136
50-54	180	496	121	*	138	243
55-59	126	294	90	224	93	195
60-64	101	177	71	38	73	166
65 or Older	195	395	112	163	156	357

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.1D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2003 and 2004

Age Category	Total		GENDER			
			Male		Female	
	2003	2004	2003	2004	2003	2004
TOTAL	0.20	0.41	0.26	0.60	0.31	0.54
18	0.80	1.18	1.08	1.38	1.17	2.02
19	0.95	1.33	1.18	1.70	1.37	2.03
20	0.86	1.64	1.14	2.38	1.26	2.45
21	0.85	1.85	1.16	2.46	1.22	2.62
22	0.78	1.33	0.99	1.61	1.15	2.13
23	0.77	1.10	1.03	1.40	1.19	1.75
24	0.84	1.24	1.21	1.30	1.33	2.07
25	0.84	1.41	0.90	1.70	1.33	2.11
26-29	0.81	1.16	0.97	1.69	1.30	1.53
30-34	0.58	1.11	0.71	1.49	0.89	1.58
35-39	0.60	1.05	0.70	1.42	1.03	1.48
40-44	0.61	1.12	0.72	1.64	0.93	1.52
45-49	0.68	0.95	0.92	1.41	1.01	1.23
50-54	0.90	2.54	1.24	*	1.35	2.28
55-59	0.85	1.74	1.22	2.66	1.23	2.28
60-64	0.86	1.49	1.30	0.63	1.15	2.87
65 or Older	0.57	1.14	0.77	1.11	0.79	1.79

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.1P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2004 Versus 2003

Age Category	Total		GENDER	
			Male	Female
	2003	2004	2003	2004
TOTAL	0.1052		0.1422	0.4026
18	0.9316		0.3395	0.4537
19	0.3962		0.9279	0.4791
20	0.4645		0.8087	0.4701
21	0.2046		0.3098	0.4014
22	0.7243		0.7715	0.4994
23	0.0947		0.3721	0.1854
24	0.0132		0.0112	0.1777
25	0.5033		0.5788	0.9611
26-29	0.9139		0.3983	0.4983
30-34	0.0440		0.1243	0.1525
35-39	0.5035		0.6330	0.5897
40-44	0.8016		0.4693	0.2941
45-49	0.2310		0.8192	0.0412
50-54	0.3809		*	0.8440
55-59	0.0095		0.0440	0.1043
60-64	0.3033		0.1348	0.0503
65 or Older	0.6433		0.5072	0.3892

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.2C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	432	872	100	164	280	482	307	713
GENDER								
Male	264	618	64	105	175	337	194	512
Female	345	601	71	124	223	335	238	497
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	415	736	88	148	269	456	287	541
White	388	670	76	125	243	401	275	517
Black or African American	143	206	36	54	98	150	88	137
American Indian or Alaska Native	21	*	5	*	18	*	*	*
Native Hawaiian or Other Pacific Islander	15	*	*	*	*	*	*	*
Asian	83	137	19	50	73	76	21	*
Two or More Races	48	36	10	*	22	20	*	*
Hispanic or Latino	151	485	41	77	112	180	103	*
EDUCATION								
< High School	223	519	46	74	138	184	166	484
High School Graduate	244	465	54	85	159	286	181	362
Some College	212	369	52	103	153	212	133	273
College Graduate	186	324	31	49	139	248	130	215
CURRENT EMPLOYMENT								
Full-Time	291	438	62	103	224	341	182	267
Part-Time	151	261	47	88	108	214	101	113
Unemployed	85	183	31	76	64	132	*	*
Other ¹	271	645	45	62	135	203	222	611

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.2D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:
Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.20	0.41	0.32	0.51	0.28	0.48	0.38	0.86
GENDER								
Male	0.26	0.60	0.40	0.65	0.36	0.68	0.52	1.35
Female	0.31	0.54	0.45	0.78	0.44	0.66	0.54	1.11
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.22	0.39	0.34	0.56	0.31	0.54	0.38	0.70
White	0.26	0.44	0.39	0.63	0.36	0.61	0.43	0.79
Black or African American	0.60	0.85	0.84	1.24	0.83	1.22	1.16	1.77
American Indian or Alaska Native	1.92	*	2.57	*	3.20	*	*	*
Native Hawaiian or Other Pacific Islander	3.53	*	*	*	*	*	*	*
Asian	0.94	1.51	1.34	3.25	1.49	1.49	0.87	*
Two or More Races	2.22	1.90	2.80	*	2.18	2.83	*	*
Hispanic or Latino	0.58	1.81	0.73	1.34	0.78	1.20	1.81	*
EDUCATION								
< High School	0.59	1.43	0.68	1.09	0.94	1.32	1.02	3.10
High School Graduate	0.36	0.68	0.49	0.77	0.53	0.98	0.69	1.29
Some College	0.40	0.66	0.52	0.98	0.60	0.82	0.78	1.41
College Graduate	0.34	0.59	0.78	1.21	0.46	0.80	0.61	1.07
CURRENT EMPLOYMENT								
Full-Time	0.25	0.37	0.42	0.69	0.31	0.48	0.58	0.87
Part-Time	0.53	0.98	0.58	1.10	0.96	2.02	1.10	1.40
Unemployed	1.05	2.28	1.16	2.76	1.56	3.53	*	*
Other ¹	0.46	1.02	0.71	0.96	1.01	1.42	0.56	1.43

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.2P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.1052		0.7137		0.9705		0.0279	
GENDER								
Male	0.1422		0.5455		0.1952		0.2804	
Female	0.4026		0.9670		0.2171		0.0440	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.1986		0.9308		0.9125		0.0784	
White	0.1493		0.9470		0.4842		0.1411	
Black or African American	0.7694		0.6302		0.2525		0.2940	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.7244		0.3097		0.5019		*	
Two or More Races	0.1845		*		0.9834		*	
Hispanic or Latino	0.3406		0.3123		0.7423		*	
EDUCATION								
< High School	0.3207		0.7020		0.7330		0.2030	
High School Graduate	0.3566		0.7820		0.5892		0.3150	
Some College	0.7171		0.7030		0.0741		0.2553	
College Graduate	0.0560		0.3301		0.1510		0.2220	
CURRENT EMPLOYMENT								
Full-Time	0.2496		0.1172		0.1568		0.7228	
Part-Time	0.2624		0.2109		0.3540		0.7463	
Unemployed	0.1365		0.3806		0.2147		*	
Other ¹	0.0320		0.2548		0.6431		0.0177	

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.3C Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	432	872	100	164	280	482	307	713
GEOGRAPHIC DIVISION								
Northeast	194	397	48	74	117	180	133	326
New England	87	169	23	31	63	59	43	*
Middle Atlantic	173	360	41	68	99	170	125	290
Midwest	168	289	42	70	120	192	116	249
East North Central	144	222	36	58	106	164	100	167
West North Central	86	185	22	39	56	103	59	184
South	269	432	52	93	162	323	191	249
South Atlantic	198	312	38	64	116	249	143	179
East South Central	105	197	17	43	74	150	74	114
West South Central	150	224	32	51	86	139	103	133
West	220	575	57	88	155	242	164	524
Mountain	102	205	26	55	63	120	89	122
Pacific	196	536	50	68	142	202	137	*
COUNTY TYPE								
Large Metro	300	630	80	111	193	322	228	529
Small Metro	233	532	49	107	162	286	150	438
250K - 1 Mil. Pop.	189	410	39	88	127	197	127	348
< 250K Pop.	122	343	30	53	84	202	85	270
Nonmetro	183	295	33	59	109	189	133	192
Urbanized	135	191	22	44	68	143	100	90
Less Urbanized	113	206	24	47	78	100	84	164
Completely Rural	47	94	7	9	29	*	33	*

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.3D Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.20	0.41	0.32	0.51	0.28	0.48	0.38	0.86
GEOGRAPHIC DIVISION								
Northeast	0.47	0.97	0.84	1.30	0.62	0.95	0.82	1.99
New England	0.81	1.57	1.59	2.10	1.26	1.17	1.03	*
Middle Atlantic	0.58	1.19	0.99	1.61	0.71	1.22	1.05	2.39
Midwest	0.35	0.60	0.57	0.94	0.54	0.86	0.63	1.32
East North Central	0.43	0.66	0.70	1.13	0.68	1.05	0.78	1.27
West North Central	0.60	1.26	0.99	1.69	0.85	1.56	1.06	3.22
South	0.36	0.56	0.46	0.80	0.46	0.90	0.66	0.84
South Atlantic	0.50	0.77	0.68	1.13	0.63	1.34	0.90	1.10
East South Central	0.82	1.52	0.86	2.22	1.26	2.55	1.48	2.23
West South Central	0.65	0.95	0.83	1.30	0.78	1.24	1.24	1.57
West	0.46	1.18	0.77	1.17	0.67	1.04	0.95	2.94
Mountain	0.74	1.44	1.17	2.41	0.96	1.81	1.75	2.30
Pacific	0.58	1.56	0.97	1.29	0.85	1.21	1.13	*
COUNTY TYPE								
Large Metro	0.27	0.55	0.48	0.68	0.34	0.57	0.57	1.29
Small Metro	0.36	0.82	0.48	1.00	0.56	1.01	0.60	1.68
250K - 1 Mil. Pop.	0.45	0.97	0.58	1.27	0.65	1.05	0.81	2.12
< 250K Pop.	0.55	1.49	0.85	1.39	0.89	2.10	0.93	2.80
Nonmetro	0.51	0.80	0.68	1.16	0.73	1.22	0.83	1.19
Urbanized	0.84	1.24	0.92	1.92	1.00	2.22	1.45	1.37
Less Urbanized	0.70	1.12	1.08	1.90	1.17	1.29	1.15	2.02
Completely Rural	1.28	3.13	1.96	2.74	1.90	*	1.87	*

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.3P Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.1052		0.7137		0.9705		0.0279	
GEOGRAPHIC DIVISION								
Northeast	0.1644		0.8072		0.9706		0.1091	
New England	0.8161		0.6311		0.0286		*	
Middle Atlantic	0.0983		0.9650		0.3891		0.1557	
Midwest	0.0739		0.6076		0.3083		0.2079	
East North Central	0.0587		0.2591		0.0926		0.6026	
West North Central	0.5886		0.4264		0.4646		0.2003	
South	0.9972		0.9519		0.8977		0.9256	
South Atlantic	0.8751		0.9184		0.4532		0.4842	
East South Central	0.5972		0.8987		0.8782		0.4128	
West South Central	0.4735		0.9150		0.2337		0.9812	
West	0.4979		0.2275		0.3442		0.1335	
Mountain	0.3293		0.9705		0.1133		0.8734	
Pacific	0.7520		0.1354		0.0525		*	
COUNTY TYPE								
Large Metro	0.8642		0.3423		0.3386		0.2488	
Small Metro	0.0106		0.2892		0.3822		0.0165	
250K - 1 Mil. Pop.	0.1115		0.1793		0.8659		0.1301	
< 250K Pop.	0.0298		0.9462		0.2861		0.0475	
Nonmetro	0.8563		0.3513		0.7241		0.6714	
Urbanized	0.8732		0.8536		0.1238		0.1392	
Less Urbanized	0.5902		0.5285		0.1776		0.6820	
Completely Rural	0.9408		0.0305		*		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).