

**Table 6.67C Substance Use in the Past Year and Past Month among Persons Aged 12 or Older, by Past Year Major Depressive Episode (MDE):
Standard Errors of Numbers in Thousands, 2004**

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	741	300	708
Marijuana and Hashish	627	236	610
Cocaine	320	121	286
Crack	173	86	129
Heroin	88	39	79
Hallucinogens	209	87	188
LSD	70	23	67
PCP	32	8	31
Ecstasy	155	67	137
Inhalants	123	47	112
Nonmedical Use of Psychotherapeutics ⁴	476	218	412
Pain Relievers	401	196	340
OxyContin®	99	44	85
Tranquilizers	269	146	218
Stimulants	229	83	210
Methamphetamine	159	73	137
Sedatives	97	59	74
Illicit Drug Other Than Marijuana ³	569	245	503
PAST MONTH USE			
Daily Cigarette Use ⁵	898	325	910
Heavy Alcohol Use ⁶	592	171	589

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 12 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.67D Substance Use in the Past Year and Past Month among Persons Aged 12 or Older, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	0.31	1.38	0.30
Marijuana and Hashish	0.26	1.16	0.27
Cocaine	0.13	0.61	0.13
Crack	0.07	0.44	0.06
Heroin	0.04	0.20	0.04
Hallucinogens	0.09	0.43	0.09
LSD	0.03	0.12	0.03
PCP	0.01	0.04	0.01
Ecstasy	0.06	0.34	0.06
Inhalants	0.05	0.25	0.05
Nonmedical Use of Psychotherapeutics ⁴	0.20	1.08	0.19
Pain Relievers	0.17	0.97	0.15
OxyContin®	0.04	0.23	0.04
Tranquilizers	0.11	0.75	0.10
Stimulants	0.10	0.44	0.10
Methamphetamine	0.07	0.38	0.06
Sedatives	0.04	0.31	0.03
Illicit Drug Other Than Marijuana ³	0.24	1.16	0.22
PAST MONTH USE			
Daily Cigarette Use ⁵	0.37	1.45	0.38
Heavy Alcohol Use ⁶	0.25	0.86	0.26

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 12 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.68C Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	87	43	91
Marijuana and Hashish	77	35	76
Cocaine	27	11	25
Crack	10	6	8
Heroin	10	4	9
Hallucinogens	36	17	33
LSD	17	9	15
PCP	12	8	9
Ecstasy	23	12	21
Inhalants	45	21	44
Nonmedical Use of Psychotherapeutics ⁴	59	32	54
Pain Relievers	56	30	49
OxyContin®	18	9	15
Tranquilizers	29	14	27
Stimulants	28	14	26
Methamphetamine	17	8	15
Sedatives	15	7	13
Illicit Drug Other Than Marijuana ³	71	36	73
PAST MONTH USE			
Daily Cigarette Use ⁵	42	17	35
Heavy Alcohol Use ⁶	36	12	34

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004.

Table 6.68D Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	0.34	1.42	0.35
Marijuana and Hashish	0.31	1.29	0.31
Cocaine	0.11	0.47	0.11
Crack	0.04	0.26	0.04
Heroin	0.04	0.17	0.04
Hallucinogens	0.14	0.71	0.14
LSD	0.07	0.41	0.07
PCP	0.05	0.34	0.04
Ecstasy	0.09	0.53	0.09
Inhalants	0.18	0.86	0.18
Nonmedical Use of Psychotherapeutics ⁴	0.23	1.24	0.23
Pain Relievers	0.22	1.16	0.21
OxyContin®	0.07	0.40	0.07
Tranquilizers	0.12	0.61	0.12
Stimulants	0.11	0.62	0.11
Methamphetamine	0.07	0.36	0.07
Sedatives	0.06	0.31	0.06
Illicit Drug Other Than Marijuana ³	0.28	1.34	0.29
PAST MONTH USE			
Daily Cigarette Use ⁵	0.17	0.75	0.15
Heavy Alcohol Use ⁶	0.14	0.52	0.15

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004.

**Table 6.69C Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE):
Standard Errors of Numbers in Thousands, 2004**

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	734	296	703
Marijuana and Hashish	620	233	602
Cocaine	317	121	284
Crack	172	85	129
Heroin	88	39	79
Hallucinogens	207	85	186
LSD	68	22	65
PCP	30	3	30
Ecstasy	153	66	134
Inhalants	116	43	105
Nonmedical Use of Psychotherapeutics ⁴	465	213	403
Pain Relievers	390	192	332
OxyContin®	98	43	84
Tranquilizers	267	146	215
Stimulants	226	80	207
Methamphetamine	156	70	135
Sedatives	96	59	72
Illicit Drug Other Than Marijuana ³	560	241	493
PAST MONTH USE			
Daily Cigarette Use ⁵	898	324	906
Heavy Alcohol Use ⁶	593	171	587

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.69D Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	0.34	1.55	0.33
Marijuana and Hashish	0.29	1.30	0.29
Cocaine	0.15	0.70	0.14
Crack	0.08	0.50	0.07
Heroin	0.04	0.23	0.04
Hallucinogens	0.10	0.48	0.09
LSD	0.03	0.13	0.03
PCP	0.01	0.01	0.02
Ecstasy	0.07	0.37	0.07
Inhalants	0.05	0.25	0.05
Nonmedical Use of Psychotherapeutics ⁴	0.22	1.20	0.20
Pain Relievers	0.18	1.08	0.17
OxyContin®	0.05	0.25	0.04
Tranquilizers	0.12	0.84	0.11
Stimulants	0.10	0.47	0.11
Methamphetamine	0.07	0.41	0.07
Sedatives	0.04	0.34	0.04
Illicit Drug Other Than Marijuana ³	0.26	1.30	0.24
PAST MONTH USE			
Daily Cigarette Use ⁵	0.42	1.64	0.43
Heavy Alcohol Use ⁶	0.28	0.97	0.29

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).