

**Table 6.67A Substance Use in the Past Year and Past Month among Persons Aged 12 or Older, by Past Year Major Depressive Episode (MDE):
Numbers in Thousands, 2004**

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	36,093	5,561	30,243
Marijuana and Hashish	26,716	3,730	22,811
Cocaine	5,678	1,049	4,552
Crack	1,372	440	870
Heroin	441	118	313
Hallucinogens	3,997	736	3,238
LSD	634	135	494
PCP	181	26	150
Ecstasy	1,971	341	1,621
Inhalants	2,319	432	1,863
Nonmedical Use of Psychotherapeutics ⁴	14,349	3,003	11,224
Pain Relievers	11,026	2,306	8,636
OxyContin®	1,148	246	900
Tranquilizers	4,751	1,237	3,454
Stimulants	3,078	677	2,370
Methamphetamine	1,465	344	1,112
Sedatives	734	224	480
Illicit Drug Other Than Marijuana ³	19,572	3,681	15,703
PAST MONTH USE			
Daily Cigarette Use ⁵	38,227	4,927	33,026
Heavy Alcohol Use ⁶	16,963	1,784	15,106

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 12 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.67B Substance Use in the Past Year and Past Month among Persons Aged 12 or Older, by Past Year Major Depressive Episode (MDE): Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	15.0	28.8	13.8
Marijuana and Hashish	11.1	19.3	10.4
Cocaine	2.4	5.4	2.1
Crack	0.6	2.3	0.4
Heroin	0.2	0.6	0.1
Hallucinogens	1.7	3.8	1.5
LSD	0.3	0.7	0.2
PCP	0.1	0.1	0.1
Ecstasy	0.8	1.8	0.7
Inhalants	1.0	2.2	0.9
Nonmedical Use of Psychotherapeutics ⁴	6.0	15.6	5.1
Pain Relievers	4.6	11.9	3.9
OxyContin®	0.5	1.3	0.4
Tranquilizers	2.0	6.4	1.6
Stimulants	1.3	3.5	1.1
Methamphetamine	0.6	1.8	0.5
Sedatives	0.3	1.2	0.2
Illicit Drug Other Than Marijuana ³	8.1	19.1	7.2
PAST MONTH USE			
Daily Cigarette Use ⁵	15.9	25.5	15.1
Heavy Alcohol Use ⁶	7.1	9.2	6.9

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 12 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.68A Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Numbers in Thousands, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	5,300	905	4,316
Marijuana and Hashish	3,656	601	2,998
Cocaine	411	66	335
Crack	63	15	44
Heroin	48	9	37
Hallucinogens	753	162	574
LSD	157	36	117
PCP	72	21	47
Ecstasy	310	68	235
Inhalants	1,168	214	947
Nonmedical Use of Psychotherapeutics ⁴	2,225	481	1,707
Pain Relievers	1,865	406	1,425
OxyContin®	194	44	147
Tranquilizers	532	120	408
Stimulants	496	116	372
Methamphetamine	163	38	117
Sedatives	133	30	99
Illicit Drug Other Than Marijuana ³	3,269	631	2,587
PAST MONTH USE			
Daily Cigarette Use ⁵	828	171	628
Heavy Alcohol Use ⁶	671	96	559

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004.

Table 6.68B Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	21.0	40.7	19.1
Marijuana and Hashish	14.5	27.0	13.3
Cocaine	1.6	3.0	1.5
Crack	0.3	0.7	0.2
Heroin	0.2	0.4	0.2
Hallucinogens	3.0	7.3	2.5
LSD	0.6	1.6	0.5
PCP	0.3	0.9	0.2
Ecstasy	1.2	3.1	1.0
Inhalants	4.6	9.6	4.2
Nonmedical Use of Psychotherapeutics ⁴	8.8	21.6	7.5
Pain Relievers	7.4	18.3	6.3
OxyContin®	0.8	2.0	0.7
Tranquilizers	2.1	5.4	1.8
Stimulants	2.0	5.2	1.6
Methamphetamine	0.6	1.7	0.5
Sedatives	0.5	1.4	0.4
Illicit Drug Other Than Marijuana ³	13.0	28.3	11.4
PAST MONTH USE			
Daily Cigarette Use ⁵	3.3	7.7	2.8
Heavy Alcohol Use ⁶	2.7	4.3	2.5

*Low precision; no estimate reported.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004.

**Table 6.69A Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE):
Numbers in Thousands, 2004**

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	30,793	4,656	25,928
Marijuana and Hashish	23,060	3,129	19,813
Cocaine	5,267	983	4,217
Crack	1,309	425	826
Heroin	393	108	277
Hallucinogens	3,244	574	2,664
LSD	476	98	378
PCP	109	5	104
Ecstasy	1,662	273	1,386
Inhalants	1,151	218	916
Nonmedical Use of Psychotherapeutics ⁴	12,124	2,522	9,517
Pain Relievers	9,161	1,900	7,211
OxyContin®	953	201	752
Tranquilizers	4,219	1,117	3,046
Stimulants	2,582	561	1,998
Methamphetamine	1,302	306	994
Sedatives	600	193	381
Illicit Drug Other Than Marijuana ³	16,303	3,050	13,117
PAST MONTH USE			
Daily Cigarette Use ⁵	37,399	4,756	32,397
Heavy Alcohol Use ⁶	16,292	1,688	14,547

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.69B Substance Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Percentages, 2004

Substance	Total ¹	PAST YEAR MAJOR DEPRESSIVE EPISODE ²	
		Yes	No
PAST YEAR USE			
Illicit Drug ³	14.3	27.2	13.2
Marijuana and Hashish	10.7	18.3	10.1
Cocaine	2.4	5.8	2.1
Crack	0.6	2.5	0.4
Heroin	0.2	0.6	0.1
Hallucinogens	1.5	3.4	1.4
LSD	0.2	0.6	0.2
PCP	0.1	0.0	0.1
Ecstasy	0.8	1.6	0.7
Inhalants	0.5	1.3	0.5
Nonmedical Use of Psychotherapeutics ⁴	5.6	14.8	4.8
Pain Relievers	4.3	11.1	3.7
OxyContin®	0.4	1.2	0.4
Tranquilizers	2.0	6.5	1.6
Stimulants	1.2	3.3	1.0
Methamphetamine	0.6	1.8	0.5
Sedatives	0.3	1.1	0.2
Illicit Drug Other Than Marijuana ³	7.6	17.9	6.7
PAST MONTH USE			
Daily Cigarette Use ⁵	17.4	27.8	16.5
Heavy Alcohol Use ⁶	7.6	9.9	7.4

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. Therefore, estimates in the Total column are not comparable to similar estimates in other detailed tables. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year MDE information.

² Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

⁵ Past Month Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).