

Table 4.27B Past Year Initiation of Substance Use among Persons Aged 12 or Older, Persons Aged 12 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 or Older: Numbers in Thousands and Percentages, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ^{1,2} | | PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS ¹ | |
|---|---|-------|---|------|---|------|--|------|
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLCIT DRUG³ | 2,627 | 2,784 | 1.1 | 1.2 | 2.0 | 2.1 | 7.5 | 8.0 |
| Marijuana and Hashish | 1,973 | 2,142 | 0.8 | 0.9 | 1.4 | 1.5 | 7.8 | 8.4 |
| Cocaine | 986 | 998 | 0.4 | 0.4 | 0.5 | 0.5 | 16.7 | 17.6 |
| Crack | 269 | 215 | 0.1 | 0.1 | 0.1 | 0.1 | 19.1 | 16.5 |
| Heroin | 92 | 118 | 0.0 | 0.0 | 0.0 | 0.0 | * | 29.6 |
| Hallucinogens | 886 | 934 | 0.4 | 0.4 | 0.4 | 0.5 | 22.5 | 24.1 |
| LSD | 200 | 235 | 0.1 | 0.1 | 0.1 | 0.1 | 35.6 | 39.7 |
| PCP | 105 | 106 | 0.0 | 0.0 | 0.0 | 0.0 | 48.1 | * |
| Ecstasy | 642 | 607 | 0.3 | 0.3 | 0.3 | 0.3 | 29.9 | 31.7 |
| Inhalants | 871 | 857 | 0.4 | 0.4 | 0.4 | 0.4 | 41.9 | 38.0 |
| Nonmedical Use of Psychotherapeutics ⁴ | 2,583 | 2,836 | 1.1 | 1.2 | 1.3 | 1.5 | 17.2 ^a | 19.4 |
| Pain Relievers | 2,456 | 2,422 | 1.0 | 1.0 | 1.2 | 1.1 | 21.0 | 21.5 |
| OxyContin [®] | -- | 615 | -- | 0.3 | -- | 0.3 | -- | 50.7 |
| Tranquilizers | 1,071 | 1,180 | 0.5 | 0.5 | 0.5 | 0.5 | 21.2 | 23.3 |
| Stimulants | 715 | 793 | 0.3 | 0.3 | 0.3 | 0.4 | 26.0 | 27.2 |
| Methamphetamine | 260 | 318 | 0.1 | 0.1 | 0.1 | 0.1 | 19.7 | 22.1 |
| Sedatives | 194 | 240 | 0.1 | 0.1 | 0.1 | 0.1 | 23.3 | 32.5 |
| ILLCIT DRUG OTHER THAN MARIJUANA³ | 2,523 | 2,664 | 1.1 | 1.1 | 1.5 | 1.5 | 12.4 | 13.5 |
| CIGARETTES | 1,983 | 2,122 | 0.8 | 0.9 | 2.6 | 2.6 | 2.8 | 3.0 |
| Daily Cigarette Use ⁵ | 1,064 | 1,101 | 0.4 | 0.5 | 0.7 | 0.7 | -- | -- |
| SMOKELESS TOBACCO | 928 | 999 | 0.4 | 0.4 | 0.5 | 0.5 | 9.0 | 10.2 |
| CIGARS | 2,736 ^a | 3,058 | 1.2 | 1.3 | 1.8 | 2.0 | 10.8 | 11.8 |
| ALCOHOL | 4,082 ^a | 4,396 | 1.7 | 1.8 | 9.2 | 9.4 | 2.6 | 2.8 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.28B Past Year Initiation of Substance Use among Persons Aged 12 to 17, Persons Aged 12 to 17 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 12 to 17: Numbers in Thousands and Percentages, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ^{1,2} | | PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS ¹ | |
|---|---|-------|---|------|---|------|--|------|
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLCIT DRUG³ | 1,562 | 1,487 | 6.2 | 5.9 | 8.3 | 7.8 | 28.6 | 28.1 |
| Marijuana and Hashish | 1,219 | 1,252 | 4.9 | 5.0 | 5.7 | 5.8 | 32.5 | 34.2 |
| Cocaine | 282 | 274 | 1.1 | 1.1 | 1.1 | 1.1 | 64.3 | 66.7 |
| Crack | 76 ^a | 42 | 0.3 ^a | 0.2 | 0.3 ^a | 0.2 | * | * |
| Heroin | 25 | 31 | 0.1 | 0.1 | 0.1 | 0.1 | * | * |
| Hallucinogens | 417 | 437 | 1.7 | 1.7 | 1.7 | 1.8 | 53.5 | 58.0 |
| LSD | 96 | 99 | 0.4 | 0.4 | 0.4 | 0.4 | 64.1 | 62.8 |
| PCP | 59 | 43 | 0.2 | 0.2 | 0.2 | 0.2 | * | * |
| Ecstasy | 219 | 225 | 0.9 | 0.9 | 0.9 | 0.9 | 66.4 | 72.7 |
| Inhalants | 600 | 603 | 2.4 | 2.4 | 2.6 | 2.6 | 53.5 | 51.6 |
| Nonmedical Use of Psychotherapeutics ⁴ | 913 | 980 | 3.7 | 3.9 | 4.0 | 4.3 | 39.9 ^a | 44.1 |
| Pain Relievers | 890 | 908 | 3.6 | 3.6 | 3.9 | 3.9 | 46.3 | 48.7 |
| OxyContin [®] | -- | 141 | -- | 0.6 | -- | 0.6 | -- | 72.8 |
| Tranquilizers | 319 | 310 | 1.3 | 1.2 | 1.3 | 1.3 | 54.5 | 58.3 |
| Stimulants | 292 | 271 | 1.2 | 1.1 | 1.2 | 1.1 | 51.4 | 54.7 |
| Methamphetamine | 107 | 99 | 0.4 | 0.4 | 0.4 | 0.4 | 61.8 | 60.6 |
| Sedatives | 66 | 66 | 0.3 | 0.3 | 0.3 | 0.3 | 50.3 | 49.6 |
| ILLCIT DRUG OTHER THAN MARIJUANA³ | 1,205 | 1,171 | 4.8 | 4.6 | 5.8 | 5.6 | 36.0 | 35.8 |
| CIGARETTES | 1,226 | 1,294 | 4.9 | 5.1 | 6.6 | 6.8 | 25.9 | 27.9 |
| Daily Cigarette Use ⁵ | 439 | 417 | 1.8 | 1.7 | 1.9 | 1.7 | -- | -- |
| SMOKELESS TOBACCO | 487 | 447 | 1.9 | 1.8 | 2.1 | 1.9 | 47.0 ^a | 41.2 |
| CIGARS | 1,163 | 1,246 | 4.7 | 4.9 | 5.2 | 5.5 | 46.8 | 48.5 |
| ALCOHOL | 2,593 | 2,743 | 10.4 | 10.9 | 15.4 | 15.8 | 30.2 | 32.1 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.29B Past Year Initiation of Substance Use among Persons Aged 18 to 25, Persons Aged 18 to 25 At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 18 to 25: Numbers in Thousands and Percentages, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ^{1,2} | | PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS ¹ | |
|---|---|-------|---|------|---|------|--|------|
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLCIT DRUG³ | 742 | 818 | 2.3 | 2.5 | 5.6 | 5.9 | 6.8 | 7.5 |
| Marijuana and Hashish | 666 | 714 | 2.1 | 2.2 | 4.4 | 4.5 | 7.4 | 8.0 |
| Cocaine | 576 | 592 | 1.8 | 1.8 | 2.1 | 2.1 | 27.5 | 27.7 |
| Crack | 109 | 120 | 0.3 | 0.4 | 0.4 | 0.4 | 39.8 | 45.8 |
| Heroin | 42 | 46 | 0.1 | 0.1 | 0.1 | 0.1 | * | * |
| Hallucinogens | 425 | 412 | 1.3 | 1.3 | 1.7 | 1.6 | 20.1 | 21.2 |
| LSD | 98 | 112 | 0.3 | 0.3 | 0.4 | 0.4 | 28.9 | 35.3 |
| PCP | 41 | 49 | 0.1 | 0.2 | 0.1 | 0.2 | * | * |
| Ecstasy | 337 | 303 | 1.1 | 0.9 | 1.2 | 1.1 | 28.9 | 30.4 |
| Inhalants | 247 | 225 | 0.8 | 0.7 | 0.9 | 0.8 | 37.4 | 33.4 |
| Nonmedical Use of Psychotherapeutics ⁴ | 962 | 945 | 3.0 | 2.9 | 4.1 | 4.0 | 20.8 | 19.9 |
| Pain Relievers | 930 | 830 | 2.9 | 2.6 | 3.7 | 3.3 | 24.3 | 21.6 |
| OxyContin [®] | -- | 290 | -- | 0.9 | -- | 0.9 | -- | 51.6 |
| Tranquilizers | 457 | 422 | 1.4 | 1.3 | 1.6 | 1.5 | 27.2 | 25.5 |
| Stimulants | 240 ^b | 344 | 0.8 ^b | 1.1 | 0.8 ^b | 1.2 | 21.5 ^b | 29.1 |
| Methamphetamine | 92 ^a | 139 | 0.3 ^a | 0.4 | 0.3 ^a | 0.5 | 18.2 ^a | 27.0 |
| Sedatives | 42 | 53 | 0.1 | 0.2 | 0.1 | 0.2 | 26.5 | 32.5 |
| ILLCIT DRUG OTHER THAN MARIJUANA³ | 823 | 885 | 2.6 | 2.7 | 4.2 | 4.3 | 13.0 | 14.3 |
| CIGARETTES | 659 | 765 | 2.1 | 2.4 | 6.5 | 7.1 | 4.4 | 5.0 |
| Daily Cigarette Use ⁵ | 474 | 566 | 1.5 | 1.8 | 2.3 | 2.6 | -- | -- |
| SMOKELESS TOBACCO CIGARS | 353 ^a | 432 | 1.1 | 1.3 | 1.4 | 1.7 | 14.2 | 16.8 |
| ALCOHOL | 1,055 | 1,199 | 3.3 | 3.7 | 5.7 | 6.2 | 14.6 | 15.5 |
| | 1,430 | 1,484 | 4.5 | 4.6 | 25.8 | 25.0 | 5.8 | 5.9 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.30B Past Year Initiation of Substance Use among Persons Aged 26 or Older, Persons Aged 26 or Older At Risk for Initiation of Substance Use, and Past Year Substance Users Aged 26 or Older: Numbers in Thousands and Percentages, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG TOTAL POPULATION ¹ | | PERCENTAGE OF PAST YEAR INITIATES AMONG PERSONS AT RISK FOR INITIATION ^{1,2} | | PERCENTAGE OF PAST YEAR INITIATES AMONG PAST YEAR USERS ¹ | |
|---|---|------|---|------|---|------|--|------|
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLCIT DRUG³ | 324 | 479 | 0.2 | 0.3 | 0.3 | 0.5 | 1.7 | 2.6 |
| Marijuana and Hashish | 88 | 176 | 0.0 | 0.1 | 0.1 | 0.2 | 0.6 | 1.4 |
| Cocaine | 128 | 133 | 0.1 | 0.1 | 0.1 | 0.1 | 3.8 | 4.3 |
| Crack | 83 | 53 | 0.0 | 0.0 | 0.0 | 0.0 | 8.1 | 5.4 |
| Heroin | 25 | 40 | 0.0 | 0.0 | 0.0 | 0.0 | * | * |
| Hallucinogens | 44 | 85 | 0.0 | 0.0 | 0.0 | 0.1 | 4.2 | 7.2 |
| LSD | * | 24 | * | 0.0 | * | 0.0 | * | * |
| PCP | * | 14 | * | 0.0 | * | 0.0 | * | * |
| Ecstasy | 86 | 78 | 0.0 | 0.0 | 0.0 | 0.0 | * | * |
| Inhalants | 24 | 29 | 0.0 | 0.0 | 0.0 | 0.0 | * | * |
| Nonmedical Use of Psychotherapeutics ⁴ | 708 | 911 | 0.4 | 0.5 | 0.5 | 0.6 | 8.7 | 11.9 |
| Pain Relievers | 636 | 684 | 0.4 | 0.4 | 0.4 | 0.4 | 10.7 | 12.3 |
| OxyContin [®] | -- | 183 | -- | 0.1 | -- | 0.1 | -- | * |
| Tranquilizers | 295 ^a | 448 | 0.2 | 0.2 | 0.2 | 0.3 | 10.6 | 15.6 |
| Stimulants | 183 | 178 | 0.1 | 0.1 | 0.1 | 0.1 | 17.1 | 14.3 |
| Methamphetamine | 61 | 80 | 0.0 | 0.0 | 0.0 | 0.0 | * | 10.5 |
| Sedatives | 86 | 121 | 0.0 | 0.1 | 0.1 | 0.1 | * | * |
| ILLCIT DRUG OTHER THAN MARIJUANA³ | 496 | 608 | 0.3 | 0.3 | 0.4 | 0.5 | 4.6 | 6.0 |
| CIGARETTES | 98 | 63 | 0.1 | 0.0 | 0.2 | 0.1 | 0.2 | 0.1 |
| Daily Cigarette Use ⁵ | 150 | 118 | 0.1 | 0.1 | 0.2 | 0.1 | -- | -- |
| SMOKELESS TOBACCO CIGARS | 88 | 121 | 0.0 | 0.1 | 0.1 | 0.1 | 1.3 | 2.0 |
| CIGARS | 518 | 614 | 0.3 | 0.3 | 0.5 | 0.5 | 3.3 | 3.9 |
| ALCOHOL | 60 | 169 | 0.0 | 0.1 | 0.3 | 0.7 | 0.0 | 0.1 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² At Risk for Initiation is defined as persons who did not use the substance(s) in their lifetime or used the substance(s) for the first time in the past year.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁵ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.31A Past Year Initiation of Substance Use among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | | | | |
|--|---|-------|------------------|-------|--------------------|-------|
| | Total | | Male | | Female | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLICIT DRUG² | 2,627 | 2,784 | 1,137 | 1,173 | 1,490 | 1,611 |
| Marijuana and Hashish | 1,973 | 2,142 | 929 | 918 | 1,044 ^a | 1,224 |
| Cocaine | 986 | 998 | 548 | 590 | 438 | 408 |
| Crack | 269 | 215 | 135 | 102 | 134 | 112 |
| Heroin | 92 | 118 | 59 | 61 | 33 | 56 |
| Hallucinogens | 886 | 934 | 480 | 506 | 406 | 427 |
| LSD | 200 | 235 | 123 | 141 | 76 | 94 |
| PCP | 105 | 106 | 59 | 55 | 46 | 51 |
| Ecstasy | 642 | 607 | 328 | 336 | 314 | 271 |
| Inhalants | 871 | 857 | 457 | 434 | 414 | 422 |
| Nonmedical Use of Psychotherapeutics ³ | 2,583 | 2,836 | 995 | 1,189 | 1,588 | 1,648 |
| Pain Relievers | 2,456 | 2,422 | 1,007 | 1,093 | 1,448 | 1,329 |
| OxyContin [®] | -- | 615 | -- | 339 | -- | 276 |
| Tranquilizers | 1,071 | 1,180 | 415 | 452 | 656 | 729 |
| Stimulants | 715 | 793 | 272 | 341 | 442 | 452 |
| Methamphetamine | 260 | 318 | 130 | 142 | 129 | 176 |
| Sedatives | 194 | 240 | 66 | 79 | 128 | 160 |
| ILLICIT DRUG OTHER THAN MARIJUANA² | 2,523 | 2,664 | 1,083 | 1,224 | 1,441 | 1,439 |
| CIGARETTES | 1,983 | 2,122 | 919 | 973 | 1,063 | 1,150 |
| Daily Cigarette Use ⁴ | 1,064 | 1,101 | 483 ^a | 594 | 581 | 507 |
| SMOKELESS TOBACCO | 928 | 999 | 708 | 727 | 220 | 272 |
| CIGARS | 2,736 ^a | 3,058 | 1,482 | 1,589 | 1,254 ^a | 1,469 |
| ALCOHOL | 4,082 ^a | 4,396 | 1,827 | 1,985 | 2,255 | 2,411 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.32A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Gender: Numbers in Thousands, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | | | | |
|--|---|-------|-------|-------|--------|-------|
| | Total | | Male | | Female | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLICIT DRUG² | 1,699 | 1,617 | 781 | 763 | 919 | 854 |
| Marijuana and Hashish | 1,330 | 1,367 | 603 | 642 | 728 | 725 |
| Cocaine | 341 | 341 | 162 | 179 | 179 | 162 |
| Crack | 86 | 67 | 36 | 31 | 49 | 35 |
| Heroin | 28 | 35 | 14 | 20 | 15 | 16 |
| Hallucinogens | 467 | 483 | 209 | 239 | 258 | 244 |
| LSD | 114 | 119 | 57 | 69 | 57 | 50 |
| PCP | 63 | 48 | 32 | 22 | 31 | 27 |
| Ecstasy | 244 | 257 | 101 | 111 | 143 | 146 |
| Inhalants | 665 | 643 | 321 | 323 | 344 | 319 |
| Nonmedical Use of Psychotherapeutics ³ | 1,036 | 1,097 | 469 | 476 | 567 | 622 |
| Pain Relievers | 1,020 | 1,018 | 474 | 446 | 546 | 572 |
| OxyContin [®] | -- | 165 | -- | 83 | -- | 82 |
| Tranquilizers | 366 | 348 | 153 | 137 | 213 | 212 |
| Stimulants | 308 | 301 | 127 | 120 | 182 | 182 |
| Methamphetamine | 111 | 112 | 49 | 42 | 62 | 70 |
| Sedatives | 73 | 74 | 35 | 30 | 38 | 44 |
| ILLICIT DRUG OTHER THAN MARIJUANA² | 1,325 | 1,293 | 629 | 608 | 696 | 685 |
| CIGARETTES | 1,352 | 1,438 | 628 | 645 | 724 | 793 |
| Daily Cigarette Use ⁴ | 526 | 545 | 249 | 257 | 278 | 287 |
| SMOKELESS TOBACCO | 544 | 495 | 440 | 378 | 104 | 117 |
| CIGARS | 1,297 | 1,409 | 752 | 782 | 545 | 627 |
| ALCOHOL | 2,819 | 3,002 | 1,282 | 1,411 | 1,537 | 1,590 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 4.33A Past Year Initiation of Substance Use among Persons Aged 12 or Older Who Initiated Use Prior to the Age of 21, by Gender: Numbers in Thousands, 2003 and 2004

| Substance | NUMBER OF PAST YEAR INITIATES (1,000s) ¹ | | | | | |
|--|---|-------|-------|-------|------------------|-------|
| | Total | | Male | | Female | |
| | 2003 | 2004 | 2003 | 2004 | 2003 | 2004 |
| ILLICIT DRUG² | 2,126 | 2,128 | 970 | 955 | 1,156 | 1,173 |
| Marijuana and Hashish | 1,744 | 1,858 | 808 | 845 | 936 | 1,012 |
| Cocaine | 662 | 681 | 363 | 396 | 299 | 285 |
| Crack | 146 | 122 | 76 | 66 | 69 | 55 |
| Heroin | 49 | 56 | 30 | 33 | 19 | 23 |
| Hallucinogens | 756 | 731 | 392 | 398 | 364 | 334 |
| LSD | 174 | 185 | 102 | 109 | 72 | 75 |
| PCP | 92 | 75 | 51 | 44 | 41 | 31 |
| Ecstasy | 461 | 428 | 213 | 219 | 248 | 209 |
| Inhalants | 791 | 779 | 412 | 401 | 379 | 378 |
| Nonmedical Use of Psychotherapeutics ³ | 1,544 | 1,621 | 701 | 703 | 843 | 917 |
| Pain Relievers | 1,495 | 1,457 | 706 | 661 | 789 | 797 |
| OxyContin [®] | -- | 306 | -- | 163 | -- | 143 |
| Tranquilizers | 582 | 545 | 253 | 229 | 329 | 316 |
| Stimulants | 453 | 526 | 203 | 206 | 250 ^a | 320 |
| Methamphetamine | 166 | 175 | 90 | 71 | 76 | 104 |
| Sedatives | 91 | 97 | 43 | 42 | 48 | 55 |
| ILLICIT DRUG OTHER THAN MARIJUANA² | 1,766 | 1,778 | 842 | 849 | 924 | 929 |
| CIGARETTES | 1,758 | 1,928 | 824 | 897 | 934 | 1,030 |
| Daily Cigarette Use ⁴ | 808 | 887 | 398 | 471 | 410 | 417 |
| SMOKELESS TOBACCO | 750 | 767 | 593 | 585 | 157 | 182 |
| CIGARS | 1,964 ^a | 2,179 | 1,157 | 1,234 | 807 ^a | 945 |
| ALCOHOL | 3,644 | 3,818 | 1,660 | 1,790 | 1,984 | 2,028 |

*Low precision, no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Past Year Initiates are defined as persons who used the substance(s) for the first time in the 12 months prior to date of interview.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.⁴ Daily Cigarette Use is defined as ever smoking every day for at least 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.