

**Table 7.17A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	5,247	5,381	6,216	6,272	4,136	3,803	3,871	3,615
Marijuana and Hashish	4,266	4,508	5,185	5,005	2,999	2,948	2,189	2,114
Cocaine	1,096	948	646	735	316 <sup>a</sup>	167	222	170
Crack	219	154	133	216	123	31	129	65
Heroin	24	56	39	40	28	12	28	58
Hallucinogens	424	385	325	311	163	111	131	123
LSD	68	67	46	40	5	12	14	22
PCP	24	23	26	9	*	*	*	15
Ecstasy	158	165	156	178	111	73	44	34
Inhalants	125	102	132	190	97	84	216	261
Nonmedical Use of Psychotherapeutics <sup>3</sup>	1,697	1,680	1,670	1,786	1,330	1,132	1,639	1,410
Pain Relievers	1,281	1,271	1,228	1,296	874	740	1,310	1,096
OxyContin <sup>®</sup>	--	135	--	106	--	41	--	43
Tranquilizers	563	486	517	481	454	337	296	313
Stimulants	480	438	336	379	197	181	179	192
Methamphetamine	269	206	160	151	99	115	79	110
Sedatives	98	45	66	62	84	81	47	77
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	2,612	2,447	2,452	2,558	1,730 <sup>a</sup>	1,349	2,055	1,893

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.17B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLICIT DRUG<sup>2</sup></b>	32.5	32.2	16.5	16.5	6.3	5.7	3.3	3.0
Marijuana and Hashish	26.4	27.0	13.8	13.2	4.6	4.5	1.8	1.8
Cocaine	6.8	5.7	1.7	1.9	0.5 <sup>a</sup>	0.3	0.2	0.1
Crack	1.4	0.9	0.4	0.6	0.2	0.0	0.1	0.1
Heroin	0.1	0.3	0.1	0.1	0.0	0.0	0.0	0.0
Hallucinogens	2.6	2.3	0.9	0.8	0.2	0.2	0.1	0.1
LSD	0.4	0.4	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.1	0.1	0.1	0.0	*	*	*	0.0
Ecstasy	1.0	1.0	0.4	0.5	0.2	0.1	0.0	0.0
Inhalants	0.8	0.6	0.4	0.5	0.1	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics <sup>3</sup>	10.5	10.1	4.4	4.7	2.0	1.7	1.4	1.2
Pain Relievers	7.9	7.6	3.3	3.4	1.3	1.1	1.1	0.9
OxyContin <sup>®</sup>	--	0.8	--	0.3	--	0.1	--	0.0
Tranquilizers	3.5	2.9	1.4	1.3	0.7	0.5	0.2	0.3
Stimulants	3.0	2.6	0.9	1.0	0.3	0.3	0.2	0.2
Methamphetamine	1.7	1.2	0.4	0.4	0.2	0.2	0.1	0.1
Sedatives	0.6	0.3	0.2	0.2	0.1	0.1	0.0	0.1
<b>ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	16.2	14.7	6.5	6.7	2.7 <sup>a</sup>	2.0	1.7	1.6

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.18A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	421	441	913	815	428	374	1,049	1,044
Marijuana and Hashish	366	385	763	683	311	284	531	557
Cocaine	56	53	57	54	20	11	20	7
Crack	21	11	6 <sup>a</sup>	2	1	4	4	*
Heroin	7	9	8	4	0	*	3	2
Hallucinogens	89	77	86	69	27	19	57	46
LSD	22	11	16	13	1	6	12	9
PCP	10	7	8	3	*	1	2	*
Ecstasy	40	32	28	29	11	5	13	11
Inhalants	39	32	76	67	53	46	152	168
Nonmedical Use of Psychotherapeutics <sup>3</sup>	188	176	282	236	114	106	413	396
Pain Relievers	151	154	213	179	99	89	339	328
OxyContin <sup>®</sup>	--	32	--	18	--	8	--	8
Tranquilizers	68	40	78 <sup>a</sup>	47	20	19	61	56
Stimulants	66	46	74	71	23	12	58	49
Methamphetamine	29 <sup>a</sup>	11	12	27	10	3	18	15
Sedatives	12	3	14	4	6	5	15	13
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	242	247	396	341	181	156	598	584

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.18B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLICIT DRUG<sup>2</sup></b>	64.5	65.6	45.5 <sup>b</sup>	38.4	24.2	22.8	5.1	5.0
Marijuana and Hashish	56.1	57.3	38.0 <sup>b</sup>	32.2	17.6	17.3	2.6	2.7
Cocaine	8.5	8.0	2.9	2.6	1.2	0.7	0.1	0.0
Crack	3.2	1.6	0.3 <sup>a</sup>	0.1	0.1	0.2	0.0	*
Heroin	1.1	1.4	0.4	0.2	0.0	*	0.0	0.0
Hallucinogens	13.7	11.5	4.3	3.3	1.5	1.1	0.3	0.2
LSD	3.3	1.6	0.8	0.6	0.1	0.4	0.1	0.0
PCP	1.6	1.0	0.4	0.1	*	0.1	0.0	*
Ecstasy	6.1	4.7	1.4	1.4	0.6	0.3	0.1	0.1
Inhalants	6.0	4.7	3.8	3.1	3.0	2.8	0.7	0.8
Nonmedical Use of Psychotherapeutics <sup>3</sup>	28.8	26.1	14.1 <sup>a</sup>	11.1	6.5	6.4	2.0	1.9
Pain Relievers	23.1	23.0	10.6	8.5	5.6	5.4	1.6	1.6
OxyContin <sup>®</sup>	--	4.8	--	0.9	--	0.5	--	0.0
Tranquilizers	10.4 <sup>a</sup>	5.9	3.9 <sup>a</sup>	2.2	1.1	1.2	0.3	0.3
Stimulants	10.1	6.9	3.7	3.3	1.3	0.7	0.3	0.2
Methamphetamine	4.5 <sup>a</sup>	1.7	0.6	1.3	0.6	0.2	0.1	0.1
Sedatives	1.8	0.5	0.7	0.2	0.3	0.3	0.1	0.1
<b>ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	37.2	36.7	19.7 <sup>a</sup>	16.1	10.2	9.5	2.9	2.8

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.19A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	2,336	2,217	2,307	2,263	1,013	949	778	809
Marijuana and Hashish	2,015	1,960	1,990	1,883	872	805	522	548
Cocaine	425	408	196	207	42	29	34	47
Crack	23 <sup>a</sup>	46	21	19	7	5	4	11
Heroin	10	16	8	16	*	1	6	10
Hallucinogens	262	228	180	173	57	40	46	36
LSD	35	50	26	24	4	5	2	2
PCP	11	9	16	7	*	*	3	1
Ecstasy	95	109	81	84	28	22	26	16
Inhalants	58	59	35	55	22	4	19	25
Nonmedical Use of Psychotherapeutics <sup>3</sup>	766	763	623	654	229	244	295	298
Pain Relievers	585	567	480	509	175	186	246	242
OxyContin <sup>®</sup>	--	63	--	52	--	18	--	9
Tranquilizers	252	253	165	190	68	75	40	47
Stimulants	191	226	137	126	33	40	39	54
Methamphetamine	74	84	70	54	17	18	24	30
Sedatives	26	21	19	14	3	4	10	10
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	1,147	1,076	874	893	296	285	348	365

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.19B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLICIT DRUG<sup>2</sup></b>	48.8	45.7	27.5	26.9	16.1	15.3	6.3	6.4
Marijuana and Hashish	42.1	40.4	23.7	22.4	13.9	13.0	4.3	4.3
Cocaine	8.9	8.4	2.3	2.5	0.7	0.5	0.3	0.4
Crack	0.5 <sup>a</sup>	0.9	0.2	0.2	0.1	0.1	0.0	0.1
Heroin	0.2	0.3	0.1	0.2	*	0.0	0.1	0.1
Hallucinogens	5.5	4.7	2.1	2.1	0.9	0.6	0.4	0.3
LSD	0.7	1.0	0.3	0.3	0.1	0.1	0.0	0.0
PCP	0.2	0.2	0.2	0.1	*	*	0.0	0.0
Ecstasy	2.0	2.3	1.0	1.0	0.4	0.4	0.2	0.1
Inhalants	1.2	1.2	0.4	0.7	0.3	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics <sup>3</sup>	16.0	15.7	7.4	7.8	3.6	3.9	2.4	2.3
Pain Relievers	12.2	11.7	5.7	6.0	2.8	3.0	2.0	1.9
OxyContin <sup>®</sup>	--	1.3	--	0.6	--	0.3	--	0.1
Tranquilizers	5.3	5.2	2.0	2.3	1.1	1.2	0.3	0.4
Stimulants	4.0	4.7	1.6	1.5	0.5	0.6	0.3	0.4
Methamphetamine	1.5	1.7	0.8	0.6	0.3	0.3	0.2	0.2
Sedatives	0.5	0.4	0.2	0.2	0.0	0.1	0.1	0.1
<b>ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	24.0	22.2	10.4	10.6	4.7	4.6	2.8	2.9

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.20A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	2,490	2,724	2,997	3,194	2,694	2,480	2,043	1,762
Marijuana and Hashish	1,885	2,164	2,431	2,439	1,816	1,859	1,136	1,009
Cocaine	616	486	393	474	254	127	168	116
Crack	175	98	107	196	115	23	120	54
Heroin	7	31	23	21	28	11	19	46
Hallucinogens	73	80	60	68	78	52	28	42
LSD	11	5	4	2	*	*	*	11
PCP	3	8	2	*	*	*	*	13
Ecstasy	24	24	47	65	73	46	5	7
Inhalants	28	12	21	68	23	34	44	67
Nonmedical Use of Psychotherapeutics <sup>3</sup>	743	742	765	895	987	782	931	715
Pain Relievers	545	550	535	608	600	465	726	526
OxyContin <sup>®</sup>	--	40	--	36	--	15	--	27
Tranquilizers	243	193	274	244	366	242	195	210
Stimulants	224	166	125	181	141	129	81	88
Methamphetamine	166	111	78	70	72	94	36	64
Sedatives	61	21	33	44	75	72	21	53
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	1,222	1,125	1,181	1,323	1,254	909	1,110	945

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 7.20B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2003 and 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>ILLCIT DRUG<sup>2</sup></b>	23.3	24.4	11.0	11.6	4.7	4.2	2.4	2.0
Marijuana and Hashish	17.6	19.4	8.9	8.9	3.2	3.2	1.3	1.2
Cocaine	5.8	4.4	1.4	1.7	0.4	0.2	0.2	0.1
Crack	1.6	0.9	0.4	0.7	0.2	0.0	0.1	0.1
Heroin	0.1	0.3	0.1	0.1	0.0	0.0	0.0	0.1
Hallucinogens	0.7	0.7	0.2	0.2	0.1	0.1	0.0	0.0
LSD	0.1	0.0	0.0	0.0	*	*	*	0.0
PCP	0.0	0.1	0.0	*	*	*	*	0.0
Ecstasy	0.2	0.2	0.2	0.2	0.1	0.1	0.0	0.0
Inhalants	0.3	0.1	0.1	0.2	0.0	0.1	0.1	0.1
Nonmedical Use of Psychotherapeutics <sup>3</sup>	6.9	6.6	2.8	3.3	1.7	1.3	1.1	0.8
Pain Relievers	5.1	4.9	2.0	2.2	1.1	0.8	0.8	0.6
OxyContin <sup>®</sup>	--	0.4	--	0.1	--	0.0	--	0.0
Tranquilizers	2.3	1.7	1.0	0.9	0.6	0.4	0.2	0.2
Stimulants	2.1	1.5	0.5	0.7	0.2	0.2	0.1	0.1
Methamphetamine	1.6	1.0	0.3	0.3	0.1	0.2	0.0	0.1
Sedatives	0.6	0.2	0.1	0.2	0.1	0.1	0.0	0.1
<b>ILLCIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	11.4	10.1	4.3	4.8	2.2	1.6	1.3	1.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>3</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.