

**Table 6.28C Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004**

| Source of Payment <sup>1</sup>            | Total |      | AGE GROUP |      |       |      |             |      |
|-------------------------------------------|-------|------|-----------|------|-------|------|-------------|------|
|                                           |       |      | 18-25     |      | 26-49 |      | 50 or Older |      |
|                                           | 2003  | 2004 | 2003      | 2004 | 2003  | 2004 | 2003        | 2004 |
| Self or Family Member Living in Household | 150   | 296  | 34        | 59   | 102   | 218  | *           | *    |
| Family Member Not Living in Household     | 30    | 54   | 16        | 14   | 26    | 36   | 1           | *    |
| Private Health Insurance                  | 192   | 191  | 33        | 44   | 125   | 155  | *           | *    |
| Medicare                                  | 103   | 199  | 9         | 17   | 66    | 112  | *           | *    |
| Medicaid                                  | 92    | 128  | 13        | 28   | 68    | 90   | 51          | *    |
| Rehabilitation Program                    | 19    | 34   | 2         | 2    | 15    | 34   | *           | *    |
| Employer                                  | 70    | 68   | 11        | 16   | 54    | 49   | 38          | *    |
| VA or Other Military Program              | 52    | 66   | 6         | 7    | 38    | 40   | 35          | *    |
| Other Public Source                       | 49    | 141  | 9         | 18   | 48    | 49   | 3           | *    |
| Other Private Source                      | 20    | 23   | 8         | 8    | 17    | 22   | *           | *    |
| Free Treatment                            | 74    | 67   | 11        | 28   | 25    | 39   | *           | *    |

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.28D Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2003 and 2004**

| Source of Payment <sup>1</sup>            | Total |      | AGE GROUP |      |       |      |             |      |
|-------------------------------------------|-------|------|-----------|------|-------|------|-------------|------|
|                                           |       |      | 18-25     |      | 26-49 |      | 50 or Older |      |
|                                           | 2003  | 2004 | 2003      | 2004 | 2003  | 2004 | 2003        | 2004 |
| Self or Family Member Living in Household | 2.00  | 3.37 | 2.25      | 4.12 | 2.32  | 4.19 | *           | *    |
| Family Member Not Living in Household     | 0.49  | 0.84 | 1.53      | 1.27 | 0.73  | 1.00 | 0.09        | *    |
| Private Health Insurance                  | 2.15  | 2.69 | 2.55      | 3.35 | 2.56  | 3.78 | *           | *    |
| Medicare                                  | 1.59  | 2.76 | 0.89      | 1.57 | 1.71  | 2.84 | *           | *    |
| Medicaid                                  | 1.45  | 1.89 | 1.22      | 2.38 | 1.83  | 2.40 | 3.43        | *    |
| Rehabilitation Program                    | 0.31  | 0.53 | 0.17      | 0.19 | 0.42  | 0.94 | *           | *    |
| Employer                                  | 1.14  | 1.06 | 1.04      | 1.45 | 1.46  | 1.35 | 2.65        | *    |
| VA or Other Military Program              | 0.87  | 1.01 | 0.53      | 0.64 | 1.08  | 1.09 | 2.51        | *    |
| Other Public Source                       | 0.82  | 2.09 | 0.89      | 1.67 | 1.35  | 1.33 | 0.20        | *    |
| Other Private Source                      | 0.34  | 0.37 | 0.81      | 0.74 | 0.49  | 0.61 | *           | *    |
| Free Treatment                            | 1.19  | 1.03 | 1.06      | 2.36 | 0.71  | 1.08 | *           | *    |

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

**Table 6.28P Source of Payment for Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2004 Versus 2003**

| Source of Payment <sup>1</sup>            | Total  |      | AGE GROUP |      |        |      |             |      |
|-------------------------------------------|--------|------|-----------|------|--------|------|-------------|------|
|                                           |        |      | 18-25     |      | 26-49  |      | 50 or Older |      |
|                                           | 2003   | 2004 | 2003      | 2004 | 2003   | 2004 | 2003        | 2004 |
| Self or Family Member Living in Household | 0.3875 |      | 0.1700    |      | 0.1112 |      | *           |      |
| Family Member Not Living in Household     | 0.5353 |      | 0.0115    |      | 0.5561 |      | *           |      |
| Private Health Insurance                  | 0.0000 |      | 0.4383    |      | 0.0600 |      | *           |      |
| Medicare                                  | 0.2884 |      | 0.3136    |      | 0.2276 |      | *           |      |
| Medicaid                                  | 0.6339 |      | 0.9328    |      | 0.3887 |      | *           |      |
| Rehabilitation Program                    | 0.8999 |      | 0.8910    |      | 0.5887 |      | *           |      |
| Employer                                  | 0.1639 |      | 0.7893    |      | 0.1977 |      | *           |      |
| VA or Other Military Program              | 0.9494 |      | 0.8869    |      | 0.7533 |      | *           |      |
| Other Public Source                       | 0.2718 |      | 0.7025    |      | 0.4127 |      | *           |      |
| Other Private Source                      | 0.3098 |      | 0.2582    |      | 0.7691 |      | *           |      |
| Free Treatment                            | 0.9277 |      | 0.0033    |      | 1.0000 |      | *           |      |

\*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Respondents could indicate multiple sources of payment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).