

Table 6.4A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Numbers in Thousands, 2003 and 2004

Drug	ILLCIT DRUG USE			
	Yes		No	
	2003	2004	2003	2004
ILLCIT DRUG¹	5,341	5,905	14,248	15,512
Marijuana and Hashish	3,698	4,203	15,890	17,214
Cocaine	1,077	1,277	18,511	20,139
Crack	325	*	19,263	21,005
Heroin	*	*	19,533	21,228
Hallucinogens	589	700	18,999	20,717
LSD	85	*	19,503	21,271
PCP	*	*	19,560	21,358
Ecstasy	302	344	19,287	21,073
Inhalants	219	*	19,369	21,114
Nonmedical Use of Psychotherapeutics ²	2,761	3,259	16,827	18,157
Pain Relievers	2,107	2,298	17,482	19,118
OxyContin [®]	--	*	--	20,978
Tranquilizers	1,172 ^a	1,670	18,416	19,746
Stimulants	593	756	18,995	20,661
Methamphetamine	331	*	19,257	21,013
Sedatives	*	*	19,298	21,231
ILLCIT DRUG OTHER THAN MARIJUANA¹	3,500	3,972	16,089	17,444

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.4B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Percentages, 2003 and 2004

Drug	ILLCIT DRUG USE			
	Yes		No	
	2003	2004	2003	2004
ILLCIT DRUG¹	18.1	20.6	7.8	8.3
Marijuana and Hashish	17.2 ^a	20.2	8.3	8.9
Cocaine	19.7	23.3	8.9	9.6
Crack	24.8	*	9.1	9.8
Heroin	*	*	9.2	9.9
Hallucinogens	18.7	22.3	9.1	9.8
LSD	20.8	*	9.2	9.9
PCP	*	*	9.2	9.9
Ecstasy	16.9	21.7	9.1	9.9
Inhalants	22.9	*	9.1	9.9
Nonmedical Use of Psychotherapeutics ²	21.7	25.4	8.4	9.0
Pain Relievers	21.6	24.1	8.6	9.3
OxyContin [®]	--	*	--	9.8
Tranquilizers	26.2	32.7	8.8	9.4
Stimulants	27.2	33.9	9.0	9.7
Methamphetamine	29.0	*	9.1	9.8
Sedatives	*	*	9.1	9.9
ILLCIT DRUG OTHER THAN MARIJUANA¹	20.6	23.3	8.2	8.8

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.5A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Numbers in Thousands, 2003 and 2004

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG²	29,538	28,728	5,341	5,905	24,197	22,823
Marijuana and Hashish	21,486	20,818	3,698	4,203	17,788	16,615
Cocaine	5,470	5,486	1,077	1,277	4,392	4,209
Crack	1,310	1,445	325	411	985	1,034
Heroin	278	380	55	188	222	191
Hallucinogens	3,158	3,133	589	700	2,568	2,433
LSD	409	390	85	146	324	244
PCP	130	147	29	58	101	89
Ecstasy	1,790	1,586	302	344	1,488	1,243
Inhalants	956	1,090	219	303	737	787
Nonmedical Use of Psychotherapeutics ³	12,697	12,829	2,761	3,259	9,935	9,569
Pain Relievers	9,753	9,524	2,107	2,298	7,647	7,225
OxyContin [®]	--	1,116	--	439	--	676
Tranquilizers	4,465	5,101	1,172 ^a	1,670	3,293	3,431
Stimulants	2,183	2,230	593	756	1,589	1,474
Methamphetamine	1,141	1,119	331	404	810	715
Sedatives	700	665	290	185	410	480
ILLCIT DRUG OTHER THAN MARIJUANA²	16,954	17,013	3,500	3,972	13,455	13,041

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.5B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Percentages, 2003 and 2004

Drug	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG²	13.9	13.3	27.3	27.6	12.5	11.8
Marijuana and Hashish	10.1	9.7	18.9	19.6	9.2	8.6
Cocaine	2.6	2.5	5.5	6.0	2.3	2.2
Crack	0.6	0.7	1.7	1.9	0.5	0.5
Heroin	0.1	0.2	0.3	0.9	0.1	0.1
Hallucinogens	1.5	1.5	3.0	3.3	1.3	1.3
LSD	0.2	0.2	0.4	0.7	0.2	0.1
PCP	0.1	0.1	0.1	0.3	0.1	0.0
Ecstasy	0.8	0.7	1.5	1.6	0.8	0.6
Inhalants	0.4	0.5	1.1	1.4	0.4	0.4
Nonmedical Use of Psychotherapeutics ³	6.0	6.0	14.1	15.2	5.1	4.9
Pain Relievers	4.6	4.4	10.8	10.7	4.0	3.7
OxyContin [®]	--	0.5	--	2.1	--	0.3
Tranquilizers	2.1	2.4	6.0	7.8	1.7	1.8
Stimulants	1.0	1.0	3.0	3.5	0.8	0.8
Methamphetamine	0.5	0.5	1.7	1.9	0.4	0.4
Sedatives	0.3	0.3	1.5	0.9	0.2	0.2
ILLCIT DRUG OTHER THAN MARIJUANA²	8.0	7.9	17.9	18.5	7.0	6.7

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.6A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Numbers in Thousands, 2003 and 2004

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2003	2004	2003	2004
PAST YEAR				
Tobacco Product ¹	10,533	10,367	9,055 ^a	11,049
Cigarettes	9,715	9,698	9,874 ^a	11,719
Smokeless Tobacco	733	869	18,856	20,548
Cigars	2,747	2,560	16,841 ^a	18,856
Pipe Tobacco ²	--	--	--	--
Alcohol	14,039	14,358	5,550 ^a	7,059
Binge Alcohol Use ³	--	--	--	--
Heavy Alcohol Use ³	--	--	--	--
PAST MONTH				
Tobacco Product ¹	9,270	9,321	10,318 ^a	12,096
Cigarettes	8,666	8,736	10,923 ^a	12,681
Smokeless Tobacco	480	564	19,109	20,853
Cigars	1,355	1,373	18,233	20,044
Pipe Tobacco	259	249	19,329	21,168
Alcohol	10,412	11,164	9,176	10,252
Binge Alcohol Use ³	5,663	6,485	13,925	14,932
Heavy Alcohol Use ³	1,876	2,071	17,712	19,346

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.6B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco Product and Alcohol Use: Percentages, 2003 and 2004

Substance	TOBACCO PRODUCT OR ALCOHOL USE			
	Yes		No	
	2003	2004	2003	2004
PAST YEAR				
Tobacco Product ¹	13.5	13.4	6.7 ^a	8.0
Cigarettes	14.9	14.9	6.7 ^a	7.8
Smokeless Tobacco	7.9	10.1	9.3	9.9
Cigars	12.0	11.2	8.9	9.8
Pipe Tobacco ²	--	--	--	--
Alcohol	9.6	9.8	8.3 ^a	10.3
Binge Alcohol Use ³	--	--	--	--
Heavy Alcohol Use ³	--	--	--	--
PAST MONTH				
Tobacco Product ¹	13.8	14.1	7.1	8.1
Cigarettes	15.1	15.6	7.0	8.0
Smokeless Tobacco	6.6	9.0	9.3	10.0
Cigars	11.6	10.7	9.1	9.9
Pipe Tobacco	17.8	14.6	9.2	9.9
Alcohol	9.1	9.7	9.3	10.2
Binge Alcohol Use ³	11.1	12.6	8.6	9.1
Heavy Alcohol Use ³	12.1	13.3	9.0	9.7

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.7A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Numbers in Thousands, 2003 and 2004

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ²	77,781	77,088	10,533	10,367	67,247	66,721
Cigarettes	65,116	64,884	9,715	9,698	55,401	55,186
Smokeless Tobacco	9,312	8,599	733	869	8,579	7,730
Cigars	22,898	22,956	2,747	2,560	20,151	20,396
Pipe Tobacco ³	--	--	--	--	--	--
Alcohol	145,978	146,589	14,039	14,358	131,939	132,232
Binge Alcohol Use ⁴	--	--	--	--	--	--
Heavy Alcohol Use ⁴	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ²	67,147	66,090	9,270	9,321	57,877	56,769
Cigarettes	57,391	56,144	8,666	8,736	48,725	47,408
Smokeless Tobacco	7,214 ^a	6,243	480	564	6,734 ^a	5,680
Cigars	11,722	12,868	1,355	1,373	10,367	11,495
Pipe Tobacco	1,458	1,700	259	249	1,199	1,452
Alcohol	114,539	114,542	10,412	11,164	104,127	103,378
Binge Alcohol Use ⁴	51,112	51,492	5,663	6,485	45,449	45,007
Heavy Alcohol Use ⁴	15,492	15,569	1,876	2,071	13,616	13,498

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Information about past year use of pipe tobacco was not collected.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.7B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Percentages, 2003 and 2004

Substance	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ²	36.6	35.8	53.8 ^a	48.4	34.8	34.4
Cigarettes	30.6	30.1	49.6	45.3	28.7	28.5
Smokeless Tobacco	4.4	4.0	3.7	4.1	4.4	4.0
Cigars	10.8	10.7	14.0	12.0	10.4	10.5
Pipe Tobacco ³	--	--	--	--	--	--
Alcohol	68.6	68.1	71.7	67.0	68.3	68.2
Binge Alcohol Use ⁴	--	--	--	--	--	--
Heavy Alcohol Use ⁴	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ²	31.6	30.7	47.3	43.5	30.0	29.3
Cigarettes	27.0	26.1	44.2	40.8	25.2	24.5
Smokeless Tobacco	3.4 ^a	2.9	2.4	2.6	3.5 ^a	2.9
Cigars	5.5	6.0	6.9	6.4	5.4	5.9
Pipe Tobacco	0.7	0.8	1.3	1.2	0.6	0.7
Alcohol	53.9	53.2	53.2	52.1	53.9	53.3
Binge Alcohol Use ⁴	24.0	23.9	28.9	30.3	23.5	23.2
Heavy Alcohol Use ⁴	7.3	7.2	9.6	9.7	7.1	7.0

*Low precision; no estimate reported.

-- Not available.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Information about past year use of pipe tobacco was not collected.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.8A Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Use and Demographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	19,588	21,417	5,341	5,905	14,248	15,512
AGE						
18-25	4,420	4,413	2,181	2,107	2,239	2,305
26-49	10,418	10,408	2,840	3,206	7,578	7,202
50 or Older	4,750 ^a	6,596	320	*	4,431 ^a	6,004
GENDER						
Male	6,887	7,976	2,134	2,616	4,753	5,359
Female	12,702	13,441	3,207	3,289	9,495	10,152
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	17,257	18,520	4,801	5,373	12,456	13,147
White	14,308	15,543	3,965	4,508	10,343	11,035
Black or African American	1,992	1,972	604	536	1,387	1,436
American Indian or Alaska Native	96	*	*	*	62	*
Native Hawaiian or Other Pacific Islander	52	*	*	*	29	*
Asian	536	608	*	*	425	464
Two or More Races	273	166	63	*	211 ^a	86
Hispanic or Latino	2,331	2,897	540	532	1,792	2,365
EDUCATION						
< High School	4,276	4,635	1,184	1,299	3,092	3,337
High School Graduate	6,360	6,955	1,706	1,986	4,654	4,969
Some College	5,388	5,543	1,679	1,605	3,709	3,938
College Graduate	3,564	4,283	771	1,015	2,793	3,268
CURRENT EMPLOYMENT						
Full-Time	9,583	8,986	2,795	2,507	6,788	6,479
Part-Time	2,861	3,003	878	1,225	1,983	1,778
Unemployed	1,230	1,526	495	793	736	732
Other ²	5,914 ^a	7,902	1,174	1,380	4,740 ^a	6,522

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.8B Serious Psychological Distress in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug Use and Demographic Characteristics: Percentages, 2003 and 2004

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	9.2	9.9	18.1	20.6	7.8	8.3
AGE						
18-25	13.9	13.7	19.9	19.8	10.8	10.7
26-49	10.4	10.4	17.9	21.1	9.0	8.5
50 or Older	5.9 ^a	7.9	11.7	*	5.7	7.5
GENDER						
Male	6.7	7.7	12.5 ^a	15.7	5.6	6.2
Female	11.5	12.0	25.7	27.3	9.7	10.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	9.2	9.8	18.5	21.0	7.7	8.1
White	9.5	10.3	18.6 ^a	21.8	8.0	8.4
Black or African American	8.4	8.1	17.3	15.0	6.9	6.9
American Indian or Alaska Native	8.9	*	*	*	6.9	*
Native Hawaiian or Other Pacific Islander	12.4	*	*	*	8.6	*
Asian	6.1	6.7	*	*	5.2	5.5
Two or More Races	12.6	8.7	14.8	*	12.0	5.7
Hispanic or Latino	9.0	10.8	15.3	16.9	8.0	10.0
EDUCATION						
< High School	11.3	12.8	21.2	26.2	9.6	10.7
High School Graduate	9.5	10.2	17.9	21.2	8.1	8.4
Some College	10.3	10.0	19.3	19.1	8.5	8.4
College Graduate	6.5	7.8	13.5	16.9	5.7	6.6
CURRENT EMPLOYMENT						
Full-Time	8.2	7.7	16.2	15.1	6.8	6.4
Part-Time	10.0	11.2	16.5 ^a	25.0	8.5	8.2
Unemployed	15.2	19.0	21.3 ^a	33.8	12.8	12.8
Other ²	10.0 ^a	12.5	25.5	28.4	8.7 ^a	11.1

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.9A Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress and Demographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic Characteristic	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	29,538	28,728	5,341	5,905	24,197	22,823
AGE						
18-25	10,976	10,626	2,181	2,107	8,795	8,519
26-49	15,835	15,172	2,840	3,206	12,995	11,966
50 or Older	2,727	2,929	320	591	2,407	2,338
GENDER						
Male	17,044	16,675	2,134	2,616	14,910	14,059
Female	12,494	12,053	3,207	3,289	9,287	8,764
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	26,020	25,576	4,801	5,373	21,219	20,203
White	21,275	20,652	3,965	4,508	17,310	16,144
Black or African American	3,496	3,574	604	536	2,891	3,038
American Indian or Alaska Native	179	309	*	*	145	*
Native Hawaiian or Other Pacific Islander	74	*	*	*	52	*
Asian	575	556	*	*	463	412
Two or More Races	421	413	*	*	359	333
Hispanic or Latino	3,518	3,152	540	532	2,978	2,620
EDUCATION						
< High School	5,579	4,957	1,184	1,299	4,395 ^a	3,659
High School Graduate	9,549	9,365	1,706	1,986	7,843	7,379
Some College	8,678	8,416	1,679	1,605	6,999	6,811
College Graduate	5,732	5,989	771	1,015	4,960	4,974
CURRENT EMPLOYMENT						
Full-Time	17,305	16,611	2,795	2,507	14,510	14,104
Part-Time	5,310	4,905	878	1,225	4,433 ^b	3,680
Unemployed	2,320	2,345	495	*	1,826	1,552
Other ²	4,602	4,866	1,174	1,380	3,428	3,487

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).

Table 6.9B Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress and Demographic Characteristics: Percentages, 2003 and 2004

Demographic Characteristic	Total		SERIOUS PSYCHOLOGICAL DISTRESS ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	13.9	13.3	27.3	27.6	12.5	11.8
AGE						
18-25	34.6	33.0	49.3	47.8	32.2	30.7
26-49	15.8	15.2	27.3	30.8	14.5	13.3
50 or Older	3.4	3.5	6.7	9.0	3.2	3.1
GENDER						
Male	16.7	16.1	31.0	32.8	15.6	14.7
Female	11.3	10.8	25.2	24.5	9.5	8.9
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13.9	13.6	27.8	29.0	12.5	11.9
White	14.1	13.6	27.7	29.0	12.7	11.9
Black or African American	14.7	14.7	30.3	27.2	13.3	13.6
American Indian or Alaska Native	16.6	26.4	*	*	14.7	*
Native Hawaiian or Other Pacific Islander	17.8	*	*	*	14.2	*
Asian	6.5	6.2	*	*	5.6	4.9
Two or More Races	19.4	21.6	*	*	18.9	19.1
Hispanic or Latino	13.6	11.8	23.2	18.4	12.7	11.0
EDUCATION						
< High School	14.7	13.7	27.7	28.0	13.1	11.6
High School Graduate	14.2	13.7	26.8	28.6	12.9	12.0
Some College	16.5	15.1	31.2	29.0	14.9	13.6
College Graduate	10.4	10.9	21.6	23.7	9.6	9.8
CURRENT EMPLOYMENT						
Full-Time	14.8	14.2	29.2	27.9	13.5	13.0
Part-Time	18.5	18.4	30.7 ^a	40.8	17.2	15.5
Unemployed	28.7	29.1	40.2	*	26.6	23.8
Other ²	7.8	7.7	19.8	17.5	6.5	6.3

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on a subsample of respondents aged 18 or older. See Section B.4.4 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Serious Psychological Distress (SPD), referred to as Serious Mental Illness (SMI) in prior NSDUH reports, is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 of Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

² Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004 (subsample).