

Table 1.118C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	517	564	299	300	207	170
GEOGRAPHIC DIVISION						
Northeast	214	252	99	133	71	62
New England	130	116	65	45	46	24
Middle Atlantic	173	227	76	125	54	57
Midwest	195	241	110	123	67	77
East North Central	152	199	96	104	60	64
West North Central	122	136	53	66	29	44
South	301	322	166	167	125	107
South Atlantic	211	248	122	134	96	77
East South Central	133	113	57	58	43	46
West South Central	167	171	96	81	67	59
West	303	306	199	171	133	86
Mountain	120	119	76	86	55	43
Pacific	277	282	184	148	121	74
COUNTY TYPE						
Large Metro	406	441	244	227	163	120
Small Metro	261	255	154	144	104	96
250K – 1 Mil. Pop.	223	200	128	119	77	76
< 250K Pop.	151	157	89	84	70	58
Nonmetro	178	200	105	113	69	64
Urbanized	121	136	69	75	48	39
Less Urbanized	108	132	73	82	49	51
Completely Rural	56	62	22	34	8	17

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.118D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.22	0.23	0.13	0.12	0.09	0.07
GEOGRAPHIC DIVISION						
Northeast	0.47	0.55	0.22	0.29	0.16	0.14
New England	1.09	0.97	0.54	0.38	0.39	0.20
Middle Atlantic	0.52	0.68	0.23	0.37	0.16	0.17
Midwest	0.36	0.45	0.20	0.23	0.12	0.14
East North Central	0.40	0.53	0.25	0.27	0.16	0.17
West North Central	0.76	0.84	0.33	0.41	0.18	0.27
South	0.36	0.37	0.20	0.19	0.15	0.12
South Atlantic	0.48	0.55	0.28	0.30	0.22	0.17
East South Central	0.93	0.78	0.40	0.40	0.30	0.32
West South Central	0.64	0.64	0.37	0.30	0.26	0.22
West	0.56	0.56	0.37	0.31	0.25	0.16
Mountain	0.77	0.74	0.49	0.54	0.35	0.27
Pacific	0.73	0.73	0.48	0.38	0.32	0.19
COUNTY TYPE						
Large Metro	0.32	0.34	0.19	0.18	0.13	0.09
Small Metro	0.37	0.36	0.22	0.20	0.15	0.13
250K – 1 Mil. Pop.	0.48	0.42	0.27	0.25	0.16	0.16
< 250K Pop.	0.62	0.64	0.36	0.34	0.29	0.24
Nonmetro	0.44	0.49	0.26	0.28	0.17	0.16
Urbanized	0.68	0.82	0.39	0.45	0.27	0.23
Less Urbanized	0.59	0.65	0.40	0.41	0.27	0.25
Completely Rural	1.37	1.75	0.55	0.96	0.20	0.48

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.118P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.8008		0.1920		0.2092	
GEOGRAPHIC DIVISION						
Northeast	0.1556		0.2255		0.5451	
New England	0.6915		0.0900		0.3139	
Middle Atlantic	0.0494		0.0163		0.1634	
Midwest	0.6815		0.8136		0.6571	
East North Central	0.5233		0.5434		0.6953	
West North Central	0.8630		0.1783		0.0998	
South	0.7155		0.2800		0.7464	
South Atlantic	0.4299		0.2791		0.6298	
East South Central	0.7981		0.1208		0.3999	
West South Central	0.4920		0.1218		0.5693	
West	0.2556		0.0903		0.0624	
Mountain	0.1888		0.4375		0.7295	
Pacific	0.4737		0.1298		0.0587	
COUNTY TYPE						
Large Metro	0.4658		0.0601		0.1235	
Small Metro	0.2094		0.9479		0.9457	
250K – 1 Mil. Pop.	0.2450		0.7226		0.5875	
< 250K Pop.	0.5934		0.7050		0.5412	
Nonmetro	0.5118		0.5710		0.7587	
Urbanized	0.2623		0.8339		0.2365	
Less Urbanized	0.8395		0.8307		0.9852	
Completely Rural	0.9316		0.0697		0.0672	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.119C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	70	67	59	56	39	36
GEOGRAPHIC DIVISION						
Northeast	30	29	23	26	14	17
New England	17	16	14	11	7	9
Middle Atlantic	25	24	18	24	13	14
Midwest	28	28	25	24	15	14
East North Central	24	22	21	19	13	13
West North Central	15	18	13	15	7	6
South	44	42	36	34	24	23
South Atlantic	32	30	27	25	15	17
East South Central	20	19	15	13	11	8
West South Central	22	22	19	19	15	13
West	37	33	33	28	22	18
Mountain	21	15	20	11	10	7
Pacific	30	30	27	25	19	17
COUNTY TYPE						
Large Metro	50	51	41	43	29	28
Small Metro	37	38	30	30	17	19
250K – 1 Mil. Pop.	32	29	25	23	14	14
< 250K Pop.	21	25	17	19	10	12
Nonmetro	29	26	25	21	17	16
Urbanized	16	16	15	12	11	9
Less Urbanized	22	20	17	17	12	12
Completely Rural	8	7	7	4	5	4

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.119D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.28	0.27	0.24	0.22	0.15	0.14
GEOGRAPHIC DIVISION						
Northeast	0.66	0.64	0.50	0.58	0.32	0.37
New England	1.41	1.39	1.17	0.93	0.57	0.78
Middle Atlantic	0.74	0.72	0.54	0.71	0.38	0.42
Midwest	0.49	0.49	0.44	0.42	0.27	0.25
East North Central	0.60	0.54	0.53	0.46	0.34	0.33
West North Central	0.87	1.08	0.75	0.88	0.44	0.36
South	0.49	0.47	0.40	0.37	0.27	0.25
South Atlantic	0.71	0.66	0.60	0.54	0.34	0.36
East South Central	1.41	1.34	1.03	0.88	0.73	0.57
West South Central	0.75	0.75	0.64	0.63	0.50	0.44
West	0.63	0.56	0.57	0.46	0.38	0.30
Mountain	1.24	0.84	1.15	0.64	0.60	0.38
Pacific	0.71	0.70	0.64	0.59	0.47	0.39
COUNTY TYPE						
Large Metro	0.37	0.38	0.31	0.32	0.22	0.21
Small Metro	0.50	0.49	0.41	0.40	0.23	0.24
250K – 1 Mil. Pop.	0.63	0.57	0.51	0.45	0.28	0.28
< 250K Pop.	0.88	0.97	0.68	0.75	0.41	0.48
Nonmetro	0.68	0.63	0.59	0.49	0.40	0.37
Urbanized	0.89	0.92	0.83	0.71	0.58	0.52
Less Urbanized	1.11	0.95	0.86	0.77	0.61	0.57
Completely Rural	1.79	1.93	1.57	1.17	1.20	1.09

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.119P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.5730		0.4035		0.2709	
GEOGRAPHIC DIVISION						
Northeast	0.5409		0.6724		0.9043	
New England	0.6566		0.5053		0.2266	
Middle Atlantic	0.6594		0.9346		0.5564	
Midwest	0.2255		0.9180		0.3523	
East North Central	0.1711		0.9214		0.4365	
West North Central	0.8688		0.9582		0.5843	
South	0.6770		0.3552		0.4477	
South Atlantic	0.8946		0.6080		0.9071	
East South Central	0.9141		0.7174		0.4667	
West South Central	0.3344		0.4786		0.5450	
West	0.9393		0.4545		0.5760	
Mountain	0.2585		0.1464		0.0544	
Pacific	0.4430		0.9905		0.7720	
COUNTY TYPE						
Large Metro	0.6402		0.9675		0.9367	
Small Metro	0.9754		0.4818		0.1591	
250K – 1 Mil. Pop.	0.8392		0.7792		0.5702	
< 250K Pop.	0.8277		0.3844		0.1112	
Nonmetro	0.5702		0.2511		0.3969	
Urbanized	0.9562		0.2718		0.3489	
Less Urbanized	0.6626		0.5782		0.7459	
Completely Rural	0.5778		0.6595		0.8747	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.120C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	130	129	97	94	61	60
GEOGRAPHIC DIVISION						
Northeast	47	55	39	40	22	21
New England	24	23	22	19	12	10
Middle Atlantic	41	50	33	35	19	18
Midwest	53	57	40	40	26	24
East North Central	43	48	31	33	22	21
West North Central	32	32	25	22	14	12
South	80	75	59	53	37	35
South Atlantic	57	49	43	33	31	25
East South Central	36	32	22	21	14	14
West South Central	44	46	33	35	16	20
West	73	71	53	53	34	36
Mountain	37	29	28	21	12	12
Pacific	62	64	44	48	32	34
COUNTY TYPE						
Large Metro	98	95	74	66	46	40
Small Metro	72	72	52	58	32	36
250K – 1 Mil. Pop.	61	60	40	48	23	28
< 250K Pop.	40	41	34	35	22	24
Nonmetro	42	49	34	33	23	25
Urbanized	28	35	23	23	16	17
Less Urbanized	27	31	22	23	15	16
Completely Rural	9	14	6	11	3	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 1.120D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Percentages, 2003 and 2004**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.41	0.40	0.30	0.29	0.19	0.19
GEOGRAPHIC DIVISION						
Northeast	0.83	0.96	0.70	0.70	0.39	0.37
New England	1.62	1.54	1.50	1.25	0.80	0.70
Middle Atlantic	0.98	1.18	0.80	0.83	0.45	0.44
Midwest	0.73	0.77	0.54	0.54	0.36	0.32
East North Central	0.85	0.93	0.62	0.65	0.43	0.40
West North Central	1.40	1.38	1.08	0.96	0.64	0.53
South	0.71	0.65	0.52	0.46	0.33	0.31
South Atlantic	1.01	0.86	0.77	0.59	0.55	0.44
East South Central	1.85	1.66	1.16	1.09	0.73	0.74
West South Central	1.16	1.20	0.88	0.90	0.43	0.51
West	0.98	0.94	0.71	0.70	0.46	0.47
Mountain	1.63	1.27	1.25	0.92	0.55	0.50
Pacific	1.20	1.23	0.85	0.92	0.62	0.64
COUNTY TYPE						
Large Metro	0.59	0.57	0.45	0.39	0.28	0.24
Small Metro	0.71	0.68	0.51	0.55	0.32	0.34
250K – 1 Mil. Pop.	0.92	0.87	0.60	0.70	0.34	0.40
< 250K Pop.	1.13	1.10	0.95	0.93	0.63	0.63
Nonmetro	0.86	1.00	0.69	0.66	0.47	0.50
Urbanized	1.19	1.51	0.95	0.98	0.67	0.73
Less Urbanized	1.24	1.34	1.01	0.98	0.71	0.68
Completely Rural	2.52	4.23	1.60	3.43	0.94	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.120P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.3035		0.8556		0.9697	
GEOGRAPHIC DIVISION						
Northeast	0.0728		0.2156		0.1609	
New England	0.5618		0.9122		0.8820	
Middle Atlantic	0.0185		0.1522		0.1361	
Midwest	0.5501		0.8631		0.5436	
East North Central	0.4220		0.4673		0.3766	
West North Central	0.9205		0.1916		0.0240	
South	0.7617		0.5244		0.4969	
South Atlantic	0.9325		0.5225		0.6269	
East South Central	0.4405		0.7302		0.7428	
West South Central	0.2373		0.5681		0.7196	
West	0.6528		0.6748		0.8074	
Mountain	0.6743		0.6428		0.6204	
Pacific	0.4706		0.8239		0.6614	
COUNTY TYPE						
Large Metro	0.9926		0.2869		0.3093	
Small Metro	0.1224		0.3200		0.1602	
250K – 1 Mil. Pop.	0.5102		0.1381		0.1282	
< 250K Pop.	0.0653		0.7699		0.6755	
Nonmetro	0.7333		0.8837		0.7066	
Urbanized	0.9901		0.4718		0.3685	
Less Urbanized	0.8134		0.5828		0.5413	
Completely Rural	0.0505		0.0141		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.121C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	477	529	275	265	193	152
GEOGRAPHIC DIVISION						
Northeast	208	250	87	112	61	51
New England	133	113	63	35	43	15
Middle Atlantic	163	225	61	106	44	49
Midwest	189	218	97	107	55	70
East North Central	146	183	85	91	48	56
West North Central	121	120	48	57	27	42
South	274	297	149	154	115	104
South Atlantic	194	231	111	123	87	68
East South Central	115	106	49	60	42	51
West South Central	155	155	87	69	62	59
West	268	285	188	151	130	69
Mountain	115	115	71	76	54	39
Pacific	241	261	174	130	119	57
COUNTY TYPE						
Large Metro	376	412	229	196	154	109
Small Metro	238	241	138	125	95	83
250K – 1 Mil. Pop.	193	196	115	107	70	67
< 250K Pop.	139	142	77	67	65	49
Nonmetro	171	184	90	105	62	57
Urbanized	112	121	58	69	42	34
Less Urbanized	103	122	63	74	46	44
Completely Rural	58	62	21	32	4	12

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.121D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.26	0.29	0.15	0.14	0.11	0.08
GEOGRAPHIC DIVISION						
Northeast	0.59	0.71	0.25	0.32	0.17	0.15
New England	1.43	1.22	0.68	0.37	0.46	0.16
Middle Atlantic	0.63	0.87	0.24	0.41	0.17	0.19
Midwest	0.46	0.53	0.24	0.26	0.14	0.17
East North Central	0.51	0.63	0.30	0.32	0.17	0.19
West North Central	1.00	0.98	0.40	0.46	0.23	0.34
South	0.42	0.45	0.23	0.23	0.18	0.16
South Atlantic	0.56	0.66	0.32	0.35	0.25	0.20
East South Central	1.06	0.96	0.45	0.55	0.38	0.46
West South Central	0.80	0.79	0.45	0.35	0.32	0.30
West	0.66	0.69	0.46	0.37	0.32	0.17
Mountain	0.99	0.96	0.61	0.63	0.46	0.33
Pacific	0.84	0.89	0.60	0.45	0.41	0.19
COUNTY TYPE						
Large Metro	0.39	0.42	0.24	0.20	0.16	0.11
Small Metro	0.44	0.45	0.26	0.23	0.18	0.16
250K – 1 Mil. Pop.	0.55	0.56	0.33	0.30	0.20	0.19
< 250K Pop.	0.75	0.78	0.41	0.37	0.35	0.27
Nonmetro	0.55	0.59	0.29	0.33	0.20	0.18
Urbanized	0.82	0.95	0.43	0.54	0.31	0.27
Less Urbanized	0.73	0.77	0.45	0.47	0.33	0.28
Completely Rural	1.76	2.15	0.65	1.13	0.13	0.41

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.121P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.9214		0.2184		0.2550	
GEOGRAPHIC DIVISION						
Northeast	0.3515		0.4359		0.8570	
New England	0.7059		0.0352		0.1312	
Middle Atlantic	0.1722		0.0284		0.2251	
Midwest	0.9446		0.8587		0.9525	
East North Central	0.8212		0.6470		0.7265	
West North Central	0.8425		0.3070		0.4741	
South	0.5837		0.4691		0.9611	
South Atlantic	0.4041		0.3799		0.7587	
East South Central	0.9486		0.1141		0.3100	
West South Central	0.8542		0.1959		0.7193	
West	0.1587		0.1016		0.0362	
Mountain	0.3032		0.7497		0.9080	
Pacific	0.2758		0.0915		0.0135	
COUNTY TYPE						
Large Metro	0.4178		0.0973		0.1930	
Small Metro	0.4429		0.6555		0.7551	
250K – 1 Mil. Pop.	0.3070		0.7452		0.9118	
< 250K Pop.	0.9138		0.7419		0.5237	
Nonmetro	0.6086		0.2883		0.9291	
Urbanized	0.2584		0.8617		0.4356	
Less Urbanized	0.9120		0.3890		0.6576	
Completely Rural	0.6921		0.2326		0.1694	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.122C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	510	559	298	289	205	164
GEOGRAPHIC DIVISION						
Northeast	215	251	98	125	71	56
New England	133	119	67	43	48	20
Middle Atlantic	173	224	73	117	53	52
Midwest	191	233	108	117	64	76
East North Central	149	192	93	100	57	63
West North Central	119	132	53	61	28	44
South	300	320	164	164	122	105
South Atlantic	208	247	120	133	93	73
East South Central	136	109	58	58	43	48
West South Central	166	170	95	77	65	59
West	294	305	200	166	134	83
Mountain	114	121	74	85	54	43
Pacific	271	280	186	143	123	71
COUNTY TYPE						
Large Metro	401	436	247	217	163	116
Small Metro	259	251	153	139	101	93
250K – 1 Mil. Pop.	219	199	129	117	76	75
< 250K Pop.	147	154	85	80	68	55
Nonmetro	176	198	99	111	68	62
Urbanized	119	136	67	74	46	38
Less Urbanized	106	129	67	80	48	47
Completely Rural	58	62	22	34	5	17

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.122D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.24	0.26	0.14	0.13	0.10	0.08
GEOGRAPHIC DIVISION						
Northeast	0.53	0.61	0.24	0.30	0.17	0.14
New England	1.24	1.11	0.63	0.40	0.45	0.19
Middle Atlantic	0.57	0.74	0.24	0.39	0.17	0.17
Midwest	0.40	0.48	0.22	0.24	0.13	0.16
East North Central	0.44	0.57	0.28	0.29	0.17	0.19
West North Central	0.83	0.90	0.37	0.42	0.19	0.30
South	0.39	0.41	0.22	0.21	0.16	0.14
South Atlantic	0.52	0.61	0.30	0.33	0.23	0.18
East South Central	1.06	0.84	0.46	0.45	0.34	0.37
West South Central	0.72	0.72	0.41	0.33	0.28	0.25
West	0.61	0.63	0.42	0.34	0.28	0.17
Mountain	0.82	0.85	0.53	0.60	0.39	0.30
Pacific	0.79	0.81	0.55	0.42	0.36	0.21
COUNTY TYPE						
Large Metro	0.36	0.38	0.22	0.19	0.14	0.10
Small Metro	0.41	0.39	0.24	0.22	0.16	0.14
250K – 1 Mil. Pop.	0.52	0.47	0.31	0.28	0.18	0.18
< 250K Pop.	0.67	0.70	0.39	0.36	0.31	0.25
Nonmetro	0.49	0.54	0.28	0.31	0.19	0.17
Urbanized	0.74	0.91	0.42	0.49	0.29	0.26
Less Urbanized	0.65	0.71	0.41	0.44	0.29	0.26
Completely Rural	1.58	1.94	0.60	1.06	0.15	0.52

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 1.122P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2003	2004	2003	2004	2003	2004
TOTAL	0.8587		0.2465		0.2870	
GEOGRAPHIC DIVISION						
Northeast	0.1823		0.2487		0.5517	
New England	0.6493		0.0743		0.2133	
Middle Atlantic	0.0561		0.0119		0.1112	
Midwest	0.8094		0.8311		0.8047	
East North Central	0.6494		0.5116		0.5590	
West North Central	0.8344		0.1733		0.1236	
South	0.6685		0.3743		0.8641	
South Atlantic	0.4357		0.3217		0.6342	
East South Central	0.7835		0.1104		0.3422	
West South Central	0.5728		0.1422		0.6545	
West	0.2441		0.1141		0.0718	
Mountain	0.2520		0.6525		0.9949	
Pacific	0.4199		0.1219		0.0464	
COUNTY TYPE						
Large Metro	0.4208		0.0593		0.1235	
Small Metro	0.1982		0.8645		0.7685	
250K – 1 Mil. Pop.	0.2269		0.6967		0.5257	
< 250K Pop.	0.6027		0.7951		0.7242	
Nonmetro	0.5694		0.3636		0.9379	
Urbanized	0.2603		0.9759		0.3200	
Less Urbanized	0.7739		0.6786		0.9040	
Completely Rural	0.9892		0.0523		0.0356	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.