

Table 6.78C Perception of Help Received from Treatment or Counseling among Persons Aged 12 or Older with a Past Year Major Depressive Episode (MDE) Who Saw or Talked to a Medical Doctor or Other Professional in the Past Year about Depression, by Age Group: Standard Errors of Numbers in Thousands, 2004

| Perception of Help Received ¹ | Total | AGE GROUP | |
|--|-------|-----------|-------------|
| | | 12-17 | 18 or Older |
| TOTAL | 518 | 39 | 513 |
| No Help at All | 172 | 17 | 171 |
| A Little Help | 182 | 18 | 181 |
| Some Help | 281 | 19 | 279 |
| A Lot of Help | 254 | 17 | 252 |
| Extremely Helpful | 224 | 14 | 222 |

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total row represent persons aged 12 or older with a past year MDE who saw or talked to a medical doctor or other professional in the past year about depression, including those with unknown perception of help information. Estimates of levels of perceived help *exclude* persons with unknown perception of help information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.78D Perception of Help Received from Treatment or Counseling among Persons Aged 12 or Older with a Past Year Major Depressive Episode (MDE) Who Saw or Talked to a Medical Doctor or Other Professional in the Past Year about Depression, by Age Group: Standard Errors of Percentages, 2004

| Perception of Help Received ¹ | Total | AGE GROUP | |
|--|-------|-----------|-------------|
| | | 12-17 | 18 or Older |
| TOTAL | 0.00 | 0.00 | 0.00 |
| No Help at All | 1.42 | 1.91 | 1.53 |
| A Little Help | 1.45 | 1.97 | 1.55 |
| Some Help | 2.14 | 2.00 | 2.30 |
| A Lot of Help | 1.98 | 1.94 | 2.13 |
| Extremely Helpful | 1.75 | 1.62 | 1.87 |

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total row represent persons aged 12 or older with a past year MDE who saw or talked to a medical doctor or other professional in the past year about depression, including those with unknown perception of help information. Estimates of levels of perceived help *exclude* persons with unknown perception of help information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.79C Perception of Help Received from Prescription Medication among Persons Aged 12 or Older with a Past Year Major Depressive Episode (MDE) Who Received Prescription Medication in the Past Year for Depression, by Age Group: Standard Errors of Numbers in Thousands, 2004

| Perception of Help Received ¹ | Total | AGE GROUP | |
|--|-------|-----------|-------------|
| | | 12-17 | 18 or Older |
| TOTAL | 437 | 27 | 435 |
| No Help at All | 96 | 10 | 96 |
| A Little Help | 133 | 12 | 133 |
| Some Help | 240 | 13 | 239 |
| A Lot of Help | 262 | 14 | 261 |
| Extremely Helpful | 174 | 11 | 172 |

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total row represent persons aged 12 or older with a past year MDE who received prescription medication in the past year for depression, including those with unknown perception of help information. Estimates of levels of perceived help *exclude* persons with unknown perception of help information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).

Table 6.79D Perception of Help Received from Prescription Medication among Persons Aged 12 or Older with a Past Year Major Depressive Episode (MDE) Who Received Prescription Medication in the Past Year for Depression, by Age Group: Standard Errors of Percentages, 2004

| Perception of Help Received ¹ | Total | AGE GROUP | |
|--|-------|-----------|-------------|
| | | 12-17 | 18 or Older |
| TOTAL | 0.00 | 0.00 | 0.00 |
| No Help at All | 1.07 | 2.27 | 1.12 |
| A Little Help | 1.45 | 2.57 | 1.51 |
| Some Help | 2.30 | 2.76 | 2.41 |
| A Lot of Help | 2.32 | 2.88 | 2.43 |
| Extremely Helpful | 1.83 | 2.41 | 1.90 |

*Low precision; no estimate reported.

NOTE: Estimates in this table are based on all respondents aged 12 to 17 and a subsample of respondents aged 18 or older. See Section B.4.5 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

NOTE: Major Depressive Episode (MDE) is defined as a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of the symptoms for depression as described in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Estimates in the Total row represent persons aged 12 or older with a past year MDE who received prescription medication in the past year for depression, including those with unknown perception of help information. Estimates of levels of perceived help *exclude* persons with unknown perception of help information.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 (subsample).