

**Table 6.33C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2003 and 2004**

Age Category	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	561	554	150	149	398	413	529	500
18	30	34	13	15	22	27	26	28
19	28	29	9	9	22	22	25	24
20	28	30	9	14	21	22	24	25
21	30	30	12	13	23	26	28	26
22	31	29	9	11	24	23	27	28
23	32	25	8	7	27	18	28	22
24	29	27	8	10	26	21	26	24
25	31	27	6	10	24	21	26	23
26-29	106	113	38	34	92	92	95	98
30-34	147	132	45	31	122	112	127	115
35-39	166	171	34	33	126	126	152	158
40-44	160	169	45	44	130	135	149	158
45-49	185	159	56	53	144	130	168	146
50-54	236	222	59	64	170	186	217	202
55-59	195	184	50	54	128	131	190	177
60-64	155	164	25	45	99	112	152	160
65 or Older	236	252	80	77	146	147	220	232

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.33D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Category: Standard Errors of Percentages, 2003 and 2004**

Age Category	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.26	0.26	0.07	0.07	0.19	0.19	0.25	0.23
18	0.65	0.74	0.29	0.34	0.48	0.58	0.57	0.61
19	0.73	0.75	0.23	0.22	0.58	0.56	0.64	0.62
20	0.71	0.76	0.23	0.36	0.52	0.55	0.60	0.63
21	0.76	0.75	0.32	0.32	0.59	0.63	0.70	0.64
22	0.77	0.73	0.21	0.27	0.61	0.57	0.67	0.70
23	0.79	0.63	0.21	0.19	0.67	0.47	0.69	0.56
24	0.79	0.67	0.22	0.25	0.71	0.52	0.70	0.59
25	0.87	0.75	0.18	0.27	0.67	0.57	0.73	0.63
26-29	0.72	0.74	0.26	0.22	0.62	0.60	0.64	0.64
30-34	0.73	0.67	0.22	0.16	0.61	0.57	0.63	0.58
35-39	0.80	0.82	0.16	0.16	0.60	0.60	0.73	0.75
40-44	0.71	0.75	0.20	0.19	0.57	0.60	0.66	0.70
45-49	0.87	0.74	0.26	0.25	0.67	0.60	0.79	0.68
50-54	1.18	1.15	0.30	0.33	0.86	0.96	1.09	1.04
55-59	1.31	1.13	0.34	0.33	0.86	0.80	1.28	1.08
60-64	1.32	1.31	0.21	0.36	0.85	0.89	1.29	1.27
65 or Older	0.69	0.73	0.23	0.22	0.43	0.43	0.64	0.67

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.33P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003**

Age Category	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.3206		0.6165		0.9495		0.2168	
18	0.6371		0.7710		0.2607		0.8113	
19	0.5492		0.8258		0.8891		0.4316	
20	0.4110		0.4323		0.6433		0.4650	
21	0.4676		0.9517		0.5021		0.2292	
22	0.9816		0.2657		0.7278		0.6061	
23	0.0142		0.4918		0.0005		0.0620	
24	0.3003		0.1353		0.0457		0.5917	
25	0.7460		0.0804		0.4926		0.6604	
26-29	0.2996		0.8773		0.9080		0.2737	
30-34	0.4177		0.1979		0.2875		0.4833	
35-39	0.5695		0.8903		0.6335		0.6552	
40-44	0.8622		0.6724		0.9159		0.5972	
45-49	0.1675		0.4963		0.9318		0.0538	
50-54	0.5592		0.5887		0.8678		0.5976	
55-59	0.4468		0.6783		0.9634		0.2874	
60-64	0.9594		0.6293		0.7938		0.8338	
65 or Older	0.7691		0.8119		0.4871		0.5798	

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.34C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	561	554	150	149	398	413	529	500
<b>AGE</b>								
18-25	89	85	28	33	66	66	78	74
26-49	354	338	95	88	281	268	322	305
50 or Older	425	423	113	117	280	296	409	396
<b>GENDER</b>								
Male	319	327	112	104	226	241	285	300
Female	448	443	102	115	320	338	433	406
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	527	552	131	138	381	406	495	502
White	503	503	101	105	360	372	473	460
Black or African American	151	167	82	71	110	108	133	159
American Indian or Alaska Native	33	34	19	6	19	22	26	29
Native Hawaiian or Other Pacific Islander	*	*	0	1	3	*	*	*
Asian	89	82	11	57	62	56	81	76
Two or More Races	58	49	9	8	48	41	49	40
Hispanic or Latino	163	156	69	71	116	106	159	138
<b>EDUCATION</b>								
< High School	200	223	80	82	141	143	194	197
High School Graduate	308	283	100	108	218	211	294	267
Some College	278	290	67	65	190	209	271	275
College Graduate	286	287	48	33	229	247	251	265
<b>CURRENT EMPLOYMENT</b>								
Full-Time	400	348	56	80	287	257	364	318
Part-Time	200	201	38	45	160	167	183	177
Unemployed	87	100	32	42	66	74	73	96
Other <sup>4</sup>	337	352	131	113	230	257	320	328

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.34D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.26	0.26	0.07	0.07	0.19	0.19	0.25	0.23
<b>AGE</b>								
18-25	0.28	0.26	0.09	0.10	0.21	0.21	0.25	0.23
26-49	0.35	0.34	0.10	0.09	0.28	0.27	0.32	0.31
50 or Older	0.53	0.51	0.14	0.14	0.35	0.36	0.51	0.48
<b>GENDER</b>								
Male	0.31	0.32	0.11	0.10	0.22	0.23	0.28	0.29
Female	0.41	0.40	0.09	0.10	0.29	0.30	0.39	0.36
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.28	0.29	0.07	0.07	0.20	0.22	0.27	0.27
White	0.33	0.33	0.07	0.07	0.24	0.25	0.31	0.30
Black or African American	0.64	0.69	0.34	0.29	0.47	0.45	0.56	0.66
American Indian or Alaska Native	3.02	2.96	1.72	0.54	1.76	1.92	2.41	2.59
Native Hawaiian or Other Pacific Islander	*	*	0.02	0.11	0.82	*	*	*
Asian	1.02	0.92	0.13	0.64	0.71	0.63	0.93	0.85
Two or More Races	2.74	2.29	0.44	0.40	2.24	1.94	2.29	1.87
Hispanic or Latino	0.64	0.58	0.27	0.27	0.45	0.40	0.61	0.52
<b>EDUCATION</b>								
< High School	0.53	0.61	0.21	0.23	0.37	0.39	0.51	0.54
High School Graduate	0.46	0.42	0.15	0.16	0.33	0.31	0.44	0.39
Some College	0.53	0.53	0.13	0.12	0.36	0.38	0.52	0.50
College Graduate	0.52	0.51	0.09	0.06	0.42	0.44	0.46	0.47
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.34	0.29	0.05	0.07	0.25	0.22	0.31	0.27
Part-Time	0.70	0.73	0.13	0.16	0.56	0.61	0.64	0.64
Unemployed	1.08	1.36	0.39	0.57	0.82	1.00	0.91	1.30
Other <sup>4</sup>	0.57	0.57	0.22	0.18	0.39	0.42	0.54	0.53

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.34P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2004 Versus 2003**

Demographic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>			
			Inpatient		Outpatient <sup>2</sup>	
	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.3206		0.6165		0.9495	0.2168
<b>AGE</b>						
18-25	0.4278		0.1781		0.1172	0.5630
26-49	0.7406		0.4915		0.8415	0.6741
50 or Older	0.4141		0.4357		0.5803	0.2669
<b>GENDER</b>						
Male	0.6110		0.6356		0.4280	0.5037
Female	0.0991		0.8583		0.5011	0.0482
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.4053		0.8636		1.0000	0.3461
White	0.4422		0.9184		0.9466	0.4456
Black or African American	0.9641		0.9501		0.7033	0.6874
American Indian or Alaska Native	0.9887		0.2855		0.8515	0.8178
Native Hawaiian or Other Pacific Islander	*		0.2741		*	*
Asian	0.9362		0.2088		0.6736	0.8746
Two or More Races	0.6290		0.8346		0.9301	0.5964
Hispanic or Latino	0.6425		0.4900		0.9203	0.3605
<b>EDUCATION</b>						
< High School	0.3082		0.4124		0.7722	0.9833
High School Graduate	0.0965		0.4748		0.8340	0.1387
Some College	0.9563		0.7457		0.6489	0.5709
College Graduate	0.2421		0.4297		0.5387	0.6432
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.0081		0.1667		0.0115	0.0200
Part-Time	0.7528		0.6770		0.3337	0.7854
Unemployed	0.0064		0.9957		0.0930	0.0008
Other <sup>4</sup>	0.7475		0.6853		0.1184	0.7633

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Retired persons, disabled persons, homemakers, students, or other persons not in the labor force are included in the Other Employment category.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.35C Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	657	601	151	149	444	430	599	539
<b>GEOGRAPHIC REGION</b>								
Northeast	263	273	83	47	200	193	244	249
Midwest	251	225	46	69	189	168	237	197
South	410	367	93	113	260	267	381	325
West	362	317	72	52	232	221	313	289
<b>COUNTY TYPE</b>								
Large Metro	477	459	119	112	332	318	435	412
Small Metro	404	349	82	83	254	259	359	321
250K - 1 Mil. Pop.	321	294	70	64	216	199	287	266
< 250K Pop.	248	198	43	54	143	153	221	189
Nonmetro	245	251	51	56	163	175	230	210
Urbanized	184	169	46	43	113	129	176	136
Less Urbanized	178	173	19	36	117	104	166	165
Completely Rural	85	82	9	8	51	46	85	59
<b>FAMILY INCOME</b>								
Less Than \$20,000	299	294	113	101	208	209	272	264
\$20,000 - \$49,999	359	330	88	77	239	238	330	294
\$50,000 - \$74,999	262	277	48	70	186	201	246	265
\$75,000 or More	321	313	24	37	230	220	279	269
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	257	264	104	90	175	199	248	231
No	607	528	113	119	406	374	541	483

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.35D Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2003 and 2004**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
<b>TOTAL</b>	0.26	0.26	0.07	0.07	0.19	0.19	0.25	0.23
<b>GEOGRAPHIC REGION</b>								
Northeast	0.62	0.61	0.20	0.12	0.47	0.44	0.60	0.56
Midwest	0.45	0.45	0.09	0.14	0.36	0.34	0.43	0.40
South	0.45	0.44	0.12	0.14	0.30	0.34	0.43	0.39
West	0.63	0.59	0.15	0.11	0.43	0.42	0.56	0.54
<b>COUNTY TYPE</b>								
Large Metro	0.37	0.36	0.10	0.10	0.27	0.26	0.35	0.32
Small Metro	0.51	0.46	0.13	0.13	0.34	0.37	0.46	0.44
250K - 1 Mil. Pop.	0.60	0.61	0.17	0.15	0.43	0.44	0.56	0.56
< 250K Pop.	0.92	0.73	0.19	0.24	0.56	0.65	0.83	0.72
Nonmetro	0.56	0.64	0.14	0.15	0.40	0.47	0.54	0.53
Urbanized	0.89	0.97	0.28	0.28	0.64	0.82	0.89	0.76
Less Urbanized	0.86	0.87	0.12	0.20	0.64	0.55	0.83	0.85
Completely Rural	2.02	2.37	0.26	0.25	1.28	1.41	2.03	1.72
<b>FAMILY INCOME</b>								
Less Than \$20,000	0.60	0.63	0.27	0.24	0.45	0.47	0.57	0.57
\$20,000 - \$49,999	0.41	0.39	0.11	0.10	0.28	0.30	0.38	0.35
\$50,000 - \$74,999	0.61	0.67	0.12	0.18	0.45	0.50	0.57	0.64
\$75,000 or More	0.55	0.50	0.05	0.07	0.41	0.37	0.50	0.45
<b>GOVERNMENT ASSISTANCE<sup>4</sup></b>								
Yes	0.77	0.78	0.37	0.29	0.58	0.63	0.77	0.69
No	0.29	0.26	0.06	0.06	0.20	0.19	0.26	0.24

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.



**Table 6.35P Specific Types of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Geographic or Socioeconomic Characteristics: P Values from Tests of Differences of Percentages, 2004 Versus 2003**

Geographic/Socioeconomic Characteristic	Received Mental Health Treatment/Counseling <sup>1,2</sup>		TYPE OF MENTAL HEALTH TREATMENT/COUNSELING <sup>3</sup>					
			Inpatient		Outpatient <sup>2</sup>		Prescription Medication	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.3206		0.6165		0.9495		0.2168	
GEOGRAPHIC REGION								
Northeast	0.6103		0.2298		0.8097		0.9640	
Midwest	0.2374		0.0756		0.7339		0.0735	
South	0.1648		0.3259		0.4874		0.0536	
West	0.3522		0.3024		0.2141		0.2685	
COUNTY TYPE								
Large Metro	0.2766		0.8413		0.4710		0.2784	
Small Metro	0.6216		0.7797		0.3216		0.7134	
250K - 1 Mil. Pop.	0.5878		0.6224		0.4147		0.8661	
< 250K Pop.	0.9321		0.2286		0.5623		0.7039	
Nonmetro	0.2427		0.6294		0.8075		0.0906	
Urbanized	0.5422		0.8280		0.6235		0.3041	
Less Urbanized	0.3102		0.3174		0.5308		0.2741	
Completely Rural	0.9112		0.4249		0.4712		0.5618	
FAMILY INCOME								
Less Than \$20,000	0.7450		0.7362		0.5220		0.7467	
\$20,000 - \$49,999	0.2990		0.9171		0.3987		0.1735	
\$50,000 - \$74,999	0.7414		0.4916		0.8096		0.7413	
\$75,000 or More	0.3440		0.7149		0.0429		0.7060	
GOVERNMENT ASSISTANCE <sup>4</sup>								
Yes	0.6706		0.4160		0.8894		0.4406	
No	0.2283		0.3526		0.8173		0.1931	

\*Low precision; no estimate reported.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded.

<sup>2</sup> Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Respondents could report receiving multiple types of mental health treatment/counseling; thus, these response categories are not mutually exclusive.

<sup>4</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.