

Table 7.121A No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drug or Alcohol ¹		Illicit Drug ¹		Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	19,521	19,734	22,184	22,540	20,570	20,779
RACE²						
American Indian or Alaska Native	244	231	271	259	266	261
Asian or Pacific Islander	946	954	1,000	1,024	975	983
Asian Only	871	885	923	945	898	910
Native Hawaiian or Other Pacific Islander Only	*	*	*	78	*	*
Black or African American	3,270	3,350	3,531	3,598	3,499	3,580
White	14,720	14,832	17,010	17,241	15,466	15,563
Two or More Races	341	368	373	419	364	392
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	3,187	3,283	3,618	3,800	3,405	3,471
Not Hispanic or Latino	16,333	16,451	18,566	18,741	17,165	17,307
Black or African American	3,143	3,215	3,383	3,449	3,363	3,432
White	11,822	11,857	13,727	13,781	12,370	12,419
GENDER						
Male	10,033	10,145	11,314	11,517	10,586	10,661
Female	9,487	9,589	10,870	11,024	9,984	10,117
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	3,265	3,453	3,589	3,809	3,518	3,715
100-199%	4,425	4,442	5,005	4,982	4,675	4,678
200% or More	11,831	11,840	13,590	13,749	12,377	12,386
REGION						
Northeast	3,417	3,449	3,963	3,994	3,609	3,663
Midwest	4,444	4,390	5,086	5,098	4,639	4,622
South	7,046	7,262	7,995	8,178	7,440	7,603
West	4,614	4,633	5,141	5,270	4,881	4,891

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.121B No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drug or Alcohol ¹		Illicit Drug ¹		Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	78.1	78.3	88.8	89.4	82.3	82.4
RACE²						
American Indian or Alaska Native	75.5	72.8	83.6	81.7	82.2	82.3
Asian or Pacific Islander	87.9	87.5	92.9	94.0	90.6	90.2
Asian Only	88.3	88.0	93.5	94.0	91.0	90.5
Native Hawaiian or Other Pacific Islander Only	*	*	*	93.6	*	*
Black or African American	83.7	84.3	90.4	90.5	89.6	90.1
White	76.5	76.6	88.4	89.1	80.3	80.4
Two or More Races	77.4	77.5	84.9	88.3	82.7	82.5
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	78.4	77.6	89.0	89.8	83.8	82.0
Not Hispanic or Latino	78.0	78.4	88.7	89.3	82.0	82.5
Black or African American	84.0	84.5	90.4	90.7	89.9	90.2
White	76.0	76.5	88.2	88.9	79.5	80.1
GENDER						
Male	78.6	78.8	88.6	89.4	82.9	82.8
Female	77.6	77.7	88.9	89.4	81.7	82.0
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	79.8	79.6	87.7	87.8	86.0	85.6
100-199%	78.7	80.2	89.0	90.0	83.2	84.5
200% or More	77.4	77.2	88.9	89.6	81.0	80.8
REGION						
Northeast	76.1	76.0	88.2	88.1	80.4	80.8
Midwest	78.0	77.3	89.3	89.7	81.5	81.4
South	78.9 ^a	80.7	89.6 ^a	90.9	83.4	84.5
West	78.4	77.3	87.4	87.9	83.0	81.6

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.122A Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	2,811	2,674	1,971	1,909	2,658	2,794
RACE³						
American Indian or Alaska Native	53	58	40	36	42	42
Asian or Pacific Islander	77	66	39	46	43	56
Asian Only	64	60	30	43	33	50
Native Hawaiian or Other Pacific Islander Only	*	5	*	3	*	*
Black or African American	374	376	233	257	188	198
White	2,241	2,119	1,612	1,525	2,337	2,443
Two or More Races	67	56	47	45	49	54
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	447	433	278	284	383 ^b	519
Not Hispanic or Latino	2,365	2,240	1,694	1,624	2,276	2,275
Black or African American	358	354	221	243	172	186
White	1,838	1,727	1,366	1,266	1,999	1,957
GENDER						
Male	1,457	1,360	1,093	1,037	1,421	1,496
Female	1,355	1,314	878	872	1,237	1,298
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	503	530	315	346	356	412
100-199%	618	557	434	407	572	547
200% or More	1,690	1,587	1,223	1,157	1,731	1,835
REGION						
Northeast	528	542	393	389	519	519
Midwest	609	584	457	417	678	700
South	932	823	611	544	852	845
West	743	726	511	559	609	730

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.122B Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	11.2	10.6	7.9	7.6	10.6	11.1
RACE³						
American Indian or Alaska Native	16.4	18.3	12.5	11.3	12.9	13.4
Asian or Pacific Islander	7.1	6.0	3.6	4.2	4.0	5.2
Asian Only	6.5	6.0	3.0	4.3	3.4	5.0
Native Hawaiian or Other Pacific Islander Only	*	6.4	*	3.9	*	*
Black or African American	9.6	9.5	6.0	6.5	4.8	5.0
White	11.6	10.9	8.4	7.9	12.1	12.6
Two or More Races	15.1	11.7	10.7	9.5	11.1	11.3
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	11.0	10.2	6.8	6.7	9.4 ^b	12.3
Not Hispanic or Latino	11.3	10.7	8.1	7.7	10.9	10.8
Black or African American	9.6	9.3	5.9	6.4	4.6	4.9
White	11.8	11.1	8.8	8.2	12.8	12.6
GENDER						
Male	11.4	10.6	8.6	8.1	11.1	11.6
Female	11.1	10.6	7.2	7.1	10.1	10.5
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	12.3	12.2	7.7	8.0	8.7	9.5
100-199%	11.0	10.0	7.7	7.3	10.2	9.9
200% or More	11.1	10.4	8.0	7.5	11.3	12.0
REGION						
Northeast	11.8	11.9	8.7	8.6	11.6	11.4
Midwest	10.7	10.3	8.0	7.3	11.9	12.3
South	10.4 ^a	9.1	6.8	6.0	9.5	9.4
West	12.6	12.1	8.7	9.3	10.4	12.2

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.123A Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	16,659	16,397	12,667	12,667	51,112	51,931
RACE³						
American Indian or Alaska Native	163	163	131	117	558	540
Asian or Pacific Islander	358	369	191	245	1,214	1,402
Asian Only	312	255	161	156	1,066	1,202
Native Hawaiian or Other Pacific Islander Only	46	*	30	*	148	*
Black or African American	2,077	2,156	1,661	1,797	5,173	5,061
White	13,788	13,403	10,467	10,297	43,588	44,282
Two or More Races	273	306	217	211	579	646
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	1,932	1,806	1,197	1,258	6,857	6,929
Not Hispanic or Latino	14,726	14,591	11,470	11,408	44,255	45,002
Black or African American	2,029	2,075	1,627	1,725	5,042	4,936
White	11,980	11,756	9,355	9,158	37,160	37,844
GENDER						
Male	10,006	10,132	8,197	8,236	34,144	34,699
Female	6,653	6,265	4,470	4,431	16,968	17,232
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	2,714 ^a	3,161	1,963 ^b	2,507	5,701 ^b	6,627
100-199%	3,790 ^a	3,241	2,894	2,569	9,535	8,843
200% or More	10,155	9,995	7,810	7,591	35,875	36,461
EDUCATION						
< High School	3,394	3,131	2,538	2,399	9,206	8,407
High School Graduate	5,562	5,344	4,272	4,057	16,824	17,144
Some College/College Graduate	7,703	7,922	5,857	6,211	25,082	26,381

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.123B Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2003	2004	2003	2004	2003	2004
TOTAL	7.8	7.6	6.0	5.9	24.0	24.1
RACE³						
American Indian or Alaska Native	8.8	8.5	7.0	6.1	30.0	28.2
Asian or Pacific Islander	3.8	3.8	2.0	2.5	13.0	14.5
Asian Only	3.5	2.8	1.8	1.7	12.0	13.3
Native Hawaiian or Other Pacific Islander Only	10.3	*	6.8	*	33.1	*
Black or African American	8.5	8.7	6.8	7.2	21.2	20.4
White	7.9	7.6	6.0	5.8	25.0	25.1
Two or More Races	11.4	12.4	9.1	8.6	24.2	26.2
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	7.5	6.7	4.6	4.7	26.6	25.9
Not Hispanic or Latino	7.9	7.7	6.1	6.1	23.7	23.9
Black or African American	8.6	8.6	6.9	7.1	21.3	20.4
White	8.0	7.8	6.2	6.0	24.7	25.0
GENDER						
Male	9.8	9.8	8.0	7.9	33.4	33.5
Female	6.0	5.6	4.0	4.0	15.4	15.4
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	11.6	12.4	8.4 ^a	9.9	24.4	26.0
100-199%	9.4	8.3	7.2	6.6	23.7	22.6
200% or More	6.8	6.6	5.2	5.0	24.1	24.2
EDUCATION						
< High School	9.0	8.6	6.7	6.6	24.3	23.0
High School Graduate	8.3	7.8	6.4	6.0	25.1	25.2
Some College/College Graduate	7.2	7.2	5.4	5.6	23.3	23.8

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Illicit Drug Use includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.124A Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2003	2004
TOTAL	1,119	1,168
RACE¹		
American Indian or Alaska Native	12	18
Asian or Pacific Islander	40	49
Asian Only	33	47
Native Hawaiian or Other Pacific Islander Only	*	2
Black or African American	90	107
White	954	966
Two or More Races	23	29
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	172	202
Not Hispanic or Latino	947	967
Black or African American	87	98
White	795	783
GENDER		
Male	554	587
Female	565	581
FAMILY INCOME (% of Federal Poverty Threshold)		
Less than 100%	162	186
100-199%	233	216
200% or More	724	766
REGION		
Northeast	191	227
Midwest	258	260
South	370	373
West	301	308

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.124B Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2003	2004
TOTAL	4.5	4.6
RACE¹		
American Indian or Alaska Native	3.8	5.7
Asian or Pacific Islander	3.7	4.5
Asian Only	3.3	4.7
Native Hawaiian or Other Pacific Islander Only	*	2.2
Black or African American	2.3	2.7
White	5.0	5.0
Two or More Races	5.2	6.0
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	4.2	4.8
Not Hispanic or Latino	4.5	4.6
Black or African American	2.3	2.6
White	5.1	5.1
GENDER		
Male	4.3	4.6
Female	4.6	4.7
FAMILY INCOME (% of Federal Poverty Threshold)		
Less than 100%	3.9	4.3
100-199%	4.1	3.9
200% or More	4.7	5.0
REGION		
Northeast	4.3	5.0
Midwest	4.5	4.6
South	4.1	4.1
West	5.1	5.1

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.125A Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2003	2004	2003	2004	2003	2004
TOTAL	284	305	1,538	1,410	984	930
RACE¹						
American Indian or Alaska Native	8	7	18	20	16	18
Asian or Pacific Islander	2	*	32	43	18	20
Asian Only	1	*	30	41	12	13
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	51	54	245	226	118	129
White	212	231	1,208	1,082	808	735
Two or More Races	11	8	35	39	23	27
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	54	63	272	258	153	131
Not Hispanic or Latino	230	242	1,266	1,152	830	799
Black or African American	50	54	233	209	114	125
White	161	170	959	852	664	614
GENDER						
Male	182	179	817	744	534	471
Female	102	126	721	666	449	458
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	84	102	320	294	220	220
100-199%	62	68	368	317	308 ^a	232
200% or More	138	135	849	798	455	477
REGION						
Northeast	36	45	284	263	120	137
Midwest	57	74	346	344	264	246
South	99	106	553	485	394	352
West	91	80	355	318	205	195

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.125B Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2003	2004	2003	2004	2003	2004
TOTAL	3.4	3.6	18.2	16.9	11.6	11.1
RACE¹						
American Indian or Alaska Native	6.8	6.3	15.9	18.2	14.5	16.1
Asian or Pacific Islander	0.6	*	9.4	12.3	5.4	5.7
Asian Only	0.3	*	9.7	12.7	4.0	4.1
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	3.7	3.9	17.9	16.2	8.6	9.3
White	3.3	3.7	18.7	17.1	12.5	11.6
Two or More Races	6.3	4.6	20.6	21.2	13.8	14.9
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	3.8	4.2	18.9	17.5	10.7	8.9
Not Hispanic or Latino	3.3	3.5	18.0	16.7	11.8	11.6
Black or African American	3.8	4.0	17.8	15.7	8.7	9.4
White	3.1	3.4	18.6	17.1	12.9	12.3
GENDER						
Male	4.2	4.2	18.7	17.5	12.2	11.1
Female	2.5	3.0	17.6	16.2	11.0	11.1
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	5.6	6.5	21.3	18.8	14.7	14.1
100-199%	3.2	3.6	19.2	16.6	16.1 ^a	12.2
200% or More	2.7	2.8	16.9	16.3	9.0	9.8
REGION						
Northeast	2.4	3.0	18.6	17.6	7.9	9.2
Midwest	2.9	3.9	17.9	18.2	13.6	13.0
South	3.4	3.5	19.0 ^a	16.1	13.6	11.7
West	4.4	4.0	17.1	16.1	9.9	9.9

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.126A Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2003	2004	2003	2004	2003	2004
TOTAL	8,614	8,699	12,628	12,320	9,528	9,514
RACE²						
American Indian or Alaska Native	96	109	133	154	118	133
Asian or Pacific Islander	404	401	455	468	479	507
Asian Only	372	374	420	423	442	474
Native Hawaiian or Other Pacific Islander Only	*	*	35	*	37	*
Black or African American	1,338	1,395	2,152	2,134	1,917	1,922
White	6,642	6,642	9,673	9,327	6,834	6,777
Two or More Races	134	153	216	237	179	175
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	1,434	1,553	1,855	1,874	1,552	1,647
Not Hispanic or Latino	7,180	7,146	10,773	10,446	7,976	7,868
Black or African American	1,283	1,329	2,068	2,043	1,838	1,850
White	5,326	5,233	7,993	7,640	5,451	5,284
GENDER						
Male	4,183	4,130	6,581	6,337	4,484	4,336
Female	4,432	4,569	6,047	5,984	5,044	5,178
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	1,479	1,546	2,046	2,071	1,675	1,771
100-199%	2,003	2,027	2,818	2,771	2,299	2,278
200% or More	5,132	5,127	7,764	7,479	5,554	5,466
REGION						
Northeast	1,368	1,391	2,173	2,131	1,624	1,655
Midwest	1,971	1,892	2,855	2,733	2,074	1,984
South	3,329	3,436	4,801	4,657	3,621	3,616
West	1,946	1,981	2,798	2,799	2,210	2,260

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.126B Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2003	2004	2003	2004	2003	2004
TOTAL	34.9	35.0	51.4 ^b	49.6	38.5	38.1
RACE²						
American Indian or Alaska Native	30.3	34.9	42.2	49.6	37.1	42.8
Asian or Pacific Islander	38.3	37.9	43.2	43.9	45.1	47.2
Asian Only	38.5	38.4	43.4	43.0	45.3	47.8
Native Hawaiian or Other Pacific Islander Only	*	*	41.2	*	42.1	*
Black or African American	34.9	35.6	56.3	54.5	49.7	48.8
White	34.9	34.7	51.1 ^b	48.9	35.8	35.3
Two or More Races	31.3	32.6	49.5	50.9	41.1	37.4
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	36.0	37.3	46.6	45.2	38.8	39.3
Not Hispanic or Latino	34.7	34.5	52.3 ^b	50.5	38.4	37.8
Black or African American	34.9	35.4	56.5	54.5	49.8	49.1
White	34.5	34.1	52.1 ^b	50.0	35.2	34.3
GENDER						
Male	33.2	32.6	52.5 ^b	50.1	35.5	34.0
Female	36.7	37.4	50.2	49.2	41.6	42.3
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	36.9	36.2	51.3	48.7	41.8	41.3
100-199%	36.0	37.2	50.9	50.8	41.2	41.5
200% or More	34.0	33.8	51.6 ^b	49.5	36.6	35.9
REGION						
Northeast	31.0	31.0	49.5	47.7	36.5	36.9
Midwest	34.9	33.7	50.7	48.8	36.7	35.2
South	37.7	38.7	54.7	52.6	41.0	40.5
West	33.6	33.4	48.5	47.4	37.9	38.0

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.127A Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Numbers in Thousands, 2003 and 2004

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drug		Alcohol		Either Illicit Drug or Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	1,103 ^a	1,427	1,298	1,535	1,874 ^a	2,327
RACE¹						
American Indian or Alaska Native	28	8	41	*	53	*
Asian or Pacific Islander	9	*	12	*	17	*
Asian Only	9	*	*	*	11	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	205	336	259	285	315	445
White	829	983	957	1,158	1,450	1,733
Two or More Races	31	34	30	32	39	41
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	89	142	140	234	191	314
Not Hispanic or Latino	1,014	1,285	1,159	1,301	1,683	2,013
Black or African American	202	334	251	285	305	443
White	757	845	832	928	1,286	1,425
GENDER						
Male	732	914	897	1,149	1,263	1,571
Female	371	513	401	386	612	755
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	281 ^a	459	360	532	467 ^a	768
100-199%	287	383	325	337	486	556
200% or More	535	585	613	667	922	1,002
AGE GROUP						
12-17	113	134	95	126	168	185
18 or Older	990 ^a	1,293	1,203	1,409	1,707 ^a	2,142

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.127B Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Percentages, 2003 and 2004

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drug		Alcohol		Either Illicit Drug or Alcohol	
	2003	2004	2003	2004	2003	2004
TOTAL	5.0	6.1	5.9	6.5	8.5	9.9
RACE¹						
American Indian or Alaska Native	8.2 ^a	2.2	11.9	*	15.3	*
Asian or Pacific Islander	1.3	*	1.8	*	2.5	*
Asian Only	1.4	*	*	*	1.9	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	8.5	12.6	10.7	10.7	13.0	16.7
White	4.5	5.1	5.2	6.0	7.9	8.9
Two or More Races	9.1	9.2	8.8	8.7	11.5	10.9
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	3.0	4.4	4.7	7.3	6.4	9.7
Not Hispanic or Latino	5.3	6.3	6.0	6.4	8.8	9.9
Black or African American	8.7	13.0	10.8	11.1	13.1	17.3
White	4.8	5.1	5.3	5.6	8.2	8.6
GENDER						
Male	5.1	5.9	6.2	7.5	8.8	10.2
Female	4.8	6.4	5.2	4.8	7.9	9.4
FAMILY INCOME (% of Federal Poverty Threshold)						
Less than 100%	8.2	10.8	10.5	12.5	13.6	18.0
100-199%	5.7	8.7	6.5	7.6	9.7	12.6
200% or More	3.9	4.0	4.5	4.5	6.7	6.8
AGE GROUP						
12-17	5.0	5.9	4.2	5.5	7.4	8.1
18 or Older	5.0	6.1	6.0	6.7	8.6	10.1

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.