

Table 6.36A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2003 and 2004

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	29,538	29,506	5,959	5,916	23,512	23,504
Marijuana and Hashish	21,486	21,796	3,959	4,010	17,479	17,739
Cocaine	5,470	5,247	1,076	1,140	4,379	4,104
Crack	1,310	1,241	371	398	938	841
Heroin	278	350	90	126	188	223
Hallucinogens	3,158	3,125	549	596	2,598	2,518
LSD	409	434	67	77	341	351
PCP	130	138	34	24	94	113
Ecstasy	1,790	1,605	311	287	1,471	1,311
Inhalants	956	1,087	203	281	747	806
Nonmedical Use of Psychotherapeutics ⁴	12,697	12,418	3,200	3,047	9,475	9,321
Pain Relievers	9,753	9,391	2,279	2,178	7,462	7,190
OxyContin [®]	--	1,019	--	305	--	712
Tranquilizers	4,465	4,536	1,379	1,514	3,073	3,014
Stimulants	2,183	2,423	553	637	1,627	1,782
Methamphetamine	1,141	1,277	255	357	887	918
Sedatives	700	604	320	240	380	338
ILLCIT DRUG OTHER THAN MARIJUANA³	16,954	16,389	3,925	3,771	12,991	12,563

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.36B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2003 and 2004

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	13.9	13.7	21.3	21.5	12.8	12.5
Marijuana and Hashish	10.1	10.1	14.2	14.6	9.5	9.5
Cocaine	2.6	2.4	3.9	4.1	2.4	2.2
Crack	0.6	0.6	1.3	1.4	0.5	0.4
Heroin	0.1	0.2	0.3	0.5	0.1	0.1
Hallucinogens	1.5	1.5	2.0	2.2	1.4	1.3
LSD	0.2	0.2	0.2	0.3	0.2	0.2
PCP	0.1	0.1	0.1	0.1	0.1	0.1
Ecstasy	0.8	0.7	1.1	1.0	0.8	0.7
Inhalants	0.4	0.5	0.7	1.0	0.4	0.4
Nonmedical Use of Psychotherapeutics ⁴	6.0	5.8	11.5	11.1	5.1	5.0
Pain Relievers	4.6	4.4	8.2	7.9	4.1	3.8
OxyContin [®]	--	0.5	--	1.1	--	0.4
Tranquilizers	2.1	2.1	4.9	5.5	1.7	1.6
Stimulants	1.0	1.1	2.0	2.3	0.9	1.0
Methamphetamine	0.5	0.6	0.9	1.3	0.5	0.5
Sedatives	0.3	0.3	1.1	0.9	0.2	0.2
ILLCIT DRUG OTHER THAN MARIJUANA³	8.0	7.6	14.1	13.7	7.1	6.7

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs

Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.37A Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	77,781	77,504	11,622	11,594	65,950	65,727
Cigarettes	65,116	65,264	10,407	10,423	54,517	54,677
Smokeless Tobacco	9,312	8,672	953	992	8,338	7,665
Cigars	22,898	23,405	2,921	2,965	19,934	20,374
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	145,978	148,149	19,994	19,668	125,671	128,252
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	67,147	66,622	10,188	10,009	56,771	56,443
Cigarettes	57,391	56,897	9,347	9,151	47,868	47,592
Smokeless Tobacco	7,214	6,583	685	731	6,507	5,842
Cigars	11,722	12,526	1,367	1,413	10,318	11,073
Pipe Tobacco	1,458	1,657	271	239	1,188	1,412
Alcohol	114,539	116,498	15,168	15,479	99,155	100,829
Binge Alcohol Use ⁵	51,112	51,931	5,896 ^a	6,579	45,106	45,283
Heavy Alcohol Use ⁵	15,492	16,018	1,750	2,068	13,680	13,920

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.⁴ Information about past year use of pipe tobacco was not collected.⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.37B Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	36.6	36.0	41.6	42.2	35.8	35.1
Cigarettes	30.6	30.3	37.2	37.9	29.6	29.2
Smokeless Tobacco	4.4	4.0	3.4	3.6	4.5 ^a	4.1
Cigars	10.8	10.9	10.5	10.8	10.8	10.9
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	68.6	68.8	71.6	71.5	68.2	68.5
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	31.6	30.9	36.5	36.4	30.8	30.1
Cigarettes	27.0	26.4	33.5	33.3	26.0	25.4
Smokeless Tobacco	3.4	3.1	2.5	2.7	3.5 ^a	3.1
Cigars	5.5	5.8	4.9	5.1	5.6	5.9
Pipe Tobacco	0.7	0.8	1.0	0.9	0.6	0.8
Alcohol	53.9	54.1	54.3	56.3	53.8	53.8
Binge Alcohol Use ⁵	24.0	24.1	21.1 ^b	23.9	24.5	24.2
Heavy Alcohol Use ⁵	7.3	7.4	6.3 ^a	7.5	7.4	7.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.38A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Numbers in Thousands, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	19,372	20,278	3,943 ^a	4,549	15,385	15,684
Illicit Drug ⁴	5,548	5,951	1,522	1,737	4,014	4,194
Alcohol	16,334	17,137	3,103 ^a	3,630	13,191	13,475
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	9,832 ^a	10,719	2,487 ^a	2,970	7,319	7,732
Illicit Drug ⁴	3,680	4,115	1,151	1,339	2,523	2,770
Alcohol	7,051 ^a	7,821	1,655 ^a	2,112	5,373	5,699

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.38B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Percentages, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	9.1	9.4	14.1 ^a	16.5	8.4	8.4
Illicit Drug ⁴	2.6	2.8	5.4	6.3	2.2	2.2
Alcohol	7.7	8.0	11.1 ^a	13.2	7.2	7.2
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	4.6	5.0	8.9 ^a	10.8	4.0	4.1
Illicit Drug ⁴	1.7	1.9	4.1	4.9	1.4	1.5
Alcohol	3.3	3.6	5.9 ^a	7.7	2.9	3.0

*Low precision; no estimate reported.

^a Difference between estimate and 2004 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information are excluded. Due to revised editing of 2003 and 2004 outpatient mental health treatment/counseling data, these 2003 estimates may differ slightly from 2003 estimates published in prior NSDUH reports. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.