

Table 6.50C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	85	93	49	53	50	54	50	54
GENDER								
Male	56	62	35	38	32	37	32	34
Female	62	66	33	36	37	39	38	40
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	78	83	44	49	44	45	44	49
White	66	69	37	39	36	41	37	42
Black or African American	34	41	20	23	21	20	19	22
American Indian or Alaska Native	*	5	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	18	17	10	12	11	7	10	9
Two or More Races	10	11	7	8	5	6	*	7
Hispanic or Latino	36	38	21	23	22	28	23	22

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.50D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.34	0.37	0.59	0.63	0.60	0.62	0.60	0.66
GENDER								
Male	0.44	0.49	0.81	0.90	0.77	0.84	0.76	0.81
Female	0.51	0.53	0.81	0.87	0.93	0.92	0.92	0.99
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.37	0.39	0.63	0.71	0.63	0.63	0.64	0.71
White	0.42	0.45	0.71	0.79	0.70	0.77	0.72	0.81
Black or African American	0.90	1.09	1.56	1.76	1.66	1.56	1.58	1.87
American Indian or Alaska Native	*	3.14	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian	1.85	1.69	3.30	3.63	3.41	1.86	2.81	3.05
Two or More Races	2.56	2.48	4.15	4.86	3.64	4.73	*	5.13
Hispanic or Latino	0.90	0.91	1.49	1.55	1.71	1.95	1.74	1.68

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.50P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.0003		0.0534		0.0516		0.0135	
GENDER								
Male	0.0823		0.1421		0.1287		0.8564	
Female	0.0004		0.1793		0.2114		0.0005	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.0014		0.0364		0.3316		0.0121	
White	0.0005		0.0241		0.0550		0.0724	
Black or African American	0.0730		0.2624		0.8905		0.0305	
American Indian or Alaska Native	*		*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.0020		0.1269		0.0004		0.5660	
Two or More Races	0.3035		0.5580		0.3943		*	
Hispanic or Latino	0.0395		0.8121		0.0173		0.6473	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.51C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	107	116	58	60	57	64	58	63
GEOGRAPHIC REGION								
Northeast	45	50	19	27	25	26	24	26
Midwest	43	44	22	24	25	21	23	25
South	65	67	36	36	32	35	35	39
West	59	67	35	32	30	41	33	35
COUNTY TYPE								
Large Metro	87	92	45	48	44	52	48	47
Small Metro	57	68	29	35	32	36	26	35
250K - 1 Mil. Pop.	49	54	26	28	28	29	21	26
< 250K Pop.	30	49	14	26	16	20	15	24
Nonmetro	39	38	21	18	19	21	18	24
Urbanized	27	24	15	12	13	12	12	15
Less Urbanized	26	29	15	14	13	17	13	17
Completely Rural	18	14	6	6	7	7	8	*
FAMILY INCOME								
Less Than \$20,000	46	51	26	26	27	32	26	29
\$20,000 - \$49,999	61	62	34	37	34	36	35	34
\$50,000 - \$74,999	40	44	20	26	25	26	24	25
\$75,000 or More	56	61	29	28	28	32	31	32
GOVERNMENT ASSISTANCE¹								
Yes	52	57	30	34	30	32	24	28
No	96	98	50	50	50	54	52	55

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.51D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.34	0.37	0.59	0.63	0.60	0.62	0.60	0.66
GEOGRAPHIC REGION								
Northeast	0.81	0.89	1.16	1.48	1.45	1.50	1.45	1.48
Midwest	0.67	0.66	1.00	1.14	1.22	1.01	1.06	1.16
South	0.55	0.58	1.02	1.08	0.92	0.95	0.97	1.10
West	0.75	0.89	1.43	1.41	1.39	1.58	1.46	1.57
COUNTY TYPE								
Large Metro	0.47	0.54	0.80	0.91	0.85	0.91	0.90	0.95
Small Metro	0.66	0.65	1.05	1.09	1.15	1.14	0.95	1.09
250K - 1 Mil. Pop.	0.81	0.77	1.31	1.32	1.50	1.41	1.12	1.26
< 250K Pop.	1.02	1.35	1.59	2.19	1.66	2.03	1.57	2.02
Nonmetro	0.72	0.73	1.26	1.20	1.16	1.20	1.13	1.46
Urbanized	1.24	1.16	1.99	2.19	1.84	1.76	1.84	2.16
Less Urbanized	0.98	1.02	2.04	1.71	1.74	1.84	1.62	1.99
Completely Rural	2.25	2.71	3.15	3.92	2.97	4.75	3.28	*
FAMILY INCOME								
Less Than \$20,000	0.89	0.91	1.44	1.36	1.57	1.70	1.62	1.64
\$20,000 - \$49,999	0.59	0.60	1.01	1.12	1.05	1.01	1.03	1.08
\$50,000 - \$74,999	0.76	0.76	1.25	1.50	1.33	1.33	1.35	1.40
\$75,000 or More	0.64	0.67	1.04	1.03	1.10	1.11	1.16	1.11
GOVERNMENT ASSISTANCE¹								
Yes	0.91	0.97	1.50	1.59	1.74	1.66	1.57	1.65
No	0.37	0.39	0.64	0.68	0.64	0.66	0.64	0.70

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.51P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Age Group and Geographic or Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic/Socioeconomic Characteristic	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
TOTAL	0.0003		0.0534		0.0516		0.0135	
GEOGRAPHIC REGION								
Northeast	0.0235		0.0363		0.1743		0.5537	
Midwest	0.1393		0.0722		0.7129		0.1823	
South	0.0212		0.4851		0.0207		0.2437	
West	0.1695		0.9614		0.6356		0.0870	
COUNTY TYPE								
Large Metro	0.0532		0.1281		0.2089		0.6399	
Small Metro	0.0014		0.1136		0.1470		0.0034	
250K - 1 Mil. Pop.	0.1419		0.4868		0.5913		0.1292	
< 250K Pop.	0.0009		0.1210		0.0619		0.0029	
Nonmetro	0.1467		0.9248		0.6165		0.0413	
Urbanized	0.3295		0.6158		0.9265		0.1311	
Less Urbanized	0.4061		0.4144		0.6868		0.0847	
Completely Rural	0.5422		0.6900		0.4057		*	
FAMILY INCOME								
Less Than \$20,000	0.2143		0.2705		0.7298		0.4683	
\$20,000 - \$49,999	0.0021		0.5332		0.0362		0.0065	
\$50,000 - \$74,999	0.1840		0.1090		0.8649		0.3818	
\$75,000 or More	0.1166		0.3769		0.1507		0.6908	
GOVERNMENT ASSISTANCE¹								
Yes	0.1918		0.2272		0.4685		0.0453	
No	0.0007		0.1331		0.0117		0.0743	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.52C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	85	93	42	46	71	78
AGE						
12-13	49	53	16	19	45	48
14-15	50	54	26	28	42	44
16-17	50	54	32	35	37	40
GENDER						
Male	56	62	28	30	50	56
Female	62	66	32	35	52	54
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	78	83	39	41	65	70
White	66	69	34	34	55	59
Black or African American	34	41	15	19	30	33
American Indian or Alaska Native	*	5	3	*	*	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	18	17	*	*	16	16
Two or More Races	10	11	*	*	9	10
Hispanic or Latino	36	38	20	21	30	34

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.52D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: Standard Errors of Percentages, 2003 and 2004

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	0.34	0.37	0.78	0.88	0.36	0.39
AGE						
12-13	0.59	0.63	2.16	2.46	0.59	0.64
14-15	0.60	0.62	1.42	1.55	0.66	0.65
16-17	0.60	0.66	1.12	1.30	0.68	0.72
GENDER						
Male	0.44	0.49	1.02	1.12	0.50	0.55
Female	0.51	0.53	1.19	1.31	0.54	0.55
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.39	0.85	0.93	0.40	0.42
White	0.42	0.45	0.96	0.99	0.46	0.49
Black or African American	0.90	1.09	2.15	2.75	0.99	1.05
American Indian or Alaska Native	*	3.14	4.98	*	*	4.12
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.85	1.69	*	*	1.93	1.85
Two or More Races	2.56	2.48	*	*	2.85	3.06
Hispanic or Latino	0.90	0.91	2.35	2.51	0.96	1.02

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.52P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Demographic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0003		0.0000		0.0773	
AGE						
12-13	0.0534		0.0266		0.2072	
14-15	0.0516		0.0168		0.3729	
16-17	0.0135		0.0021		0.3367	
GENDER						
Male	0.0823		0.3292		0.1242	
Female	0.0004		0.0000		0.2917	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0014		0.0000		0.1401	
White	0.0005		0.0002		0.0410	
Black or African American	0.0730		0.0021		0.5933	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0020		*		0.0140	
Two or More Races	0.3035		*		0.4812	
Hispanic or Latino	0.0395		0.0384		0.2487	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.53C Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2003 and 2004

Geographic/Socioeconomic Characteristic	Total		ILLICIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	85	93	42	46	71	78
GEOGRAPHIC REGION						
Northeast	36	40	17	22	32	33
Midwest	38	37	19	18	34	30
South	49	52	24	25	40	45
West	44	53	24	27	36	45
COUNTY TYPE						
Large Metro	62	72	33	35	53	58
Small Metro	49	49	22	25	39	43
250K - 1 Mil. Pop.	40	39	18	18	33	33
< 250K Pop.	25	34	13	16	19	30
Nonmetro	31	31	17	19	25	24
Urbanized	23	20	13	11	17	17
Less Urbanized	19	22	10	14	17	16
Completely Rural	10	10	*	*	7	7
FAMILY INCOME						
Less Than \$20,000	40	42	19	20	35	35
\$20,000 - \$49,999	52	52	26	29	42	45
\$50,000 - \$74,999	36	37	17	19	31	31
\$75,000 or More	44	48	24	25	36	38
GOVERNMENT ASSISTANCE²						
Yes	40	45	20	19	33	40
No	76	81	38	44	65	68

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.53D Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: Standard Errors of Percentages, 2003 and 2004

Geographic/Socioeconomic Characteristic	Total		ILLICIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	0.34	0.37	0.78	0.88	0.36	0.39
GEOGRAPHIC REGION						
Northeast	0.81	0.89	1.85	2.19	0.92	0.92
Midwest	0.67	0.66	1.55	1.45	0.75	0.68
South	0.55	0.58	1.27	1.41	0.57	0.63
West	0.75	0.89	1.74	2.07	0.80	0.97
COUNTY TYPE						
Large Metro	0.47	0.54	1.20	1.29	0.51	0.54
Small Metro	0.66	0.65	1.26	1.53	0.68	0.72
250K - 1 Mil. Pop.	0.81	0.77	1.63	1.66	0.85	0.82
< 250K Pop.	1.02	1.35	2.13	2.90	1.03	1.50
Nonmetro	0.72	0.73	1.77	2.03	0.76	0.72
Urbanized	1.24	1.16	2.85	2.78	1.24	1.29
Less Urbanized	0.98	1.02	2.32	2.93	1.08	0.97
Completely Rural	2.25	2.71	*	*	2.04	2.32
FAMILY INCOME						
Less Than \$20,000	0.89	0.91	1.92	1.91	0.99	0.97
\$20,000 - \$49,999	0.59	0.60	1.31	1.57	0.61	0.68
\$50,000 - \$74,999	0.76	0.76	1.67	1.95	0.83	0.81
\$75,000 or More	0.64	0.67	1.68	1.78	0.66	0.66
GOVERNMENT ASSISTANCE²						
Yes	0.91	0.97	1.94	1.78	0.98	1.11
No	0.37	0.39	0.87	1.04	0.40	0.42

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.53P Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17, by Past Year Illicit Drug Use and Geographic or Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Geographic/Socioeconomic Characteristic	Total		ILLCIT DRUG USE ¹			
			Yes		No	
	2003	2004	2003	2004	2003	2004
TOTAL	0.0003		0.0000		0.0773	
GEOGRAPHIC REGION						
Northeast	0.0235		0.0039		0.3337	
Midwest	0.1393		0.0307		0.6028	
South	0.0212		0.0190		0.1031	
West	0.1695		0.0261		0.6723	
COUNTY TYPE						
Large Metro	0.0532		0.0108		0.3783	
Small Metro	0.0014		0.0002		0.0622	
250K - 1 Mil. Pop.	0.1419		0.1132		0.3414	
< 250K Pop.	0.0009		0.0000		0.0931	
Nonmetro	0.1467		0.0549		0.6847	
Urbanized	0.3295		0.7669		0.2721	
Less Urbanized	0.4061		0.0300		0.5170	
Completely Rural	0.5422		*		0.8447	
FAMILY INCOME						
Less Than \$20,000	0.2143		0.0357		0.8344	
\$20,000 - \$49,999	0.0021		0.0002		0.1270	
\$50,000 - \$74,999	0.1840		0.0068		0.8522	
\$75,000 or More	0.1166		0.5841		0.0956	
GOVERNMENT ASSISTANCE²						
Yes	0.1918		0.0026		0.9866	
No	0.0007		0.0003		0.0555	

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.54C Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	107	116	58	60	57	64	58	63
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	70	73	36	37	39	44	40	43
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher ²	70	70	44	42	38	39	36	40
Mental Health Clinic or Center	31	40	19	20	17	25	19	21
In-Home Therapist, Counselor, or Family Preservation Worker	33	41	19	22	18	24	21	24
Pediatrician or Other Family Doctor	38	38	20	22	21	22	21	24
Overnight or Longer Stay in Any Type of Hospital	31	33	15	17	19	20	19	19
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	42	46	22	25	23	25	26	27
Partial Day Hospital or Day Treatment Program	26	29	16	16	14	16	15	16
Overnight or Longer Stay in a Residential Treatment Center	23	24	10	13	11	15	15	13
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	19	18	11	9	11	12	11	10

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.54D Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2003 and 2004

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	0.94	0.85	1.64	1.57	1.56	1.47	1.64	1.62
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher ²	0.92	0.85	1.75	1.55	1.51	1.46	1.58	1.65
Mental Health Clinic or Center	0.60	0.62	1.04	1.01	0.94	1.15	1.13	1.06
In-Home Therapist, Counselor, or Family Preservation Worker	0.61	0.67	1.07	1.13	0.98	1.09	1.19	1.26
Pediatrician or Other Family Doctor	0.68	0.61	1.11	1.14	1.11	1.01	1.17	1.21
Overnight or Longer Stay in Any Type of Hospital	0.58	0.54	0.82	0.90	1.01	0.92	1.12	0.99
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	0.75	0.72	1.20	1.25	1.19	1.15	1.48	1.30
Partial Day Hospital or Day Treatment Program	0.48	0.48	0.87	0.84	0.74	0.78	0.88	0.84
Overnight or Longer Stay in a Residential Treatment Center	0.43	0.40	0.56	0.68	0.59	0.74	0.91	0.72
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.37	0.32	0.63	0.47	0.59	0.60	0.66	0.57

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.54P Source of Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Source of Treatment/Counseling	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/COUNSELING FROM ANY OF FOLLOWING 10 SOURCES¹	N/A		N/A		N/A		N/A	
Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor	0.3765		0.6946		0.9723		0.2702	
School Counselor, School Psychologist, or Having Regular Meetings with a Teacher ²	0.3056		0.6359		0.3147		0.8109	
Mental Health Clinic or Center	0.7127		0.7867		0.5017		0.8844	
In-Home Therapist, Counselor, or Family Preservation Worker	0.4007		0.4475		0.0500		0.2487	
Pediatrician or Other Family Doctor	0.4495		0.9783		0.5760		0.5241	
Overnight or Longer Stay in Any Type of Hospital	0.9635		0.6123		0.7899		0.5295	
Special Education Services While in a Regular Classroom or in a Special Classroom or Placement in a Special Program or Special School ²	0.5417		0.7948		0.8544		0.2745	
Partial Day Hospital or Day Treatment Program	0.4733		0.7848		0.9089		0.3970	
Overnight or Longer Stay in a Residential Treatment Center	0.1859		0.0194		0.3086		0.5056	
Overnight or Longer Stay in Foster Care or in a Therapeutic Foster Care Home	0.1439		0.1470		0.9959		0.2446	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis.

¹ Respondents could indicate multiple sources; thus, these response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of mental health treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.55C Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2003 and 2004

Reason for Treatment/Counseling ¹	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	103	112	56	59	55	62	56	60
Thought about Killing Self or Tried to Kill Self	38	38	18	19	22	24	24	24
Felt Depressed ²	67	71	33	33	34	43	39	44
Felt Very Afraid or Tense	42	44	22	22	23	25	26	26
Breaking Rules or "Acting Out" ³	44	51	24	30	28	29	23	27
Eating Problems	27	28	11	14	18	17	18	19
Family/Home Problems ⁴	35	33	20	18	19	19	20	18
Social/Friend Problems ⁴	28	25	15	14	17	14	14	16
School-Related Issues ⁴	28	31	15	17	17	17	16	19
Other Diagnosed Mental/Neurological Disorder ⁴	14	15	7	8	8	9	8	9
Other Reason/Reasons	47	42	27	24	23	22	27	25

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other-specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other-specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.55D Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2003 and 2004

Reason for Treatment/Counseling ¹	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Thought about Killing Self or Tried to Kill Self	0.80	0.70	1.22	1.24	1.34	1.27	1.52	1.35
Felt Depressed ²	1.08	0.96	1.90	1.74	1.70	1.67	1.82	1.64
Felt Very Afraid or Tense	0.88	0.82	1.45	1.42	1.40	1.36	1.57	1.46
Breaking Rules or "Acting Out" ³	0.89	0.93	1.51	1.71	1.62	1.49	1.47	1.53
Eating Problems	0.62	0.57	0.80	1.00	1.12	0.95	1.23	1.17
Family/Home Problems ⁴	0.73	0.67	1.33	1.18	1.15	1.10	1.29	1.14
Social/Friend Problems ⁴	0.61	0.53	1.10	0.93	1.05	0.84	0.92	1.03
School-Related Issues ⁴	0.62	0.65	1.09	1.18	1.13	0.99	1.09	1.16
Other Diagnosed Mental/Neurological Disorder ⁴	0.31	0.33	0.54	0.53	0.53	0.53	0.54	0.60
Other Reason/Reasons	0.94	0.81	1.72	1.46	1.40	1.27	1.59	1.47

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other-specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other-specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.55P Reasons for Mental Health Treatment/Counseling in the Past Year among Persons Aged 12 to 17 Who Received Mental Health Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Reason for Treatment/Counseling ¹	Total		AGE GROUP					
			12-13		14-15		16-17	
	2003	2004	2003	2004	2003	2004	2003	2004
RECEIVED TREATMENT/ COUNSELING FROM ANY OF NINE SOURCES	N/A		N/A		N/A		N/A	
Thought about Killing Self or Tried to Kill Self	0.7674		0.7889		0.6634		0.7650	
Felt Depressed ²	0.3136		0.5118		0.3090		0.1567	
Felt Very Afraid or Tense	0.6532		0.3513		0.8425		0.9248	
Breaking Rules or "Acting Out" ³	0.1574		0.0545		0.8003		0.7392	
Eating Problems	0.4428		0.4342		0.2428		0.5225	
Family/Home Problems ⁴	0.2103		0.2299		0.4446		0.8635	
Social/Friend Problems ⁴	0.4123		0.7793		0.3135		0.9567	
School-Related Issues ⁴	0.3474		0.2893		0.4843		0.1863	
Other Diagnosed Mental/Neurological Disorder ⁴	0.5790		0.9289		0.8914		0.3870	
Other Reason/Reasons	0.1809		0.3768		0.1692		0.8532	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (i.e., "special education services while in a regular classroom or in a special classroom, a special program, or in a special school") were not asked for the reasons they received these services and were therefore excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each reported source of treatment and could indicate multiple reasons for the last time they received treatment; thus, these response categories are not mutually exclusive.

² Includes other-specify responses of "feeling inadequate, hopeless, lonely, or having low self-esteem."

³ Includes other-specify responses of "difficulty controlling anger or fighting."

⁴ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.56C Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling:
Standard Errors of Numbers in Thousands, 2003 and 2004**

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	110	106	51	59	92	83
Marijuana and Hashish	88	89	42	48	73	68
Cocaine	28	28	16	18	22	21
Crack	13	10	8	8	10	7
Heroin	8	10	6	7	5	7
Hallucinogens	38	37	20	24	31	29
LSD	15	17	11	13	12	12
PCP	12	12	7	8	9	9
Ecstasy	23	24	12	15	19	19
Inhalants	50	48	27	29	42	38
Nonmedical Use of Psychotherapeutics ⁴	70	63	36	40	58	48
Pain Relievers	64	60	33	37	53	46
OxyContin [®]	--	18	--	12	--	13
Tranquilizers	34	30	20	21	28	24
Stimulants	31	28	18	18	25	22
Methamphetamine	17	17	11	12	13	13
Sedatives	16	15	9	11	13	11
ILLCIT DRUG OTHER THAN MARIJUANA³	89	81	43	48	72	63

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

**Table 6.56D Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling:
Standard Errors of Percentages, 2003 and 2004**

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	0.36	0.34	0.82	0.81	0.39	0.38
Marijuana and Hashish	0.31	0.31	0.71	0.71	0.33	0.32
Cocaine	0.11	0.11	0.30	0.32	0.11	0.11
Crack	0.05	0.04	0.15	0.13	0.05	0.04
Heroin	0.03	0.04	0.11	0.12	0.03	0.04
Hallucinogens	0.15	0.14	0.37	0.40	0.15	0.14
LSD	0.06	0.07	0.21	0.22	0.06	0.06
PCP	0.05	0.05	0.13	0.15	0.05	0.05
Ecstasy	0.09	0.09	0.24	0.27	0.10	0.09
Inhalants	0.19	0.18	0.51	0.47	0.20	0.18
Nonmedical Use of Psychotherapeutics ⁴	0.26	0.23	0.65	0.64	0.27	0.24
Pain Relievers	0.24	0.22	0.61	0.60	0.25	0.23
OxyContin [®]	--	0.07	--	0.22	--	0.07
Tranquilizers	0.13	0.12	0.38	0.36	0.14	0.12
Stimulants	0.12	0.11	0.34	0.31	0.12	0.11
Methamphetamine	0.07	0.07	0.21	0.21	0.07	0.07
Sedatives	0.06	0.06	0.18	0.20	0.06	0.05
ILLCIT DRUG OTHER THAN MARIJUANA³	0.32	0.28	0.75	0.74	0.33	0.29

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.56P Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Drug	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
ILLCIT DRUG³	0.1100		0.1104		0.0020	
Marijuana and Hashish	0.2853		0.2394		0.0289	
Cocaine	0.4183		0.9815		0.1970	
Crack	0.0454		0.2834		0.0426	
Heroin	0.4001		0.6168		0.1391	
Hallucinogens	0.5433		0.8547		0.2548	
LSD	0.7486		0.4935		0.6110	
PCP	0.3041		0.9587		0.1660	
Ecstasy	0.4924		0.7396		0.4488	
Inhalants	0.5597		0.5475		0.9922	
Nonmedical Use of Psychotherapeutics ⁴	0.3390		0.3671		0.0312	
Pain Relievers	0.4035		0.3645		0.0470	
OxyContin [®]	--		--		--	
Tranquilizers	0.1897		0.4610		0.2321	
Stimulants	0.0533		0.3378		0.0276	
Methamphetamine	0.6092		0.5027		0.0869	
Sedatives	0.9469		0.9131		0.9507	
ILLCIT DRUG OTHER THAN MARIJUANA³	0.2944		0.5107		0.0282	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

⁴ Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.57C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	111	114	50	59	94	88
Cigarettes	101	104	46	56	84	81
Smokeless Tobacco	42	47	18	22	36	40
Cigars	71	73	32	40	61	57
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	137	146	61	67	116	118
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	87	89	40	49	71	69
Cigarettes	78	84	38	46	64	65
Smokeless Tobacco	30	33	12	15	26	28
Cigars	44	46	20	27	39	38
Pipe Tobacco	18	20	9	11	15	16
Alcohol	95	96	44	47	79	81
Binge Alcohol Use ⁵	73	79	37	41	59	65
Heavy Alcohol Use ⁵	34	37	18	17	29	32

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.57D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2003 and 2004

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
PAST YEAR						
Tobacco Product ³	0.37	0.38	0.82	0.86	0.40	0.40
Cigarettes	0.36	0.35	0.79	0.83	0.38	0.37
Smokeless Tobacco	0.16	0.18	0.35	0.39	0.18	0.20
Cigars	0.26	0.27	0.59	0.63	0.28	0.28
Pipe Tobacco ⁴	--	--	--	--	--	--
Alcohol	0.42	0.41	0.92	0.83	0.46	0.47
Binge Alcohol Use ⁵	--	--	--	--	--	--
Heavy Alcohol Use ⁵	--	--	--	--	--	--
PAST MONTH						
Tobacco Product ³	0.32	0.32	0.73	0.75	0.33	0.33
Cigarettes	0.29	0.30	0.71	0.72	0.30	0.31
Smokeless Tobacco	0.12	0.13	0.23	0.26	0.13	0.14
Cigars	0.17	0.18	0.40	0.45	0.19	0.19
Pipe Tobacco	0.07	0.08	0.17	0.20	0.08	0.08
Alcohol	0.33	0.32	0.73	0.72	0.35	0.37
Binge Alcohol Use ⁵	0.27	0.29	0.66	0.67	0.28	0.32
Heavy Alcohol Use ⁵	0.13	0.14	0.34	0.29	0.15	0.16

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.57P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling; *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²	
			Received	Not Received
	2003	2004	2003	2004
PAST YEAR				
Tobacco Product ³	0.3724		0.0107	0.0032
Cigarettes	0.2989		0.0120	0.0016
Smokeless Tobacco	0.4820		0.4437	0.7985
Cigars	0.5335		0.0275	0.4239
Pipe Tobacco ⁴	--		--	--
Alcohol	0.5058		0.3325	0.1346
Binge Alcohol Use ⁵	--		--	--
Heavy Alcohol Use ⁵	--		--	--
PAST MONTH				
Tobacco Product ³	0.9564		0.0020	0.0152
Cigarettes	0.4951		0.0257	0.0056
Smokeless Tobacco	0.1877		0.1255	0.5014
Cigars	0.2087		0.0092	0.8043
Pipe Tobacco	0.5409		0.6189	0.8728
Alcohol	0.8159		0.5802	0.8539
Binge Alcohol Use ⁵	0.2617		0.7405	0.1594
Heavy Alcohol Use ⁵	0.7851		0.9751	0.8064

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.58C Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	65	67	37	39	52	52
Illicit Drug ⁴	47	48	27	30	38	37
Alcohol	52	54	29	32	42	43
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	45	45	26	29	34	34
Illicit Drug ⁴	35	37	21	24	28	26
Alcohol	32	32	20	21	23	24

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.58D Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2003 and 2004

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	0.25	0.24	0.66	0.64	0.25	0.25
Illicit Drug ⁴	0.18	0.18	0.50	0.51	0.19	0.19
Alcohol	0.20	0.20	0.54	0.53	0.20	0.21
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	0.17	0.17	0.50	0.48	0.17	0.17
Illicit Drug ⁴	0.14	0.14	0.41	0.41	0.14	0.13
Alcohol	0.12	0.12	0.39	0.36	0.11	0.12

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 6.58P Substance Dependence or Abuse in the Past Year among Persons Aged 12 to 17, by Receipt of Past Year Mental Health Treatment/Counseling; *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Dependence/Abuse	Total ¹		MENTAL HEALTH TREATMENT/COUNSELING ²			
			Received		Not Received	
	2003	2004	2003	2004	2003	2004
DEPENDENCE OR ABUSE³						
Illicit Drug or Alcohol ⁴	0.9592		0.7747		0.4749	
Illicit Drug ⁴	0.4552		0.2753		0.7764	
Alcohol	0.6366		0.8048		0.9844	
DEPENDENCE³						
Illicit Drug or Alcohol ⁴	0.6523		0.7136		0.8799	
Illicit Drug ⁴	0.4579		0.4022		0.8703	
Alcohol	0.6757		1.0000		0.8569	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 12 to 17, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling for youths is defined as having received treatment or counseling from any of 10 specific sources (e.g., private therapist, school counselor, special school program) for emotional or behavioral problems NOT caused by drug or alcohol use. Youths who answered none of the source of treatment questions with a "yes" and answered "no" four or fewer times were excluded from this analysis. See Table 6.54 for a list of the 10 specific sources of treatment/counseling.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.