

Table 7.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	340	334
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	144	130	229	217
Cigars	240	229	242	231
Pipe Tobacco	86	90	102	115
ALCOHOL	401	394	842	852
Binge Alcohol Use ²	388	383	511	529
Heavy Alcohol Use ²	270	258	244	267

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	0.19	0.18
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.24	0.22	0.13	0.12
Cigars	0.40	0.38	0.14	0.13
Pipe Tobacco	0.14	0.15	0.06	0.06
ALCOHOL	0.66	0.66	0.47	0.47
Binge Alcohol Use ²	0.64	0.64	0.29	0.29
Heavy Alcohol Use ²	0.45	0.43	0.14	0.15

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A		0.7419	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.8243		0.0644	
Cigars	0.1270		0.2894	
Pipe Tobacco	0.2874		0.6291	
ALCOHOL	0.0725		0.8464	
Binge Alcohol Use ²	0.4330		0.7695	
Heavy Alcohol Use ²	0.3739		0.6331	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	33	32
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	20	22	19	21
Cigars	30	32	25	25
Pipe Tobacco	15	16	9	10
ALCOHOL	34	37	63	67
Binge Alcohol Use ²	35	38	45	51
Heavy Alcohol Use ²	26	28	16	19

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	0.15	0.15
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.65	0.74	0.09	0.10
Cigars	0.99	1.08	0.11	0.11
Pipe Tobacco	0.51	0.53	0.04	0.04
ALCOHOL	1.10	1.22	0.29	0.30
Binge Alcohol Use ²	1.14	1.26	0.20	0.23
Heavy Alcohol Use ²	0.86	0.95	0.07	0.08

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A		0.1711	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.2531		0.3142	
Cigars	0.2597		0.2113	
Pipe Tobacco	0.4876		0.8014	
ALCOHOL	0.6191		0.7326	
Binge Alcohol Use ²	0.7009		0.0773	
Heavy Alcohol Use ²	0.5390		0.9844	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.28C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	57	62
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	44	48	34	33
Cigars	65	70	48	54
Pipe Tobacco	21	27	14	12
ALCOHOL	78	76	123	122
Binge Alcohol Use ²	89	89	104	108
Heavy Alcohol Use ²	83	81	62	59

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.28D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	0.30	0.32
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.34	0.38	0.18	0.17
Cigars	0.51	0.55	0.25	0.28
Pipe Tobacco	0.16	0.21	0.07	0.06
ALCOHOL	0.61	0.60	0.65	0.63
Binge Alcohol Use ²	0.69	0.70	0.55	0.56
Heavy Alcohol Use ²	0.65	0.64	0.33	0.30

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.28P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A		0.0669	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.1563		0.9939	
Cigars	0.0071		0.0031	
Pipe Tobacco	0.0026		1.0000	
ALCOHOL	0.6835		0.3248	
Binge Alcohol Use ²	0.9543		0.7779	
Heavy Alcohol Use ²	0.2811		0.2803	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	334	334
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	132	119	225	215
Cigars	235	214	240	228
Pipe Tobacco	81	86	100	113
ALCOHOL	389	388	809	838
Binge Alcohol Use ²	364	362	497	514
Heavy Alcohol Use ²	250	239	235	261

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2003 and 2004

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A	N/A	0.24	0.24
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.30	0.27	0.17	0.15
Cigars	0.53	0.48	0.18	0.16
Pipe Tobacco	0.18	0.19	0.07	0.08
ALCOHOL	0.87	0.88	0.59	0.60
Binge Alcohol Use ²	0.82	0.82	0.36	0.37
Heavy Alcohol Use ²	0.56	0.54	0.17	0.19

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.

Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2004 Versus 2003

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2003	2004	2003	2004
TOBACCO PRODUCT¹	N/A		0.4219	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3350		0.0501	
Cigars	0.5598		0.7539	
Pipe Tobacco	0.9000		0.6402	
ALCOHOL	0.0512		0.8955	
Binge Alcohol Use ²	0.4481		0.8830	
Heavy Alcohol Use ²	0.5935		0.4713	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2003 and 2004.