





VIRGINIA

Adolescent Behavioral Health In Brief

A Short Report from the Office of Applied Studies



Adolescence (12 to 17 years) is a critical and vulnerable stage of human development, during which males and females experience different biological, social, and cognitive changes. During this life stage, millions of adolescents experiment with substance use and engage in behaviors that can affect healthy neurological and psychological development. Understanding the behavioral health differences between adolescent males and females can help to inform public health policy and build prevention and intervention programs that strategically target the different needs of adolescent males and females.

Highlights: Adolescents in Virginia

- Approximately 56,000 (9.0 percent) of adolescents in Virginia used an illicit drug in the past month; 37,000 (6.0 percent) used marijuana, and 26,000 (4.3 percent) used an illicit drug other than marijuana.
- Adolescent females were significantly more likely than males to have used any illicit drug in the past month (11.3 v. 6.7 percent).
- Rates of illicit drug use other than marijuana in the past month were also significantly higher for adolescent females than adolescent males (5.9 v. 2.7 percent).
- 17,000 adolescent males and 27,000 adolescent females used pain relievers nonmedically in the 12 months prior to the interview.

This report provides a snapshot of behavioral health among adolescents in Virginia. National-level data on behavioral and cognitive differences between U.S. adolescent males and females is provided in a separate report entitled, *Adolescent Behavioral Health in the United States*, and is referenced at the end of this report.



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration
www.samhsa.gov

- Adolescent females in Virginia were more than three times as likely as adolescent males to have experienced a major depressive episode (MDE) in the past year (15.5 v. 4.9 percent).

The data described in the Adolescent Behavioral Health reports derive principally from national surveys conducted by the Office of Applied Studies, a component of the Substance Abuse and Mental Health Services Administration. Sources for all data used in this report appear at the end.

Adolescent Risk Perceptions


One factor that can influence adolescents' decisions to use or not use tobacco, alcohol, or illicit drugs is the extent to which youths believe these substances might cause them harm. The National Survey on Drug Use and Health (NSDUH)¹ asks respondents how much they thought people risk harming themselves physically and in other ways when they use various substances in certain amounts or frequencies.

Combined 2003–2006 NSDUH revealed that Virginia adolescents were similar to those of the Nation as a whole; that is, nationwide adolescents perceive:

- Smoking one or more packs of cigarettes per day is a greater risk than binge drinking once or twice a week.
- Binge drinking once or twice a week is perceived to be more risky than smoking marijuana once a month.

Virginia adolescents demonstrate one exception to the national patterns of the rates of risk perceptions.

- Nationwide, marijuana use once a month were significantly higher for adolescent females than for adolescent males, but in Virginia, the rates of the perception of risk associated with this behavior were similar between adolescent males and females (Chart 1).




Source: NSDUH 2003–2006.

Illicit Substance Use²

Marijuana is the most commonly used illicit drug in the United States.³ According to the combined 2003–2006 NSDUH:


- Approximately 56,000 (9.0 percent) of the 619,000 adolescents in Virginia used an illicit drug in the past month; 37,000 (6.0 percent) used marijuana, and 26,000 (4.3 percent) used an illicit drug other than marijuana (Chart 2).
- In Virginia, adolescent females were significantly more likely than males to have used any illicit drug in the past month (11.3 v. 6.7 percent). Rates of illicit drug use other than marijuana in the past month were significantly higher for adolescent females than adolescent males (5.9 v. 2.7 percent).



Source: NSDUH 2003–2006.

The misuse of pain relievers among youth is also a major public health concern.⁴


- In Virginia, 17,000 adolescent males and 27,000 adolescent females used pain relievers nonmedically in the 12 months prior to the interview (Chart 3).
- Adolescent females were significantly more likely than adolescent males to have used pain relievers nonmedically in the 12 months prior to the interview (8.9 v. 5.3 percent).



Source: NSDUH 2003–2006.

Adolescent Alcohol Use and Abuse in Virginia


- 14.5 percent (90,000) of adolescents used alcohol in the past month, and 8.4 percent (52,000) engaged in binge drinking. Binge drinking is defined as 5 or more drinks on the same occasion on at least 1 day of past 30 days.
- Rates of current alcohol use and past-month binge drinking among Virginia adolescents were similar between males and females; 14.7 percent of males and 14.3 percent of females currently used alcohol, and 8.0 percent of males and 8.8 percent of females engaged in binge drinking in the month prior to the interview (Chart 4).



Source: NSDUH 2003–2006.

Adolescent Tobacco Use

- During the month prior to the interview, 37,000 females and 41,000 males in Virginia used any form of tobacco and 31,000 males and 39,000 females used cigarettes.
- Rates of past-month cigarette use and rates of past-month tobacco use were similar between Virginia males and females (Chart 5).




Source: NSDUH 2003–2006.

Adolescent Alcohol and Illicit Drug Dependence or Abuse⁵

According to the 2003–2006 NSDUH:

- Nationwide nearly 1.5 million adolescents were dependent on or abused alcohol in the past year and more than 1.2 million adolescents were dependent or abused illicit drugs.
- Overall, the rates of past-year abuse or dependence on alcohol were significantly higher for females than males (6.0 v. 5.4 percent), but rates of past-year abuse or dependence on illicit drugs were similar between males and females.
- Rates of alcohol abuse or dependence and drug or alcohol dependence were similar between males and females; 18,000 males and 27,000 females abused or were dependent on or abused alcohol or drugs in the past year (Chart 6).



Source: NSDUH 2003–2006.

ADOLESCENT SUBSTANCE ABUSE TREATMENT

State treatment data for substance use disorders are derived from two primary sources:


(1) National Survey of Substance Abuse Treatment Services (N-SSATS),⁶ an annual 1-day census of clients in treatment and (2) the Treatment Episode Data Set (TEDS),⁷ which provides information on annual treatment admissions.

According to the 2006 N-SSATS survey:

- Virginia showed a 1-day total of 22,847 clients in treatment, the majority of whom (21,311 or 93.3 percent) were in outpatient treatment. Of the total number of clients in treatment on this date, 2,879 (12.6 percent) were under the age of 18.

According to 2003–2006 TEDS data:



- Adolescent males accounted for 67.8 percent (11,998) of the total adolescent substance abuse treatment admissions.
- Of the total male admissions, 22.6 percent were drug treatment admissions, 42.3 percent were alcohol and drug treatment, 4.9 percent were alcohol treatment, and 30.2 percent did not report type of treatment.
- Of the total adolescent female admissions, 18.2 percent were drug treatment, 35.6 percent were alcohol and drug treatment, 6.8 percent were alcohol treatment, and 39.5 percent did not report type of treatment (Chart 7).



Source: TEDS 2003–2006.

Among adolescent admissions, marijuana and alcohol were the most prevalent substances of abuse.

- Of the total adolescent male admissions, 47.2 percent (5,667) reported marijuana use and 62.5 percent (7,503) reported alcohol use.
- Of the total adolescent female admissions, 48.7 percent (2,768) reported marijuana use, and 42.4 percent (2,407) reported alcohol use.
- Further, 6.0 percent of total admissions reported cocaine use, 5.0 percent (594) of male admissions and 8.1 percent (460) of female admissions (Charts 8 and 9).





Source: TEDS 2003–2006.

UNMET NEED FOR SUBSTANCE ABUSE TREATMENT

NSDUH 2003–2006 estimates that more than 1.16 million adolescents needed but did not receive treatment for illicit drug problems and more than 1.3 million needed but did not receive treatment for alcohol problems. NSDUH defines “unmet treatment need” as an individual who meets the criteria for abuse of or dependence on illicit drugs or alcohol according to the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV), but who has not received specialty treatment for that problem in the past year.

In 2003-2006,

- Rates of unmet need for past-year alcohol and drug problems were similar between adolescent males and females in Virginia.
- 10,000 males (3.3 percent) and 15,000 females (5.1 percent) needed but did not receive treatment for drug problems (Chart 10).
- 15,000 females (4.8 percent) and 14,000 males (4.6 percent) needed but did not receive treatment for alcohol problems (Chart 11).



Source: NSDUH 2003–2006.

ADOLESCENT MENTAL HEALTH

Major Depressive Episodes


NSDUH uses the DSM-IV to define a “major depressive episode” as a period of 2 or more weeks during which the individual experiences loss of interest, depressed mood, or loss of pleasure, and four or more additional symptoms.

- MDEs impacted more than 2.1 million adolescents nationwide in the year prior to the 2004–2006 NSDUH surveys.
- Overall, adolescent females were almost three times more likely than adolescent males to experience a past-year MDE (12.7 v. 4.6 percent).

Major Depressive Episodes Among Virginia Adolescents

From 2004–2006, approximately 610,000 adolescents lived in Virginia; 312,000 males and 298,000 females.

- During this time period, the prevalence patterns of MDEs among adolescent males and females in Virginia were similar to the national patterns; adolescent females in Virginia were more than three times as likely as adolescent males to have experienced an MDE in the past year (15.5 v. 4.9 percent) (Chart 12).



Adolescent Mental Health Treatment in Virginia⁸

- In 2006, Virginia State Mental Health Authority (VSMHA) served 24,974 youth aged birth to 17 years, primarily through community programs.
- VSMHA clients aged birth to 17 years accounted for 21.9 percent of VSMHA’s total client population in 2006.
- Of the 24,974 youth served by VSMHA, 52 percent met the Federal definition for a serious emotional disturbance (SED),⁹ and 11 percent had co-occurring mental health and alcohol and other drug (AOD) disorders.



For Further Information

Adolescent Behavioral Health in the United States:

Full Report is available at:

<http://www.samhsa.gov/statesinbrief/>

A comprehensive listing of all NSDUH measures for every State is available at:

<http://oas.samhsa.gov/statesList.cfm>.

Also, information about variations in incidence and prevalence of the NSDUH substance abuse and mental health measures within each State is available at:

<http://oas.samhsa.gov/metro.htm>.

Data Sources

Facility Data: National Survey of Substance Abuse Treatment Services (N-SSATS)–2006 is available at: <http://www.dasis.samhsa.gov>.

Center for Mental Health Services Uniform Reporting System Output Tables 2006 is available at: <http://mentalhealth.samhsa.gov/cmhs/MentalHealthStatistics/URS2006.asp>.

Substance Abuse Treatment Data: Treatment Episode Data Set–Concatenated File–is available from the Substance Abuse and Mental Health Data Archive: <http://www.icpsr.umich.edu/SDA/SAMHDA>.

Mental Health Treatment Data: Center for Mental Health Services Uniform Reporting System Output Tables 2006 is available at: <http://mentalhealth.samhsa.gov/cmhs/MentalHealthStatistics/URS2006.asp>.

¹NSDUH is directed by the Substance Abuse and Mental Health Services Administration and provides information on the incidence and prevalence of substance use in the population and the problems associated with use. The survey collects information on the sociodemographic characteristics of users, perceptions of risk and availability, and mental health issues. Since 1999, the NSDUH sample has been designed to provide State-level estimates, based on about 67,500 respondents per year.

²NSDUH defines “illicit drugs” to include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type drugs used nonmedically. “Nonmedical” use is defined as use not prescribed for the respondent by a physician or used only for the experience or feeling the drug(s) caused. Nonmedical use of any prescription-type pain reliever, sedative, stimulant, or tranquilizer does not include over-the-counter drugs. Non-medical use of stimulants includes methamphetamine use.

³Substance Abuse and Mental Health Services Administration. (2008). *Results from the 2007 National Survey on Drug Use and Health: National Findings* (Office of Applied Studies, NSDUH Series H-34, DHHS Publication No. SMA 08-4343). Rockville, MD.

⁴The White House, Executive Office of the President (2009). *National Drug Control Policy: 2009 Annual Report* (Chapter 1). [Available at: <http://www.whitehousedrugpolicy.gov/publications/policy/ndcs09/chapter1.pdf>]

⁵Questions in NSDUH are used to classify persons as being dependent on or abusing specific substances based on criteria specified in the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV) (American Psychiatric Association, 1994).

⁶The National Survey of Substance Abuse Treatment Services (N-SSATS) is designed to collect information from all facilities in the United States, both public and private, that provide substance abuse treatment. N-SSATS does not collect information from the following three types of facilities: nontreatment halfway houses; jails, prisons, or other organizations that treat incarcerated clients exclusively; and solo practitioners.

⁷TEDS is an admissions-based system, and TEDS admissions do not represent individuals. For example, an individual admitted to treatment twice within a calendar year would be counted as two admissions. TEDS collects information on up to three substances of abuse that lead to the treatment episode. These are not necessarily a complete enumeration of all drugs used at the time of admission.

⁸The Uniform Reporting System (URS) consists of 21 standardized tables (12 basic tables and 9 developmental tables) that State mental health agencies (SMHA) submit each December in their Community Mental Health Services Block Grant Implementation Report to the Substance Abuse and Mental Health Services Administration’s Center for Mental Health Services. The URS data submitted by the States has been used to create 14 different output tables that show performance on issues of access, appropriateness, outcomes, and system management. The intent of the URS tables is to allow both (1) the tracking of individual State performance over time, and (2) the aggregation of State information to develop a national picture of the public mental health systems of the States.

⁹The Federal definition of “serious emotional disturbance” (SED) is provided by the Individuals with Disabilities Education Act (IDEA).