

Table 1.15A (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2010 and 2011

Dependence or Abuse Status	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)
DEPENDENCE				
None	36,003	36,733	7,508	7,397
Both Illicit Drugs and Alcohol	648	695	277	348
Illicit Drugs or Alcohol	5,414 ^a	4,649	1,826	1,613
Illicit Drugs	2,230	2,029	869	809
Alcohol	3,833 ^a	3,314	1,234	1,152
DEPENDENCE OR ABUSE				
None	33,552	34,568	6,946	6,888
Both Illicit Drugs and Alcohol	1,324	1,167	503	476
Illicit Drugs or Alcohol	7,865 ^b	6,813	2,388	2,123
Illicit Drugs	3,007 ^a	2,492	1,052	951
Alcohol	6,182 ^a	5,489	1,838	1,649

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15B (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Percentages, 2010 and 2011

Dependence or Abuse Status	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)
DEPENDENCE				
None	16.5	16.5	3.4	3.3
Both Illicit Drugs and Alcohol	62.7	66.7	26.8	33.4
Illicit Drugs or Alcohol	48.7	45.2	16.4	15.7
Illicit Drugs	52.7	51.6	20.5	20.6
Alcohol	48.3	44.8	15.6	15.6
DEPENDENCE OR ABUSE				
None	16.1	16.2	3.3	3.2
Both Illicit Drugs and Alcohol	54.9	52.8	20.8	21.5
Illicit Drugs or Alcohol	38.5	36.1	11.7	11.2
Illicit Drugs	50.2	46.4	17.6	17.7
Alcohol	36.7	34.9	10.9	10.5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16A (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2010 and 2011

Dependence or Abuse Status	Total		Any	Any	Serious	Serious	Moderate	Moderate	Mild	Mild	No	No
	(2010)	(2011)	Mental Illness (2010)	Mental Illness (2011)								
DEPENDENCE												
Both Illicit Drugs and Alcohol	1,033	1,042	648	695	277	348	152	153	218	194	385	347
Illicit Drugs or Alcohol	11,128	10,290	5,414 ^a	4,649	1,826	1,613	1,445 ^a	1,087	2,144	1,948	5,714	5,641
Illicit Drugs	4,228	3,934	2,230	2,029	869	809	590	489	771	731	1,999	1,905
Alcohol	7,933	7,397	3,833 ^a	3,314	1,234	1,152	1,008 ^a	752	1,591	1,411	4,100	4,083
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	2,413	2,212	1,324	1,167	503	476	310	291	511	401	1,089	1,044
Illicit Drugs or Alcohol	20,435 ^b	18,887	7,865 ^b	6,813	2,388	2,123	2,082 ^b	1,605	3,395	3,085	12,570	12,073
Illicit Drugs	5,990 ^a	5,374	3,007 ^a	2,492	1,052	951	770	598	1,185	943	2,983	2,882
Alcohol	16,858 ^a	15,724	6,182 ^a	5,489	1,838	1,649	1,623 ^a	1,297	2,721	2,543	10,676	10,235

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16B (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2010 and 2011

Dependence or Abuse Status	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.5	0.4	1.6	1.7	3.0	3.9	1.5	1.5	1.0	0.9	0.2	0.2
Illicit Drugs or Alcohol	4.9 ^a	4.4	13.1 ^a	11.2	19.6	17.9	14.6 ^b	10.5	9.6	8.8	3.0	2.9
Illicit Drugs	1.8	1.7	5.4	4.9	9.3	9.0	6.0	4.7	3.5	3.3	1.1	1.0
Alcohol	3.5	3.2	9.3 ^a	8.0	13.2	12.8	10.2 ^a	7.3	7.2	6.4	2.2	2.1
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	1.1	1.0	3.2	2.8	5.4	5.3	3.1	2.8	2.3	1.8	0.6	0.5
Illicit Drugs or Alcohol	8.9 ^b	8.1	19.0 ^b	16.5	25.6	23.6	21.1 ^b	15.5	15.3	14.0	6.7	6.3
Illicit Drugs	2.6 ^a	2.3	7.3 ^a	6.0	11.3	10.5	7.8	5.8	5.3	4.3	1.6	1.5
Alcohol	7.4 ^a	6.8	14.9 ^a	13.3	19.7	18.3	16.4 ^a	12.6	12.2	11.5	5.7	5.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17A (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	5,990 ^a	5,374	3,007 ^a	2,492	1,052	951	770	598	1,185	943	2,983	2,882
18-25	2,677	2,574	1,123 ^a	970	322	306	316 ^a	238	486	426	1,554	1,604
26-49	2,757 ^a	2,235	1,559 ^a	1,207	608	525	317	294	633 ^a	388	1,198	1,028
50 or Older	555	565	325	315	123	119	136	67	66	129	230	250
MALE	3,894	3,615	1,645	1,336	454	423	423	343	769	571	2,249	2,278
18-25	1,668	1,726	563	490	106	124	178 ^a	114	279	252	1,105	1,236
26-49	1,833	1,509	919	676	320	249	151	178	449 ^a	249	914	833
50 or Older	393	380	163	170	*	*	*	*	41	69	230	210
FEMALE	2,096 ^a	1,760	1,362	1,156	599	528	347	256	416	372	734	604
18-25	1,009 ^a	848	560	480	215	182	138	124	207	173	449	368
26-49	924	726	640	531	289	276	166	116	185	139	285	195
50 or Older	162	185	162	145	95	70	43	*	25	60	*	40

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17B (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	2.6 ^a	2.3	7.3 ^a	6.0	11.3	10.5	7.8	5.8	5.3	4.3	1.6	1.5
18-25	7.9	7.5	18.3 ^b	15.3	24.1	23.6	19.0 ^a	14.3	15.4 ^a	12.5	5.6	5.7
26-49	2.8 ^a	2.3	7.5	6.1	11.9	10.6	6.5	5.6	6.0 ^a	4.0	1.5	1.3
50 or Older	0.6	0.6	2.2	2.1	4.3	4.3	4.1	1.9	0.8	1.5	0.3	0.3
MALE	3.5	3.2	10.0	8.4	13.5	13.1	11.4	9.4	8.2	6.3	2.4	2.4
18-25	9.7	10.0	24.3 ^a	19.7	24.0	27.8	29.6 ^a	19.8	21.9	17.2	7.4	8.4
26-49	3.8	3.1	11.1	9.0	16.7	14.7	8.7	9.7	9.6	6.3	2.3	2.1
50 or Older	0.9	0.8	2.8	2.9	*	*	*	*	1.2	1.9	0.6	0.5
FEMALE	1.8 ^a	1.5	5.5	4.5	10.0	9.1	5.6	3.8	3.3	2.9	0.8	0.6
18-25	6.0 ^b	5.0	14.6	12.4	24.2	21.4	12.9	11.4	11.0	9.0	3.5 ^a	2.8
26-49	1.9	1.5	5.2	4.3	9.0	8.5	5.3	3.4	3.1	2.4	0.8	0.5
50 or Older	0.3	0.3	1.8	1.6	5.1	4.1	2.2	*	0.5	1.1	*	0.1

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18A (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	16,858 ^a	15,724	6,182 ^a	5,489	1,838	1,649	1,623 ^a	1,297	2,721	2,543	10,676	10,235
18-25	5,342 ^a	4,936	1,629	1,577	422	370	478	438	729	768	3,713 ^a	3,359
26-49	8,345	7,867	3,600 ^b	2,962	1,184	909	893	686	1,523	1,367	4,745	4,905
50 or Older	3,171	2,921	952	950	232	370	252	173	469	407	2,219	1,971
MALE	11,253 ^b	10,040	3,278 ^a	2,800	898	754	751	632	1,629	1,415	7,975	7,239
18-25	3,377 ^b	2,950	774	752	167	151	225	181	382	420	2,603 ^b	2,198
26-49	5,638	5,156	1,993 ^b	1,503	605	420	437	303	950	780	3,646	3,653
50 or Older	2,238	1,933	512	545	*	*	*	*	297	215	1,726	1,388
FEMALE	5,605	5,685	2,903	2,688	940	895	871	666	1,092	1,128	2,702	2,996
18-25	1,965	1,986	855	825	255	220	253	257	347	348	1,110	1,161
26-49	2,707	2,711	1,607	1,459	579	488	456	383	573	587	1,099	1,252
50 or Older	933	988	441	405	106	187	163 ^a	25	172	192	493	583

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18B (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	7.4 ^a	6.8	14.9 ^a	13.3	19.7	18.3	16.4 ^a	12.6	12.2	11.5	5.7	5.4
18-25	15.7 ^b	14.4	26.5	24.8	31.6	28.6	28.7	26.3	23.1	22.6	13.3 ^a	12.0
26-49	8.5	8.0	17.4 ^a	14.9	23.1	18.4	18.3 ^a	13.1	14.3	14.0	6.1	6.3
50 or Older	3.3	2.9	6.5	6.3	8.1	13.4	7.6	5.0	5.6	4.6	2.7	2.3
MALE	10.1 ^b	9.0	19.9	17.6	26.8	23.4	20.3	17.3	17.3	15.7	8.4 ^a	7.5
18-25	19.5 ^b	17.2	33.4	30.3	37.7	33.8	37.4	31.4	30.1	28.7	17.4 ^b	15.0
26-49	11.6	10.7	24.0	20.1	31.7	24.9	25.1	16.6	20.4	19.7	9.0	9.0
50 or Older	5.0	4.1	8.8	9.2	*	*	*	*	8.5	5.9	4.4	3.4
FEMALE	4.7	4.7	11.6	10.5	15.7	15.5	14.1 ^a	9.9	8.5	8.7	2.9	3.1
18-25	11.7	11.6	22.3	21.3	28.6	25.8	23.7	23.6	18.5	18.0	8.6	8.8
26-49	5.4	5.4	13.0	11.7	18.0	15.0	14.5	11.3	9.6	10.1	2.9	3.3
50 or Older	1.8	1.8	5.0	4.4	5.7	11.1	8.2 ^a	1.1	3.5	3.6	1.1	1.3

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.19A (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	20,435 ^b	18,887	7,865 ^b	6,813	2,388	2,123	2,082 ^b	1,605	3,395	3,085	12,570	12,073
18-25	6,803 ^a	6,369	2,182	2,060	570	529	621	546	991	985	4,621	4,308
26-49	10,000	9,217	4,477 ^b	3,629	1,508 ^a	1,153	1,073	854	1,897	1,622	5,522	5,588
50 or Older	3,632	3,300	1,206	1,124	310	441	388	205	508	478	2,426	2,177
MALE	13,576 ^b	12,105	4,131 ^a	3,459	1,069	932	989	804	2,073	1,723	9,445	8,646
18-25	4,284 ^b	3,879	1,025	984	207	207	302 ^a	228	515	549	3,259 ^a	2,894
26-49	6,739 ^a	6,062	2,487 ^b	1,863	736 ^a	516	504	412	1,246 ^a	935	4,252	4,199
50 or Older	2,553	2,165	619	612	*	*	*	*	311	239	1,934	1,553
FEMALE	6,859	6,781	3,734	3,354	1,319	1,191	1,093 ^a	801	1,322	1,362	3,125	3,427
18-25	2,519	2,490	1,157	1,076	363	322	319	318	475	436	1,362	1,414
26-49	3,261	3,156	1,990	1,766	772	637	569	442	650	688	1,271	1,390
50 or Older	1,079	1,135	586	512	185	233	205 ^a	41	196	238	493	623

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.19B (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	8.9 ^b	8.1	19.0 ^b	16.5	25.6	23.6	21.1 ^b	15.5	15.3	14.0	6.7	6.3
18-25	20.0 ^a	18.6	35.5 ^a	32.4	42.7	40.8	37.3	32.9	31.5	29.0	16.6 ^a	15.4
26-49	10.1	9.4	21.7 ^b	18.2	29.4 ^a	23.3	22.0 ^a	16.3	17.8	16.6	7.1	7.2
50 or Older	3.8	3.3	8.2	7.4	10.8	16.0	11.7	5.9	6.0	5.4	3.0	2.6
MALE	12.2 ^b	10.8	25.1 ^a	21.8	31.9	29.0	26.8	22.1	22.0	19.1	10.0 ^a	9.0
18-25	24.8 ^b	22.6	44.2	39.6	46.6	46.6	50.3 ^a	39.7	40.6	37.5	21.8 ^a	19.7
26-49	13.8	12.6	29.9 ^a	24.9	38.6	30.5	28.9	22.5	26.8	23.6	10.5	10.3
50 or Older	5.7	4.6	10.6	10.3	*	*	*	*	8.9	6.6	4.9	3.8
FEMALE	5.8	5.6	15.0	13.2	22.0	20.6	17.7 ^b	12.0	10.3	10.5	3.3	3.6
18-25	15.0	14.5	30.2	27.8	40.8	37.8	29.9	29.3	25.3	22.6	10.5	10.7
26-49	6.5	6.3	16.1	14.2	23.9	19.5	18.1	13.0	10.9	11.9	3.4	3.7
50 or Older	2.1	2.1	6.7	5.6	9.9	13.8	10.4 ^b	1.8	4.0	4.5	1.1	1.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.