

**Table 1.31A (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Numbers in Thousands, 2010 and 2011**

Location of Treatment/Counseling <sup>1</sup>			Any	Any	Serious	Serious	Moderate	Moderate	Mild	Mild	No	No
	Total	Total	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental
	(2010)	(2011)	Illness	Illness	Illness	Illness	Illness	Illness	Illness	Illness	Illness	Illness
	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)
Outpatient Mental Health Clinic or Center	3,320	3,645	2,470	2,717	1,160	1,241	695	759	615	717	849	928
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	8,482	8,559	5,312	5,324	2,319	2,090	1,312	1,299	1,681	1,934	3,170	3,236
Doctor's Office - Not Part of a Clinic	3,157	3,014	2,042	2,072	799	898	703	619	539	555	1,115	942
Outpatient Medical Clinic	1,007	1,161	657	741	321	322	151	194	185	226	350	420
Partial Day Hospital or Day Treatment Program	283	284	228	227	151	146	41	36	35	44	56	58
School or University Setting/Clinic/Center <sup>2</sup>	81	84	44	54	15	16	17	16	12	23	37	30
Some Other Place <sup>3</sup>	325	373	234	270	49 <sup>a</sup>	162	101	45	85	63	91	103

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.31B (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Percentages, 2010 and 2011**

Location of Treatment/Counseling <sup>1</sup>			Any	Any	Serious	Serious	Moderate	Moderate	Mild	Mild	No	No
	Total	Total	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental	Mental
	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)	(2010)	(2011)
Outpatient Mental Health Clinic or Center	22.0	23.7	25.7	27.6	29.5	31.6	25.9	28.4	20.6	22.0	15.6	16.8
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	56.3	55.7	55.3	54.0	59.0	53.3	48.9	48.6	56.3	59.4	58.1	58.7
Doctor's Office - Not Part of a Clinic	21.0	19.6	21.3	21.0	20.3	22.9	26.2	23.2	18.1	17.0	20.4	17.1
Outpatient Medical Clinic	6.7	7.6	6.8	7.5	8.2	8.2	5.6	7.2	6.2	6.9	6.4	7.6
Partial Day Hospital or Day Treatment Program	1.9	1.9	2.4	2.3	3.8	3.7	1.5	1.4	1.2	1.4	1.0	1.0
School or University Setting/Clinic/Center <sup>2</sup>	0.5	0.6	0.5	0.6	0.4	0.4	0.6	0.6	0.4	0.7	0.7	0.5
Some Other Place <sup>3</sup>	2.2	2.4	2.5	2.8	1.3 <sup>a</sup>	4.1	3.8	1.7	2.9	2.0	1.7	1.9

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.