

**Table 1.32A (REVISED) Type and Location of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2010 and 2011**

Type and Location of Mental Health Treatment/Counseling	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
<b>ANY MENTAL HEALTH SERVICES<sup>1</sup></b>	31,490	31,642	17,510	16,847	6,283	5,838	4,636	4,726	6,591	6,283	13,980	14,795
<b>Outpatient</b>	15,146	15,477	9,660	9,902	3,961	3,962	2,699	2,672	3,001	3,268	5,486	5,576
Outpatient Mental Health Clinic or Center	3,320	3,645	2,470	2,717	1,160	1,241	695	759	615	717	849	928
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	8,482	8,559	5,312	5,324	2,319	2,090	1,312	1,299	1,681	1,934	3,170	3,236
Doctor's Office - Not Part of a Clinic	3,157	3,014	2,042	2,072	799	898	703	619	539	555	1,115	942
Outpatient Medical Clinic	1,007	1,161	657	741	321	322	151	194	185	226	350	420
Partial Day Hospital or Day Treatment Program	283	284	228	227	151	146	41	36	35	44	56	58
School or University Setting/Clinic/Center <sup>2</sup>	81	84	44	54	15	16	17	16	12	23	37	30
Some Other Place <sup>3</sup>	325	373	234	270	49 <sup>a</sup>	162	101	45	85	63	91	103
<b>Inpatient</b>	1,752	1,779	1,102	1,346	622	796	199	296	281	254	650	433
<b>Prescription Medication</b>	26,671	26,777	15,288	14,700	5,683	5,242	4,086	4,100	5,519	5,359	11,383	12,077
<b>DID NOT RECEIVE MENTAL HEALTH SERVICES</b>	197,171 <sup>b</sup>	200,369	23,823	24,453	3,031	3,155	5,188	5,591	15,604	15,707	173,347	175,916

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple service sources; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.32B (REVISED) Type and Location of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2010 and 2011**

Type and Location of Mental Health Treatment/Counseling	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
<b>ANY MENTAL HEALTH SERVICES<sup>1</sup></b>	13.8	13.6	42.4	40.8	67.5	64.9	47.2	45.8	29.7	28.6	7.5	7.8
<b>Outpatient</b>	6.6	6.7	23.4	24.0	42.5	44.1	27.6	26.0	13.5	14.9	2.9	2.9
Outpatient Mental Health Clinic or Center	1.5	1.6	6.0	6.6	12.5	13.9	7.1	7.4	2.8	3.3	0.5	0.5
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	3.7	3.7	12.9	12.9	25.0	23.4	13.4	12.6	7.6	8.8	1.7	1.7
Doctor's Office - Not Part of a Clinic	1.4	1.3	5.0	5.0	8.6	10.0	7.2	6.0	2.4	2.5	0.6	0.5
Outpatient Medical Clinic	0.4	0.5	1.6	1.8	3.5	3.6	1.5	1.9	0.8	1.0	0.2	0.2
Partial Day Hospital or Day Treatment Program	0.1	0.1	0.6	0.6	1.6	1.6	0.4	0.4	0.2	0.2	0.0	0.0
School or University Setting/Clinic/Center <sup>2</sup>	0.0	0.0	0.1	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.0	0.0
Some Other Place <sup>3</sup>	0.1	0.2	0.6	0.7	0.5 <sup>a</sup>	1.8	1.0	0.4	0.4	0.3	0.0	0.1
<b>Inpatient</b>	0.8	0.8	2.7	3.3	6.7	8.8	2.0	2.9	1.3	1.2	0.3	0.2
<b>Prescription Medication</b>	11.7	11.5	36.9	35.6	61.0	58.2	41.4	39.7	24.9	24.3	6.1	6.3
<b>DID NOT RECEIVE MENTAL HEALTH SERVICES</b>	86.2	86.4	57.6	59.2	32.5	35.1	52.8	54.2	70.3	71.4	92.5	92.2

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple service sources; thus, these response categories are not mutually exclusive.

<sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

<sup>3</sup> Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.