

**Table 1.67A Serious Psychological Distress (SPD) in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	10,196	10,508	11,018	10,888
<b>AGE</b>				
18-25	2,507	2,574	2,694	2,635
26-49	5,055	5,203	4,978	5,037
50 or Older	2,634	2,731	3,346	3,215
<b>GENDER</b>				
Male	4,287	4,056	4,373	4,521
Female	5,909	6,452	6,645	6,367
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	8,697	9,021	9,225	9,457
White	6,761	7,340	7,159	7,172
Black or African American	1,352	1,248	1,508	1,490
American Indian or Alaska Native	43	43	41	101
Native Hawaiian or Other Pacific Islander	*	32	50	*
Asian	291	191	259	311
Two or More Races	201	168	208	260
Hispanic or Latino	1,498	1,487	1,793	1,431

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 1.67B Serious Psychological Distress (SPD) in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	4.5	4.6	4.8	4.7
<b>AGE</b>				
18-25	7.6	7.7	7.9	7.7
26-49	5.1	5.2	5.1	5.1
50 or Older	2.9	2.9	3.5	3.2
<b>GENDER</b>				
Male	3.9	3.7	3.9	4.0
Female	5.1	5.5	5.6	5.3
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	4.5	4.6	4.7	4.8
White	4.4	4.7	4.6	4.6
Black or African American	5.2	4.7	5.7	5.6
American Indian or Alaska Native	4.5	4.0	3.9	7.8
Native Hawaiian or Other Pacific Islander	*	4.3	7.5	*
Asian	3.0	1.9	2.5	2.8
Two or More Races	8.0	6.9	7.8	7.9
Hispanic or Latino	4.9	4.8	5.6 <sup>a</sup>	4.2

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.