

Table 1.13C (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2010 and 2011

Drug	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
ILLCIT DRUGS¹	631	589	343	319	177	176	152	152	230	197	554	524
Marijuana and Hashish	548	503	273	261	142	141	130	123	176	165	478	456
Cocaine	202	167	119	92	58	60	49	41	88	57	165	135
Crack	92	68	71	48	45	28	31	29	42	27	59	49
Heroin	75	71	54	59	37	35	14	28	36	38	51	31
Hallucinogens	170	138	94	83	60	54	42	36	54	50	133	110
LSD	69	57	45	32	27	18	28	14	20	22	48	47
PCP	23	19	21	15	20	7	3	9	3	9	9	12
Ecstasy	121	103	68	65	34	46	34	29	44	37	94	80
Inhalants	109	99	54	61	35	32	29	39	28	32	94	77
Nonmedical Use of Psychotherapeutics ^{2,3}	389	357	240	216	132	125	119	107	157	134	317	284
Pain Relievers	329	297	210	180	122	110	109	93	129	104	258	242
OxyContin [®]	114	102	87	75	69	35	42	41	31	51	71	68
Tranquilizers	205	207	148	135	71	80	94	70	96	86	154	155
Stimulants ³	155	149	91	100	53	63	43	42	59	63	117	104
Methamphetamine ³	103	118	53	80	33	50	22	36	35	51	85	81
Sedatives	103	63	62	51	43	36	23	30	39	19	82	39
ILLCIT DRUGS OTHER THAN MARIJUANA¹	433	397	265	232	143	135	127	115	173	146	345	321

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13D (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2010 and 2011

Drug	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
ILLCIT DRUGS¹	0.28	0.25	0.74	0.69	1.61	1.63	1.46	1.34	0.95	0.85	0.28	0.25
Marijuana and Hashish	0.24	0.22	0.61	0.58	1.38	1.39	1.29	1.13	0.76	0.73	0.24	0.22
Cocaine	0.09	0.07	0.29	0.22	0.61	0.65	0.50	0.40	0.40	0.26	0.09	0.07
Crack	0.04	0.03	0.17	0.12	0.48	0.31	0.32	0.28	0.19	0.12	0.03	0.03
Heroin	0.03	0.03	0.13	0.14	0.39	0.38	0.15	0.27	0.16	0.17	0.03	0.02
Hallucinogens	0.07	0.06	0.23	0.20	0.64	0.59	0.43	0.35	0.24	0.23	0.07	0.06
LSD	0.03	0.02	0.11	0.08	0.29	0.20	0.29	0.13	0.09	0.10	0.03	0.02
PCP	0.01	0.01	0.05	0.04	0.22	0.08	0.03	0.09	0.02	0.04	0.00	0.01
Ecstasy	0.05	0.04	0.16	0.16	0.36	0.50	0.34	0.28	0.20	0.17	0.05	0.04
Inhalants	0.05	0.04	0.13	0.15	0.37	0.35	0.29	0.38	0.13	0.15	0.05	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.17	0.15	0.54	0.49	1.29	1.26	1.15	0.98	0.67	0.59	0.17	0.15
Pain Relievers	0.14	0.13	0.48	0.41	1.21	1.12	1.05	0.86	0.57	0.47	0.14	0.12
OxyContin [®]	0.05	0.04	0.21	0.18	0.72	0.39	0.42	0.39	0.14	0.23	0.04	0.04
Tranquilizers	0.09	0.09	0.35	0.32	0.75	0.86	0.92	0.66	0.42	0.39	0.08	0.08
Stimulants ³	0.07	0.06	0.22	0.24	0.57	0.69	0.43	0.40	0.26	0.28	0.06	0.05
Methamphetamine ³	0.04	0.05	0.13	0.19	0.36	0.55	0.23	0.34	0.16	0.23	0.05	0.04
Sedatives	0.04	0.03	0.15	0.12	0.45	0.40	0.23	0.29	0.18	0.09	0.04	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.19	0.17	0.60	0.53	1.39	1.34	1.23	1.04	0.75	0.64	0.18	0.16

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13P (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Drug	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
ILLCIT DRUGS¹	0.1999		0.4471		0.6922		0.5769		0.7159		0.3142	
Marijuana and Hashish	0.9121		0.8497		0.8003		0.6638		0.9078		0.9560	
Cocaine	0.0068		0.0458		0.4506		0.0145		0.0512		0.0573	
Crack	0.0169		0.0359		0.3756		0.0803		0.2511		0.2257	
Heroin	0.7385		0.3521		0.8059		0.2424		0.5898		0.0910	
Hallucinogens	0.0543		0.1839		0.9520		0.0421		0.5174		0.1475	
LSD	0.6840		0.3596		0.9803		0.0301		0.5920		0.7706	
PCP	0.5534		0.9166		0.5049		0.4219		0.3110		0.3030	
Ecstasy	0.1922		0.4329		0.5532		0.1331		0.4848		0.2809	
Inhalants	0.3631		0.7703		0.6765		0.9713		0.9284		0.3706	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.0039		0.0573		0.2981		0.3636		0.2057		0.0318	
Pain Relievers	0.0071		0.0675		0.1231		0.8562		0.1855		0.0527	
OxyContin [®]	0.0830		0.3093		0.0188		0.9917		0.2904		0.1686	
Tranquilizers	0.0611		0.0940		0.9273		0.1534		0.1786		0.3496	
Stimulants ³	0.3191		0.9022		0.2925		0.5563		0.4493		0.2467	
Methamphetamine ³	0.7343		0.1542		0.1593		0.2427		0.8813		0.4943	
Sedatives	0.0025		0.0753		0.2779		0.8458		0.0911		0.0183	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.0038		0.0304		0.3082		0.1449		0.1892		0.0422	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14C (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
PAST YEAR												
Tobacco Products ¹	906	895	484	458	235	212	227	219	344	313	982	944
Cigarettes	828	853	442	434	226	208	215	203	304	293	845	844
Smokeless Tobacco	345	319	145	118	68	52	67	36	109	100	325	295
Cigars	504	498	224	201	116	99	105	105	162	140	507	491
Alcohol	953	977	644	632	274	272	281	293	470	450	1,750	1,707
PAST MONTH												
Tobacco Products ¹	840	845	457	433	220	206	217	206	313	288	867	839
Cigarettes	772	814	422	400	214	205	205	189	281	265	770	777
Smokeless Tobacco	310	275	131	104	59	40	57	29	100	92	291	251
Cigars	358	369	169	157	81	68	86	87	126	112	330	343
Pipe Tobacco	147	143	88	65	36	26	40	49	69	34	119	129
Alcohol	974	1,006	556	542	233	227	237	246	407	400	1,502	1,516
Binge Alcohol Use ²	748	736	348	344	163	161	154	161	255	232	814	806
Heavy Alcohol Use ²	421	423	187	201	91	95	82	94	129	128	402	389

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14D (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2010 and 2011

Substance	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
PAST YEAR												
Tobacco Products ¹	0.40	0.38	0.87	0.85	1.80	1.77	1.75	1.63	1.21	1.16	0.43	0.42
Cigarettes	0.36	0.37	0.83	0.83	1.79	1.75	1.69	1.61	1.13	1.14	0.39	0.39
Smokeless Tobacco	0.15	0.14	0.34	0.28	0.72	0.58	0.68	0.35	0.47	0.44	0.17	0.15
Cigars	0.22	0.21	0.52	0.46	1.16	1.04	1.03	0.96	0.71	0.62	0.24	0.24
Alcohol	0.42	0.42	0.87	0.92	1.83	1.73	1.70	1.67	1.20	1.27	0.45	0.46
PAST MONTH												
Tobacco Products ¹	0.37	0.36	0.86	0.84	1.77	1.76	1.74	1.65	1.18	1.11	0.40	0.38
Cigarettes	0.34	0.35	0.82	0.80	1.74	1.74	1.67	1.57	1.12	1.07	0.37	0.37
Smokeless Tobacco	0.14	0.12	0.31	0.25	0.62	0.44	0.58	0.28	0.43	0.41	0.15	0.13
Cigars	0.16	0.16	0.40	0.37	0.83	0.74	0.85	0.82	0.56	0.50	0.17	0.17
Pipe Tobacco	0.06	0.06	0.21	0.16	0.38	0.29	0.40	0.47	0.31	0.15	0.06	0.07
Alcohol	0.42	0.43	0.93	0.92	1.79	1.77	1.80	1.77	1.31	1.27	0.47	0.48
Binge Alcohol Use ²	0.33	0.32	0.72	0.73	1.54	1.55	1.46	1.42	1.04	0.95	0.36	0.36
Heavy Alcohol Use ²	0.18	0.18	0.43	0.46	0.94	1.01	0.82	0.89	0.57	0.56	0.20	0.19

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14P (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
PAST YEAR												
Tobacco Products ¹	0.0437		0.0420		0.8932		0.3742		0.0376		0.2146	
Cigarettes	0.0575		0.0736		0.8346		0.4827		0.0692		0.2183	
Smokeless Tobacco	0.1199		0.0171		0.4537		0.0057		0.2713		0.5125	
Cigars	0.0355		0.0425		0.6261		0.2934		0.0750		0.1651	
Alcohol	0.8354		0.2770		0.2149		0.7678		0.0421		0.8055	
PAST MONTH												
Tobacco Products ¹	0.0482		0.1131		0.5941		0.2439		0.0850		0.1657	
Cigarettes	0.0466		0.1320		0.7274		0.2977		0.1153		0.1466	
Smokeless Tobacco	0.0817		0.0564		0.3602		0.0233		0.4728		0.2890	
Cigars	0.2026		0.1337		0.5952		0.2199		0.4070		0.5084	
Pipe Tobacco	0.7186		0.3028		0.3202		0.3323		0.0896		0.8027	
Alcohol	0.9236		0.1512		0.9541		0.4233		0.1524		0.4632	
Binge Alcohol Use ²	0.3614		0.1811		0.8088		0.5343		0.1189		0.7089	
Heavy Alcohol Use ²	0.0322		0.4370		0.9207		0.4386		0.4876		0.0511	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.