

Table 1.15C (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2010 and 2011

Dependence or Abuse Status	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)
DEPENDENCE				
None	744	761	330	300
Both Illicit Drugs and Alcohol	73	87	46	63
Illicit Drugs or Alcohol	247	206	138	122
Illicit Drugs	150	127	96	86
Alcohol	208	182	108	107
DEPENDENCE OR ABUSE				
None	719	737	322	288
Both Illicit Drugs and Alcohol	98	99	63	66
Illicit Drugs or Alcohol	287	251	155	145
Illicit Drugs	176	136	104	91
Alcohol	251	229	130	130

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15D (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2010 and 2011

Dependence or Abuse Status	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)
DEPENDENCE				
None	0.30	0.30	0.15	0.13
Both Illicit Drugs and Alcohol	3.98	3.69	3.78	4.61
Illicit Drugs or Alcohol	1.52	1.44	1.13	1.05
Illicit Drugs	2.23	2.14	1.98	1.93
Alcohol	1.85	1.75	1.29	1.27
DEPENDENCE OR ABUSE				
None	0.31	0.31	0.15	0.13
Both Illicit Drugs and Alcohol	2.58	2.75	2.26	2.58
Illicit Drugs or Alcohol	1.09	1.09	0.72	0.71
Illicit Drugs	1.85	1.79	1.54	1.53
Alcohol	1.20	1.21	0.75	0.77

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.15P (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Dependence or Abuse Status	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)
DEPENDENCE				
None	0.9677		0.5605	
Both Illicit Drugs and Alcohol	0.4672		0.2707	
Illicit Drugs or Alcohol	0.0902		0.6295	
Illicit Drugs	0.7081		0.9950	
Alcohol	0.1598		0.9908	
DEPENDENCE OR ABUSE				
None	0.8076		0.6038	
Both Illicit Drugs and Alcohol	0.5769		0.8390	
Illicit Drugs or Alcohol	0.1056		0.6555	
Illicit Drugs	0.1359		0.9569	
Alcohol	0.2801		0.6932	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16C (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2010 and 2011

Dependence or Abuse Status	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
DEPENDENCE												
Both Illicit Drugs and Alcohol	86	96	73	87	46	63	31	37	44	44	47	39
Illicit Drugs or Alcohol	361	286	247	206	138	122	128	87	157	126	254	211
Illicit Drugs	201	170	150	127	96	86	86	60	74	75	125	113
Alcohol	302	257	208	182	108	107	102	79	144	111	218	187
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	125	120	98	99	63	66	40	47	63	55	79	71
Illicit Drugs or Alcohol	460	401	287	251	155	145	144	107	183	164	385	349
Illicit Drugs	237	195	176	136	104	91	94	65	100	83	147	138
Alcohol	409	374	251	229	130	130	123	96	168	152	358	331

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.16D (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2010 and 2011

Dependence or Abuse Status	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.04	0.04	0.17	0.21	0.48	0.68	0.32	0.36	0.19	0.20	0.03	0.02
Illicit Drugs or Alcohol	0.16	0.12	0.56	0.48	1.33	1.24	1.21	0.83	0.68	0.56	0.13	0.11
Illicit Drugs	0.09	0.07	0.35	0.30	0.98	0.91	0.84	0.58	0.33	0.34	0.07	0.06
Alcohol	0.13	0.11	0.48	0.42	1.08	1.11	1.00	0.76	0.62	0.49	0.11	0.10
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.05	0.05	0.24	0.24	0.65	0.71	0.41	0.45	0.28	0.25	0.04	0.04
Illicit Drugs or Alcohol	0.20	0.17	0.64	0.57	1.47	1.40	1.34	1.01	0.78	0.72	0.19	0.17
Illicit Drugs	0.10	0.08	0.42	0.33	1.05	0.95	0.93	0.63	0.45	0.38	0.08	0.07
Alcohol	0.18	0.16	0.56	0.52	1.27	1.31	1.18	0.91	0.72	0.67	0.18	0.17

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 1.16P (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness:
P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Dependence or Abuse Status	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.9616		0.6757		0.2889		0.8967		0.7101		0.4713	
Illicit Drugs or Alcohol	0.0291		0.0111		0.3459		0.0051		0.3522		0.5924	
Illicit Drugs	0.1756		0.3031		0.7992		0.2210		0.7482		0.4444	
Alcohol	0.1005		0.0471		0.7738		0.0185		0.3360		0.7504	
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.1911		0.2724		0.9180		0.5902		0.2044		0.5554	
Illicit Drugs or Alcohol	0.0023		0.0025		0.3061		0.0009		0.2295		0.1442	
Illicit Drugs	0.0218		0.0188		0.6010		0.0701		0.0707		0.4497	
Alcohol	0.0117		0.0263		0.4389		0.0100		0.4737		0.1668	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17C (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	237	195	176	136	104	91	94	65	100	83	147	138
18-25	85	83	57	51	31	27	29	23	35	36	69	72
26-49	178	145	135	103	80	74	55	53	89	55	115	98
50 or Older	114	93	94	71	54	43	70	30	28	48	64	62
MALE	187	161	133	100	66	54	78	50	86	67	130	124
18-25	65	64	41	35	19	17	22	18	28	27	58	61
26-49	142	121	104	76	60	48	37	40	76	47	98	92
50 or Older	96	76	72	51	*	*	*	*	24	39	64	58
FEMALE	131	107	111	92	81	72	53	39	50	48	63	54
18-25	51	47	38	37	25	23	18	16	21	23	36	32
26-49	98	80	80	71	53	57	40	32	43	32	53	37
50 or Older	58	54	58	49	50	38	26	*	15	27	*	22

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17D (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.10	0.08	0.42	0.33	1.05	0.95	0.93	0.63	0.45	0.38	0.08	0.07
18-25	0.25	0.24	0.80	0.71	1.96	1.86	1.54	1.32	1.00	0.95	0.24	0.25
26-49	0.18	0.15	0.62	0.51	1.45	1.39	1.11	0.99	0.79	0.57	0.15	0.13
50 or Older	0.12	0.09	0.64	0.46	1.83	1.52	2.04	0.86	0.34	0.53	0.08	0.07
MALE	0.17	0.14	0.79	0.61	1.85	1.64	2.03	1.38	0.92	0.73	0.14	0.13
18-25	0.38	0.37	1.47	1.26	3.65	3.45	3.07	2.81	1.90	1.60	0.37	0.39
26-49	0.29	0.25	1.17	0.96	2.81	2.59	2.07	2.09	1.55	1.16	0.24	0.23
50 or Older	0.21	0.16	1.21	0.85	*	*	*	*	0.68	1.07	0.16	0.14
FEMALE	0.11	0.09	0.43	0.36	1.28	1.17	0.84	0.59	0.39	0.37	0.07	0.06
18-25	0.30	0.28	0.90	0.87	2.31	2.35	1.61	1.40	1.07	1.12	0.27	0.23
26-49	0.20	0.16	0.62	0.58	1.55	1.66	1.24	0.93	0.70	0.55	0.14	0.10
50 or Older	0.11	0.10	0.66	0.53	2.59	2.20	1.29	*	0.30	0.52	*	0.05

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.17P (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.0218		0.0188		0.6010		0.0701		0.0707		0.4497	
18-25	0.3019		0.0041		0.8449		0.0249		0.0335		0.6088	
26-49	0.0222		0.0540		0.5311		0.5473		0.0350		0.2449	
50 or Older	0.9448		0.8724		0.9864		0.3296		0.2928		0.9042	
MALE	0.2123		0.1080		0.8672		0.4071		0.1123		0.9787	
18-25	0.4502		0.0176		0.4441		0.0192		0.0541		0.0546	
26-49	0.1032		0.1667		0.5998		0.7193		0.0691		0.5261	
50 or Older	0.8290		0.9521		*		*		0.5637		0.7348	
FEMALE	0.0228		0.0967		0.5996		0.0797		0.4728		0.0756	
18-25	0.0095		0.0711		0.3710		0.4733		0.1828		0.0474	
26-49	0.1044		0.2719		0.8269		0.2222		0.4445		0.1516	
50 or Older	0.8379		0.7517		0.7829		*		0.2931		*	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18C (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	409	374	251	229	130	130	123	96	168	152	358	331
18-25	125	116	68	72	36	31	35	35	41	50	121	120
26-49	280	259	199	175	110	100	98	75	135	122	228	229
50 or Older	262	238	134	129	59	78	66	56	100	78	232	203
MALE	332	300	177	165	94	82	85	80	123	117	312	271
18-25	99	83	49	48	24	20	26	22	31	37	101	85
26-49	232	214	150	122	78	59	72	54	106	94	202	199
50 or Older	223	194	95	97	*	*	*	*	76	62	208	168
FEMALE	225	219	174	153	90	101	94	60	109	96	150	166
18-25	70	79	47	52	28	23	23	28	28	33	58	73
26-49	149	160	132	124	78	80	72	52	80	78	93	106
50 or Older	140	127	94	76	41	58	55	14	65	48	102	102

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18D (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.18	0.16	0.56	0.52	1.27	1.31	1.18	0.91	0.72	0.67	0.18	0.17
18-25	0.37	0.34	0.90	0.91	2.15	1.96	1.79	1.84	1.15	1.23	0.37	0.35
26-49	0.28	0.26	0.85	0.80	1.89	1.77	1.78	1.35	1.15	1.16	0.27	0.28
50 or Older	0.27	0.24	0.90	0.84	2.05	2.70	1.96	1.58	1.16	0.87	0.28	0.23
MALE	0.30	0.27	1.01	0.97	2.52	2.35	2.19	2.08	1.26	1.22	0.31	0.27
18-25	0.57	0.49	1.68	1.56	4.02	3.67	3.22	3.22	2.07	2.03	0.58	0.50
26-49	0.48	0.44	1.54	1.45	3.42	3.09	3.50	2.76	2.02	2.11	0.46	0.46
50 or Older	0.50	0.42	1.58	1.58	*	*	*	*	2.11	1.67	0.52	0.41
FEMALE	0.19	0.18	0.67	0.57	1.41	1.57	1.41	0.88	0.84	0.73	0.16	0.17
18-25	0.41	0.46	1.07	1.14	2.52	2.34	1.96	2.19	1.39	1.53	0.42	0.48
26-49	0.30	0.32	1.01	0.94	2.18	2.19	2.09	1.47	1.30	1.27	0.24	0.28
50 or Older	0.27	0.24	1.05	0.82	2.15	3.23	2.71	0.65	1.30	0.90	0.24	0.23

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.18P (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.0117		0.0263		0.4389		0.0100		0.4737		0.1668	
18-25	0.0092		0.1675		0.3087		0.3521		0.7457		0.0114	
26-49	0.2325		0.0184		0.0741		0.0202		0.8462		0.6151	
50 or Older	0.3028		0.8629		0.1176		0.3185		0.5021		0.2723	
MALE	0.0039		0.1049		0.3196		0.3175		0.3672		0.0295	
18-25	0.0013		0.1647		0.4868		0.1716		0.6482		0.0011	
26-49	0.1880		0.0571		0.1247		0.0549		0.8019		0.9724	
50 or Older	0.1967		0.8496		*		*		0.3459		0.1253	
FEMALE	0.9071		0.2113		0.9035		0.0128		0.8992		0.2752	
18-25	0.8635		0.5103		0.4286		0.9728		0.8102		0.7474	
26-49	0.9948		0.3465		0.3561		0.2018		0.7585		0.2551	
50 or Older	0.9286		0.6566		0.1623		0.0114		0.9134		0.6268	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.19C (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	460	401	287	251	155	145	144	107	183	164	385	349
18-25	139	128	80	85	43	38	40	39	49	59	134	135
26-49	310	281	225	194	126	111	107	84	150	130	247	239
50 or Older	280	251	157	139	76	85	91	59	103	86	239	214
MALE	362	321	205	182	104	88	111	86	137	130	335	287
18-25	106	92	56	56	26	23	30	24	37	44	111	98
26-49	250	232	170	138	90	66	77	61	121	104	214	208
50 or Older	238	204	116	102	*	*	*	*	78	67	216	180
FEMALE	251	230	197	166	116	114	106	67	113	101	158	172
18-25	78	83	57	59	34	29	26	31	34	36	64	79
26-49	167	171	146	133	90	89	81	56	83	80	101	111
50 or Older	152	135	110	87	62	66	61	21	67	54	102	105

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.19D (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2010 and 2011

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.20	0.17	0.64	0.57	1.47	1.40	1.34	1.01	0.78	0.72	0.19	0.17
18-25	0.41	0.37	0.97	1.00	2.29	2.20	1.94	1.92	1.28	1.35	0.42	0.38
26-49	0.31	0.29	0.93	0.88	2.03	1.89	1.91	1.50	1.23	1.23	0.29	0.29
50 or Older	0.29	0.25	1.04	0.90	2.55	2.91	2.59	1.68	1.18	0.94	0.29	0.25
MALE	0.33	0.29	1.15	1.05	2.69	2.52	2.71	2.24	1.41	1.35	0.33	0.29
18-25	0.62	0.54	1.73	1.65	4.03	3.99	3.37	3.36	2.21	2.23	0.63	0.56
26-49	0.51	0.48	1.66	1.59	3.60	3.28	3.65	3.03	2.23	2.28	0.49	0.48
50 or Older	0.53	0.44	1.90	1.65	*	*	*	*	2.15	1.79	0.54	0.44
FEMALE	0.21	0.19	0.74	0.62	1.74	1.72	1.58	1.00	0.87	0.78	0.17	0.18
18-25	0.47	0.48	1.21	1.20	2.78	2.59	2.17	2.34	1.62	1.62	0.45	0.50
26-49	0.33	0.34	1.09	1.01	2.40	2.39	2.34	1.58	1.33	1.31	0.26	0.29
50 or Older	0.29	0.25	1.21	0.93	3.18	3.63	2.97	0.95	1.33	1.00	0.24	0.23

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.19P (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Gender/ Age Group	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
TOTAL	0.0023		0.0025		0.3061		0.0009		0.2295		0.1442	
18-25	0.0104		0.0245		0.5488		0.1016		0.1834		0.0417	
26-49	0.0677		0.0034		0.0288		0.0185		0.4683		0.8686	
50 or Older	0.2204		0.5765		0.1820		0.0650		0.6704		0.2790	
MALE	0.0014		0.0334		0.4204		0.1773		0.1415		0.0272	
18-25	0.0056		0.0508		0.9944		0.0213		0.3265		0.0111	
26-49	0.0792		0.0233		0.0899		0.1702		0.3082		0.7966	
50 or Older	0.1286		0.9115		*		*		0.4190		0.1009	
FEMALE	0.5197		0.0609		0.5382		0.0023		0.8999		0.2989	
18-25	0.4906		0.1511		0.4252		0.8325		0.2269		0.8100	
26-49	0.6351		0.1948		0.2000		0.0690		0.5963		0.4046	
50 or Older	0.9476		0.4831		0.4123		0.0064		0.7450		0.4517	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.