

Table 1.31C (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2010 and 2011

Location of Treatment/Counseling ¹	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
Outpatient Mental Health Clinic or Center	197	221	179	189	116	125	90	117	103	91	98	108
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	357	330	267	242	188	148	121	120	144	164	212	201
Doctor's Office - Not Part of a Clinic	208	186	160	164	101	109	100	88	73	82	139	104
Outpatient Medical Clinic	110	129	88	92	69	60	37	52	41	47	65	91
Partial Day Hospital or Day Treatment Program	53	59	46	54	39	44	17	19	20	25	26	23
School or University Setting/Clinic/Center ²	16	14	11	13	6	7	7	7	6	8	11	8
Some Other Place ³	67	63	58	52	16	45	48	17	29	20	34	34

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.31D (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2010 and 2011

Location of Treatment/Counseling ¹	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
Outpatient Mental Health Clinic or Center	1.18	1.26	1.61	1.67	2.58	2.68	2.93	3.52	2.96	2.45	1.69	1.80
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	1.46	1.39	1.84	1.67	2.79	2.63	3.44	3.49	3.31	3.07	2.48	2.25
Doctor's Office - Not Part of a Clinic	1.21	1.08	1.46	1.44	2.33	2.33	3.09	2.93	2.31	2.23	2.27	1.71
Outpatient Medical Clinic	0.71	0.79	0.87	0.90	1.66	1.47	1.35	1.87	1.34	1.41	1.16	1.55
Partial Day Hospital or Day Treatment Program	0.35	0.37	0.48	0.54	0.97	1.09	0.63	0.70	0.66	0.77	0.48	0.41
School or University Setting/Clinic/Center ²	0.10	0.10	0.11	0.13	0.15	0.17	0.26	0.28	0.19	0.24	0.21	0.15
Some Other Place ³	0.44	0.41	0.60	0.53	0.41	1.16	1.74	0.63	0.94	0.63	0.61	0.62

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.31P (REVISED) Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Past Year Level of Mental Illness: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Location of Treatment/Counseling ¹	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
Outpatient Mental Health Clinic or Center	0.3349		0.4212		0.5610		0.5870		0.7158		0.6094	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.7457		0.5913		0.1375		0.9639		0.4991		0.8677	
Doctor's Office - Not Part of a Clinic	0.3907		0.9064		0.4292		0.4651		0.7521		0.2281	
Outpatient Medical Clinic	0.4210		0.5914		0.9845		0.4818		0.7085		0.5355	
Partial Day Hospital or Day Treatment Program	0.9525		0.9232		0.9365		0.8447		0.8595		0.9665	
School or University Setting/Clinic/Center ²	0.9381		0.5869		0.9319		0.9133		0.3285		0.6129	
Some Other Place ³	0.6398		0.7004		0.0191		0.2579		0.3858		0.8020	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.