

**Table 1.65C Had Serious Thoughts of Suicide in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	299	299	311	299
<b>AGE</b>				
18-25	74	66	75	85
26-49	193	200	201	169
50 or Older	210	215	214	229
<b>GENDER</b>				
Male	206	204	235	194
Female	210	211	216	228
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	282	280	297	288
White	265	256	269	267
Black or African American	88	89	103	86
American Indian or Alaska Native	9	13	24	55
Native Hawaiian or Other Pacific Islander	18	*	14	*
Asian	44	38	72	69
Two or More Races	38	48	24	25
Hispanic or Latino	92	112	76	84

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Respondents with unknown suicide information were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 1.65D Had Serious Thoughts of Suicide in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	0.13	0.13	0.14	0.13
<b>AGE</b>				
18-25	0.23	0.20	0.22	0.25
26-49	0.19	0.20	0.20	0.17
50 or Older	0.23	0.23	0.22	0.23
<b>GENDER</b>				
Male	0.19	0.19	0.21	0.17
Female	0.18	0.18	0.18	0.19
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	0.15	0.14	0.15	0.15
White	0.16	0.16	0.17	0.17
Black or African American	0.34	0.32	0.38	0.32
American Indian or Alaska Native	0.92	1.24	2.17	3.72
Native Hawaiian or Other Pacific Islander	2.21	*	2.13	*
Asian	0.46	0.38	0.69	0.61
Two or More Races	1.48	1.87	0.96	0.78
Hispanic or Latino	0.30	0.36	0.24	0.25

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Respondents with unknown suicide information were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 1.65P Had Serious Thoughts of Suicide in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2008-2010**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	0.8498	0.7810	0.4811	
<b>AGE</b>				
18-25	0.9255	0.0164	0.6098	
26-49	0.2435	0.0141	0.1292	
50 or Older	0.3816	0.2929	0.8211	
<b>GENDER</b>				
Male	0.7116	0.5291	0.1693	
Female	0.9349	0.8546	0.7095	
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	0.9243	0.6774	0.4363	
White	0.9047	0.6556	0.7304	
Black or African American	0.5623	0.6631	0.1315	
American Indian or Alaska Native	0.0080	0.0324	0.1803	
Native Hawaiian or Other Pacific Islander	*	*	*	
Asian	0.4106	0.1964	0.7159	
Two or More Races	0.4450	0.1081	0.3948	
Hispanic or Latino	0.5135	0.0559	0.7160	

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Respondents with unknown suicide information were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.