

**Table 1.67C Serious Psychological Distress (SPD) in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	347	324	363	341
<b>AGE</b>				
18-25	76	75	84	80
26-49	212	212	217	215
50 or Older	245	230	272	245
<b>GENDER</b>				
Male	234	194	235	227
Female	251	259	280	251
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	319	306	320	309
White	286	293	288	267
Black or African American	122	105	137	114
American Indian or Alaska Native	10	8	11	36
Native Hawaiian or Other Pacific Islander	*	14	21	*
Asian	74	35	53	62
Two or More Races	51	33	69	45
Hispanic or Latino	140	130	163	139

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 1.67D Serious Psychological Distress (SPD) in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2008-2011**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	0.15	0.14	0.16	0.15
<b>AGE</b>				
18-25	0.23	0.22	0.25	0.23
26-49	0.21	0.21	0.22	0.22
50 or Older	0.27	0.24	0.28	0.24
<b>GENDER</b>				
Male	0.22	0.18	0.21	0.20
Female	0.22	0.22	0.24	0.21
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	0.16	0.16	0.16	0.16
White	0.18	0.18	0.18	0.17
Black or African American	0.45	0.38	0.48	0.41
American Indian or Alaska Native	1.12	0.79	1.07	2.61
Native Hawaiian or Other Pacific Islander	*	1.98	3.13	*
Asian	0.74	0.35	0.50	0.55
Two or More Races	1.94	1.32	2.46	1.35
Hispanic or Latino	0.46	0.42	0.51	0.41

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 1.67P Serious Psychological Distress (SPD) in the Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2011 Versus 2008-2010**

<b>Demographic Characteristic</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>
<b>TOTAL</b>	0.4796	0.7828	0.5612	
<b>AGE</b>				
18-25	0.8268	0.9569	0.5033	
26-49	0.8038	0.7518	0.7764	
50 or Older	0.3330	0.3682	0.4910	
<b>GENDER</b>				
Male	0.7486	0.1939	0.7292	
Female	0.5138	0.4731	0.2723	
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic or Latino	0.1873	0.4607	0.6954	
White	0.2954	0.6753	0.9058	
Black or African American	0.5884	0.1337	0.8923	
American Indian or Alaska Native	0.2426	0.1598	0.1625	
Native Hawaiian or Other Pacific Islander	*	*	*	
Asian	0.8288	0.1964	0.7056	
Two or More Races	0.9713	0.5881	0.9789	
Hispanic or Latino	0.2393	0.3527	0.0304	

\*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.