

**Table 2.13A Had at Least One Major Depressive Episode (MDE) in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2004-2011**

Demographic Characteristic	2004	2005	2006	2007	2008	2009	2010	2011
<b>TOTAL</b>	2,225 <sup>a</sup>	2,191 <sup>a</sup>	1,970	2,016	2,027	1,954	1,911	2,011
<b>AGE</b>								
12-13	445 <sup>b</sup>	417 <sup>b</sup>	383	337	366	330	330	312
14-15	783	811	684	705	706	741	706	710
16-17	997	964	902	974	955	883	876	989
<b>GENDER</b>								
Male	637	571	539	586	540	577	536	566
Female	1,588 <sup>a</sup>	1,620 <sup>a</sup>	1,431	1,430	1,487	1,377	1,375	1,446
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	1,848 <sup>b</sup>	1,802 <sup>b</sup>	1,614	1,691	1,681	1,594	1,537	1,586
White	1,413 <sup>b</sup>	1,390 <sup>b</sup>	1,220	1,284	1,266	1,180	1,186	1,167
Black or African American	287	288	245	292	261	285	236	238
American Indian or Alaska Native	13	10	12	7	13	9	10	15
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	1	*
Asian	82	63	77	66	73	73	57	90
Two or More Races	51	41 <sup>a</sup>	52	41 <sup>a</sup>	59	41 <sup>a</sup>	48	75
Hispanic or Latino	377	389	356	325 <sup>a</sup>	347	360	374	425

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

Respondents with unknown past year MDE data were excluded.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2011.

**Table 2.13B Had at Least One Major Depressive Episode (MDE) in the Past Year among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2004-2011**

Demographic Characteristic	2004	2005	2006	2007	2008	2009	2010	2011
<b>TOTAL</b>	9.0 <sup>a</sup>	8.8	7.9	8.2	8.3	8.1	8.0	8.2
<b>AGE</b>								
12-13	5.4 <sup>b</sup>	5.2 <sup>a</sup>	4.9	4.3	4.9	4.6	4.3	4.1
14-15	9.2	9.5	7.9	8.4	8.5	8.8	9.0	8.6
16-17	12.3	11.5	10.7	11.5	11.2	10.4	10.6	11.7
<b>GENDER</b>								
Male	5.0	4.5	4.2	4.6	4.3	4.7	4.4	4.5
Female	13.1	13.3 <sup>a</sup>	11.8	11.9	12.5	11.7	11.9	12.1
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	8.9	8.7	7.9	8.4	8.5	8.2	8.1	8.3
White	9.2	9.1	8.2	8.7	8.8	8.4	8.6	8.6
Black or African American	7.7	7.6	6.4	7.8	7.1	7.9	6.8	7.0
American Indian or Alaska Native	7.8	6.1	9.3	4.6	10.1	7.5	7.4	11.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	1.8	*
Asian	8.3	6.0	7.7	6.6	7.7	7.6	5.5	7.6
Two or More Races	11.7	10.5	13.0	9.9	12.0	8.0	9.4	10.6
Hispanic or Latino	9.1	9.1	8.0	7.1	7.5	7.7	7.8	8.1

\*Low precision; no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

Respondents with unknown past year MDE data were excluded.

<sup>a</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2011.