

Table 2.9A Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Numbers in Thousands, 2010 and 2011

Substance	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
PAST YEAR USE						
Illicit Drugs ²	4,745	4,735	709	724	3,902	3,893
Marijuana and Hashish	3,414	3,553	481	523	2,836	2,962
Cocaine	239	229	46	54	178	165
Crack	18	29	9	11	9	18
Heroin	28	54	10	11	18	36
Hallucinogens	747	648	147	122	570	499
LSD	136	159	24	30	103	113
PCP	35	39	8	12	26	26
Ecstasy	465	419	97	82	353	320
Inhalants	885	820	168	174	683	612
Nonmedical Use of Psychotherapeutics ^{3,4}	1,812	1,749	325	327	1,431	1,364
Pain Relievers	1,522	1,484	264	275	1,210	1,157
OxyContin [®]	235	219	33	54	187	159
Tranquilizers	466	475	103	124	345	345
Stimulants ⁴	325	299	76	79	234	210
Methamphetamine ⁴	98	106	20	28	73	73
Sedatives	111	84	31	20	72	62
Illicit Drugs Other Than Marijuana ²	2,758	2,565	498	476	2,182	2,001
PAST MONTH USE						
Daily Cigarette Use ⁵	459	440	59	81	388	349
Heavy Alcohol Use ⁶	409	379	48	50	349	322

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9B Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Percentages, 2010 and 2011

Substance	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
PAST YEAR USE						
Illicit Drugs ²	19.5	19.0	37.1	36.0	17.8	17.4
Marijuana and Hashish	14.0	14.2	25.2	26.0	12.9	13.2
Cocaine	1.0	0.9	2.4	2.7	0.8	0.7
Crack	0.1	0.1	0.5	0.5	0.0	0.1
Heroin	0.1	0.2	0.5	0.5	0.1	0.2
Hallucinogens	3.1 ^a	2.6	7.7	6.1	2.6	2.2
LSD	0.6	0.6	1.2	1.5	0.5	0.5
PCP	0.1	0.2	0.4	0.6	0.1	0.1
Ecstasy	1.9	1.7	5.1	4.1	1.6	1.4
Inhalants	3.6	3.3	8.8	8.6	3.1	2.7
Nonmedical Use of Psychotherapeutics ^{3,4}	7.4	7.0	17.0	16.3	6.5	6.1
Pain Relievers	6.3	5.9	13.8	13.7	5.5	5.2
OxyContin [®]	1.0	0.9	1.7	2.7	0.9	0.7
Tranquilizers	1.9	1.9	5.4	6.1	1.6	1.5
Stimulants ⁴	1.3	1.2	4.0	3.9	1.1	0.9
Methamphetamine ⁴	0.4	0.4	1.1	1.4	0.3	0.3
Sedatives	0.5	0.3	1.6	1.0	0.3	0.3
Illicit Drugs Other Than Marijuana ²	11.3 ^a	10.3	26.0	23.7	10.0 ^a	8.9
PAST MONTH USE						
Daily Cigarette Use ⁵	1.9	1.8	3.1	4.0	1.8	1.6
Heavy Alcohol Use ⁶	1.7	1.5	2.5	2.5	1.6	1.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.