

Table 2.9C Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
PAST YEAR USE						
Illicit Drugs ²	94	93	34	38	102	94
Marijuana and Hashish	84	83	29	32	87	81
Cocaine	22	20	9	9	20	16
Crack	5	7	4	5	3	5
Heroin	8	11	4	4	7	9
Hallucinogens	40	35	16	14	37	31
LSD	17	18	6	7	16	14
PCP	8	7	3	5	7	6
Ecstasy	32	29	14	12	29	25
Inhalants	42	40	17	17	40	35
Nonmedical Use of Psychotherapeutics ^{3,4}	58	53	22	24	57	50
Pain Relievers	54	48	20	22	52	45
OxyContin [®]	24	18	7	10	21	16
Tranquilizers	30	31	12	14	27	28
Stimulants ⁴	24	23	11	11	21	19
Methamphetamine ⁴	15	16	6	7	13	13
Sedatives	14	12	7	5	11	11
Illicit Drugs Other Than Marijuana ²	74	65	29	30	74	64
PAST MONTH USE						
Daily Cigarette Use ⁵	30	28	9	11	27	25
Heavy Alcohol Use ⁶	28	25	9	8	26	23

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9D Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2010 and 2011

Substance	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
PAST YEAR USE						
Illicit Drugs ²	0.38	0.37	1.47	1.51	0.40	0.38
Marijuana and Hashish	0.34	0.33	1.35	1.37	0.36	0.34
Cocaine	0.09	0.08	0.45	0.45	0.09	0.07
Crack	0.02	0.03	0.19	0.23	0.01	0.02
Heroin	0.03	0.05	0.20	0.19	0.03	0.04
Hallucinogens	0.16	0.14	0.81	0.68	0.16	0.13
LSD	0.07	0.07	0.29	0.33	0.07	0.06
PCP	0.03	0.03	0.18	0.24	0.03	0.02
Ecstasy	0.13	0.12	0.72	0.58	0.13	0.11
Inhalants	0.17	0.16	0.85	0.80	0.18	0.15
Nonmedical Use of Psychotherapeutics ^{3,4}	0.24	0.21	1.11	1.09	0.24	0.21
Pain Relievers	0.22	0.19	1.01	1.02	0.22	0.19
OxyContin [®]	0.10	0.07	0.38	0.48	0.09	0.07
Tranquilizers	0.12	0.12	0.64	0.70	0.12	0.12
Stimulants ⁴	0.10	0.09	0.56	0.55	0.09	0.08
Methamphetamine ⁴	0.06	0.06	0.31	0.37	0.06	0.06
Sedatives	0.06	0.05	0.39	0.26	0.05	0.05
Illicit Drugs Other Than Marijuana ²	0.30	0.26	1.36	1.29	0.31	0.26
PAST MONTH USE						
Daily Cigarette Use ⁵	0.12	0.11	0.45	0.53	0.12	0.11
Heavy Alcohol Use ⁶	0.11	0.10	0.44	0.41	0.12	0.10

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 2.9P Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	Total ¹ (2010)	Total ¹ (2011)	MDE (2010)	MDE (2011)	No MDE (2010)	No MDE (2011)
PAST YEAR USE						
Illicit Drugs ²	0.3277		0.6056		0.4405	
Marijuana and Hashish	0.6664		0.6723		0.5570	
Cocaine	0.5837		0.6264		0.5358	
Crack	0.2407		0.8119		0.1597	
Heroin	0.0731		0.8992		0.1271	
Hallucinogens	0.0301		0.1272		0.0842	
LSD	0.4254		0.5678		0.7270	
PCP	0.7631		0.6283		0.9944	
Ecstasy	0.1975		0.2809		0.2913	
Inhalants	0.1489		0.9071		0.1159	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.1571		0.6424		0.1582	
Pain Relievers	0.2855		0.9273		0.2215	
OxyContin [®]	0.4648		0.1196		0.2238	
Tranquilizers	0.9260		0.4339		0.8289	
Stimulants ⁴	0.3223		0.9427		0.3051	
Methamphetamine ⁴	0.8063		0.4919		0.9054	
Sedatives	0.1090		0.2083		0.4690	
Illicit Drugs Other Than Marijuana ²	0.0100		0.2178		0.0122	
PAST MONTH USE						
Daily Cigarette Use ⁵	0.4622		0.1906		0.1897	
Heavy Alcohol Use ⁶	0.2621		0.9676		0.2985	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.