

Table 1.13A (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2009 and 2010

Drug	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
ILLCIT DRUGS¹	33,285	34,163	10,222	10,163	2,684	3,074	2,609	2,485	4,928	4,603	23,063	24,001
Marijuana and Hashish	25,325	25,887	7,329	7,198	1,840	2,080	1,936	1,811	3,553	3,307	17,996	18,689
Cocaine	4,564	4,294	1,682	1,512	505	387	345	402	832	723	2,882	2,782
Crack	979	866	523	469	203	158	104	140	217	170	456	397
Heroin	547	593	266	292	140	137	51	45	74	110	282	301
Hallucinogens	3,799	3,795	1,297	1,256	294	397	374	364	629	495	2,502	2,540
LSD	647	746	227	287	61	88	70	115	96	84	420	459
PCP	77	61	32	34	3	24	23	3	6	6	45	27
Ecstasy	2,388	2,188	852	751	182	207	238	216	432	328	1,536	1,436
Inhalants	1,149	1,163	452	415	89	141	171	112	193	162	697	748
Nonmedical Use of Psychotherapeutics ^{2,3}	14,169	14,238	5,350	5,333	1,556	1,747	1,323	1,308	2,471	2,278	8,819	8,905
Pain Relievers	10,822	10,720	4,076	4,072	1,262	1,440	1,056	1,002	1,758	1,629	6,747	6,648
OxyContin [®]	1,460	1,645	672	783	281	384	133	195	258	204	788	862
Tranquilizers	5,031	5,117	2,298	2,275	649	691	568	630	1,081	955	2,733	2,841
Stimulants ³	2,759	2,577	1,260	1,061	390	320	323	255	547	486	1,499	1,517
Methamphetamine ³	1,080	861	570 ^a	367	249 ^a	118	130	74	192	176	510	494
Sedatives	690	796	325	394	142	187	70	77	113	131	365	401
ILLCIT DRUGS OTHER THAN MARIJUANA¹	18,228	17,866	6,523	6,427	1,840	2,058	1,620	1,619	3,063	2,750	11,705	11,440

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.13B (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2009 and 2010

Drug	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
ILLCIT DRUGS¹	14.6	14.9	24.8	24.5	32.0	32.9	25.6	25.2	21.8	20.7	12.4	12.8
Marijuana and Hashish	11.1	11.3	17.8	17.4	21.9	22.3	19.0	18.4	15.7	14.9	9.7	9.9
Cocaine	2.0	1.9	4.1	3.7	6.0	4.1	3.4	4.1	3.7	3.3	1.5	1.5
Crack	0.4	0.4	1.3	1.1	2.4	1.7	1.0	1.4	1.0	0.8	0.2	0.2
Heroin	0.2	0.3	0.6	0.7	1.7	1.5	0.5	0.5	0.3	0.5	0.2	0.2
Hallucinogens	1.7	1.7	3.1	3.0	3.5	4.2	3.7	3.7	2.8	2.2	1.3	1.4
LSD	0.3	0.3	0.6	0.7	0.7	0.9	0.7	1.2	0.4	0.4	0.2	0.2
PCP	0.0	0.0	0.1	0.1	0.0	0.3	0.2	0.0	0.0	0.0	0.0	0.0
Ecstasy	1.1	1.0	2.1	1.8	2.2	2.2	2.3	2.2	1.9	1.5	0.8	0.8
Inhalants	0.5	0.5	1.1	1.0	1.1	1.5	1.7	1.1	0.9	0.7	0.4	0.4
Nonmedical Use of Psychotherapeutics ^{2,3}	6.2	6.2	13.0	12.9	18.6	18.7	13.0	13.3	10.9	10.3	4.7	4.7
Pain Relievers	4.8	4.7	9.9	9.8	15.1	15.4	10.4	10.2	7.8	7.3	3.6	3.5
OxyContin [®]	0.6	0.7	1.6	1.9	3.3	4.1	1.3	2.0	1.1	0.9	0.4	0.5
Tranquilizers	2.2	2.2	5.6	5.5	7.7	7.4	5.6	6.4	4.8	4.3	1.5	1.5
Stimulants ³	1.2	1.1	3.1	2.6	4.7	3.4	3.2	2.6	2.4	2.2	0.8	0.8
Methamphetamine ³	0.5	0.4	1.4 ^a	0.9	3.0 ^a	1.3	1.3	0.7	0.8	0.8	0.3	0.3
Sedatives	0.3	0.3	0.8	1.0	1.7	2.0	0.7	0.8	0.5	0.6	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	8.0	7.8	15.8	15.5	21.9	22.0	15.9	16.4	13.5	12.4	6.3	6.1

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.14A (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2009 and 2010

Substance	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
PAST YEAR												
Tobacco Products ¹	78,716	78,965	18,713	18,937	4,298 ^a	4,944	5,032	4,662	9,383	9,330	60,002	60,029
Cigarettes	65,537	65,139	16,769	16,722	3,996 ^a	4,573	4,522	4,105	8,250	8,044	48,768	48,417
Smokeless Tobacco	10,870	11,169	2,129	2,151	521	477	509	504	1,099	1,170	8,741	9,018
Cigars	24,374	24,179	5,275	5,246	1,096	1,376	1,466	1,248	2,713	2,622	19,099	18,933
Alcohol	160,620	161,399	29,515	29,515	6,013	6,591	7,605	7,072	15,897	15,852	131,105	131,884
PAST MONTH												
Tobacco Products ¹	66,932	67,051	16,397	16,359	3,905	4,292	4,406	4,091	8,086	7,976	50,534	50,692
Cigarettes	56,507	56,295	14,845	14,593	3,656	4,015	3,992	3,655	7,197	6,923	41,661	41,702
Smokeless Tobacco	8,032	8,309	1,414	1,492	374	316	337	324	703	852	6,618	6,817
Cigars	12,355	12,485	2,729	2,864	577	721	845	699	1,307	1,444	9,627	9,621
Pipe Tobacco	1,873	2,000	450	646	66 ^b	190	227	128	157 ^a	328	1,423	1,354
Alcohol	126,994	128,055	22,860	22,829	4,353 ^a	5,000	5,792	5,345	12,715	12,484	104,134	105,226
Binge Alcohol Use ²	57,501	56,623	11,673	11,155	2,277	2,594	2,971	2,657	6,425	5,904	45,828	45,467
Heavy Alcohol Use ²	16,655	16,490	3,653	3,523	712	885	951	879	1,990	1,760	13,002	12,967

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.14B (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2009 and 2010

Substance	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
PAST YEAR												
Tobacco Products ¹	34.6	34.4	45.4	45.7	51.3	53.0	49.4	47.2	41.5	42.0	32.3	32.0
Cigarettes	28.8	28.4	40.7	40.4	47.7	49.0	44.4	41.6	36.4	36.2	26.2	25.8
Smokeless Tobacco	4.8	4.9	5.2	5.2	6.2	5.1	5.0	5.1	4.9	5.3	4.7	4.8
Cigars	10.7	10.5	12.8	12.7	13.1	14.7	14.4	12.6	12.0	11.8	10.3	10.1
Alcohol	70.7	70.4	71.6	71.3	71.7	70.6	74.7	71.7	70.2	71.4	70.5	70.2
PAST MONTH												
Tobacco Products ¹	29.5	29.2	39.8	39.5	46.6	46.0	43.3	41.5	35.7	35.9	27.2	27.0
Cigarettes	24.9	24.6	36.0	35.2	43.6	43.0	39.2	37.0	31.8	31.2	22.4	22.2
Smokeless Tobacco	3.5	3.6	3.4	3.6	4.5	3.4	3.3	3.3	3.1	3.8	3.6	3.6
Cigars	5.4	5.4	6.6	6.9	6.9	7.7	8.3	7.1	5.8	6.5	5.2	5.1
Pipe Tobacco	0.8	0.9	1.1	1.6	0.8 ^b	2.0	2.2	1.3	0.7 ^a	1.5	0.8	0.7
Alcohol	55.9	55.9	55.5	55.1	51.9	53.6	56.9	54.2	56.2	56.2	56.0	56.0
Binge Alcohol Use ²	25.3	24.7	28.3	26.9	27.2	27.8	29.2	26.9	28.4	26.6	24.6	24.2
Heavy Alcohol Use ²	7.3	7.2	8.9	8.5	8.5	9.5	9.3	8.9	8.8	7.9	7.0	6.9

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.