

Table 1.15A (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2009 and 2010

Dependence or Abuse Status	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)
DEPENDENCE				
None	35,810	36,003	6,804	7,508
Both Illicit Drugs and Alcohol	789	648	249	277
Illicit Drugs or Alcohol	5,386	5,414	1,579	1,826
Illicit Drugs	2,214	2,230	803	869
Alcohol	3,961	3,833	1,024	1,234
DEPENDENCE OR ABUSE				
None	33,373	33,552	6,323	6,946
Both Illicit Drugs and Alcohol	1,551	1,324	511	503
Illicit Drugs or Alcohol	7,822	7,865	2,059	2,388
Illicit Drugs	2,966	3,007	1,039	1,052
Alcohol	6,407	6,182	1,531	1,838

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15B (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Percentages, 2009 and 2010

Dependence or Abuse Status	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)
DEPENDENCE				
None	16.6	16.5	3.2	3.4
Both Illicit Drugs and Alcohol	63.7	62.7	20.1	26.8
Illicit Drugs or Alcohol	46.7	48.7	13.7	16.4
Illicit Drugs	51.3	52.7	18.6	20.5
Alcohol	46.8	48.3	12.1 ^a	15.6
DEPENDENCE OR ABUSE				
None	16.2	16.1	3.1	3.3
Both Illicit Drugs and Alcohol	55.6	54.9	18.3	20.8
Illicit Drugs or Alcohol	37.4	38.5	9.9	11.7
Illicit Drugs	49.0	50.2	17.2	17.6
Alcohol	36.3	36.7	8.7 ^a	10.9

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16A (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2009 and 2010

Dependence or Abuse Status	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
DEPENDENCE												
Both Illicit Drugs and Alcohol	1,240	1,033	789	648	249	277	198	152	343	218	451	385
Illicit Drugs or Alcohol	11,543	11,128	5,386	5,414	1,579	1,826	1,514	1,445	2,293	2,144	6,157	5,714
Illicit Drugs	4,316	4,228	2,214	2,230	803	869	531	590	881	771	2,101	1,999
Alcohol	8,467	7,933	3,961	3,833	1,024	1,234	1,181	1,008	1,755	1,591	4,506	4,100
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	2,787	2,413	1,551	1,324	511	503	400	310	641	511	1,236	1,089
Illicit Drugs or Alcohol	20,895	20,435	7,822	7,865	2,059	2,388	2,088	2,082	3,675	3,395	13,073	12,570
Illicit Drugs	6,056	5,990	2,966	3,007	1,039	1,052	711	770	1,217	1,185	3,090	2,983
Alcohol	17,626	16,858	6,407	6,182	1,531	1,838	1,778	1,623	3,099	2,721	11,219	10,676

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16B (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2009 and 2010

Dependence or Abuse Status	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.5	0.5	1.9	1.6	3.0	3.0	1.9	1.5	1.5	1.0	0.2	0.2
Illicit Drugs or Alcohol	5.1	4.9	13.1	13.1	18.8	19.6	14.9	14.6	10.1	9.6	3.3	3.0
Illicit Drugs	1.9	1.8	5.4	5.4	9.6	9.3	5.2	6.0	3.9	3.5	1.1	1.1
Alcohol	3.7	3.5	9.6	9.3	12.2	13.2	11.6	10.2	7.8	7.2	2.4	2.2
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	1.2 ^a	1.1	3.8	3.2	6.1	5.4	3.9	3.1	2.8	2.3	0.7	0.6
Illicit Drugs or Alcohol	9.2	8.9	19.0	19.0	24.6	25.6	20.5	21.1	16.2	15.3	7.0	6.7
Illicit Drugs	2.7	2.6	7.2	7.3	12.4	11.3	7.0	7.8	5.4	5.3	1.7	1.6
Alcohol	7.8	7.4	15.6	14.9	18.3	19.7	17.5	16.4	13.7	12.2	6.0	5.7

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17A (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	6,056	5,990	2,966	3,007	1,039	1,052	711	770	1,217	1,185	3,090	2,983
18-25	2,596	2,677	985	1,123	255	322	274	316	456	486	1,611	1,554
26-49	2,735	2,757	1,581	1,559	573	608	350	317	658	633	1,154	1,198
50 or Older	725	555	399	325	211	123	86	136	103	66	325	230
MALE	4,134	3,894	1,789	1,645	527	454	417	423	845	769	2,344	2,249
18-25	1,684	1,668	503	563	115	106	109 ^a	178	280	279	1,181	1,105
26-49	1,908	1,833	1,017	919	288	320	244	151	485	449	891	914
50 or Older	542	393	269	163	*	*	*	*	80	41	273	230
FEMALE	1,922	2,096	1,176	1,362	511	599	294	347	371	416	746	734
18-25	912	1,009	482	560	140 ^a	215	166	138	177	207	430	449
26-49	827	924	564	640	286	289	106	166	172	185	264	285
50 or Older	183	162	131	162	86	95	22	43	*	25	52	*

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17B (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	2.7	2.6	7.2	7.3	12.4	11.3	7.0	7.8	5.4	5.3	1.7	1.6
18-25	7.7	7.9	16.3	18.3	22.7	24.1	16.8	19.0	13.9	15.4	5.8	5.6
26-49	2.8	2.8	7.4	7.5	11.8	11.9	6.6	6.5	5.8	6.0	1.5	1.5
50 or Older	0.8	0.6	2.9	2.2	8.8	4.3	2.6	4.1	1.3	0.8	0.4	0.3
MALE	3.8	3.5	11.5	10.0	18.5	13.5	11.1	11.4	9.4	8.2	2.5	2.4
18-25	10.0	9.7	21.7	24.3	32.9	24.0	18.3 ^b	29.6	20.3	21.9	8.1	7.4
26-49	3.9	3.8	11.9	11.1	16.7	16.7	11.9	8.7	10.2	9.6	2.2	2.3
50 or Older	1.2	0.9	5.7	2.8	*	*	*	*	2.8	1.2	0.7	0.6
FEMALE	1.6	1.8	4.6	5.5	9.2	10.0	4.6	5.6	2.7	3.3	0.8	0.8
18-25	5.5	6.0	13.0	14.6	18.0 ^a	24.2	15.9	12.9	9.3	11.0	3.3	3.5
26-49	1.6	1.9	4.4	5.2	9.1	9.0	3.3	5.3	2.6	3.1	0.7	0.8
50 or Older	0.4	0.3	1.5	1.8	5.3	5.1	1.0	2.2	*	0.5	0.1	*

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18A (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	17,626	16,858	6,407	6,182	1,531	1,838	1,778	1,623	3,099	2,721	11,219	10,676
18-25	5,422	5,342	1,708	1,629	332 ^a	422	545	478	831	729	3,715	3,713
26-49	8,828	8,345	3,637	3,600	897 ^a	1,184	1,042	893	1,698	1,523	5,191	4,745
50 or Older	3,376	3,171	1,063	952	302	232	191	252	570	469	2,314	2,219
MALE	11,694	11,253	3,530	3,278	673	898	978	751	1,879	1,629	8,164	7,975
18-25	3,381	3,377	806	774	120	167	232	225	454	382	2,575	2,603
26-49	6,021	5,638	2,108	1,993	398 ^a	605	630	437	1,080	950	3,912	3,646
50 or Older	2,293	2,238	616	512	*	*	*	*	346	297	1,677	1,726
FEMALE	5,933	5,605	2,877	2,903	858	940	800	871	1,219	1,092	3,055	2,702
18-25	2,042	1,965	902	855	211	255	313	253	377	347	1,140	1,110
26-49	2,807	2,707	1,529	1,607	499	579	411	456	618	573	1,279	1,099
50 or Older	1,083	933	447	441	147	106	75	163	225	172	636	493

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18B (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	7.8	7.4	15.6	14.9	18.3	19.7	17.5	16.4	13.7	12.2	6.0	5.7
18-25	16.1	15.7	28.3	26.5	29.5	31.6	33.4	28.7	25.3	23.1	13.5	13.3
26-49	8.9	8.5	17.0	17.4	18.5	23.1	19.7	18.3	15.0	14.3	6.7	6.1
50 or Older	3.6	3.3	7.8	6.5	12.6	8.1	5.9	7.6	7.1	5.6	2.9	2.7
MALE	10.7	10.1	22.6	19.9	23.6	26.8	25.9	20.3	20.9	17.3	8.7	8.4
18-25	20.0	19.5	34.7	33.4	34.4	37.7	39.1	37.4	32.9	30.1	17.7	17.4
26-49	12.2	11.6	24.7	24.0	23.2	31.7	30.8	25.1	22.6	20.4	9.6	9.0
50 or Older	5.3	5.0	12.9	8.8	*	*	*	*	12.1	8.5	4.3	4.4
FEMALE	5.0	4.7	11.2	11.6	15.5	15.7	12.5	14.1	8.9	8.5	3.3	2.9
18-25	12.2	11.7	24.2	22.3	27.3	28.6	30.1 ^a	23.7	19.8	18.5	8.8	8.6
26-49	5.6	5.4	11.8	13.0	15.9	18.0	12.7	14.5	9.4	9.6	3.4	2.9
50 or Older	2.1	1.8	5.0	5.0	9.1	5.7	3.5	8.2	4.3	3.5	1.5	1.1

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.19A (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	20,895	20,435	7,822	7,865	2,059	2,388	2,088	2,082	3,675	3,395	13,073	12,570
18-25	6,753	6,803	2,122	2,182	446 ^a	570	645	621	1,032	991	4,631	4,621
26-49	10,326	10,000	4,455	4,477	1,227	1,508	1,208	1,073	2,020	1,897	5,871	5,522
50 or Older	3,816	3,632	1,245	1,206	386	310	236	388	622	508	2,571	2,426
MALE	13,818	13,576	4,293	4,131	918	1,069	1,143	989	2,232	2,073	9,525	9,445
18-25	4,224	4,284	1,003	1,025	166	207	265	302	572	515	3,221	3,259
26-49	6,996	6,739	2,576	2,487	556	736	736	504	1,285	1,246	4,420	4,252
50 or Older	2,598	2,553	713	619	*	*	*	*	375	311	1,885	1,934
FEMALE	7,077	6,859	3,529	3,734	1,141	1,319	945	1,093	1,443	1,322	3,548	3,125
18-25	2,529	2,519	1,119	1,157	280 ^a	363	380	319	460	475	1,410	1,362
26-49	3,330	3,261	1,878	1,990	671	772	472	569	736	650	1,451	1,271
50 or Older	1,218	1,079	531	586	191	185	94	205	247	196	687	493

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.19B (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	9.2	8.9	19.0	19.0	24.6	25.6	20.5	21.1	16.2	15.3	7.0	6.7
18-25	20.1	20.0	35.1	35.5	39.7	42.7	39.5	37.3	31.4	31.5	16.8	16.6
26-49	10.4	10.1	20.8	21.7	25.3	29.4	22.9	22.0	17.9	17.8	7.5	7.1
50 or Older	4.0	3.8	9.1	8.2	16.1	10.8	7.2	11.7	7.7	6.0	3.2	3.0
MALE	12.6	12.2	27.5	25.1	32.2	31.9	30.3	26.8	24.8	22.0	10.1	10.0
18-25	25.0	24.8	43.2	44.2	47.5	46.6	44.7	50.3	41.5	40.6	22.1	21.8
26-49	14.2	13.8	30.2	29.9	32.4	38.6	35.9	28.9	26.9	26.8	10.9	10.5
50 or Older	6.0	5.7	15.0	10.6	*	*	*	*	13.2	8.9	4.8	4.9
FEMALE	6.0	5.8	13.8	15.0	20.6	22.0	14.8	17.7	10.6	10.3	3.9 ^a	3.3
18-25	15.2	15.0	30.1	30.2	36.1	40.8	36.5 ^a	29.9	24.1	25.3	10.9	10.5
26-49	6.6	6.5	14.5	16.1	21.4	23.9	14.6	18.1	11.3	10.9	3.9	3.4
50 or Older	2.4	2.1	5.9	6.7	11.8	9.9	4.4	10.4	4.8	4.0	1.6	1.1

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.