

**Table 1.62A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics:
Numbers in Thousands, 2002-2010**

Demographic Characteristic	2002	2003	2004	2005	2006	2007	2008	2009	2010
TOTAL	27,215 ^b	27,939 ^b	27,502 ^b	28,166 ^b	28,346 ^b	29,401 ^a	30,002	30,179	31,314
AGE									
18-25	3,256 ^b	3,511	3,459	3,615	3,514	3,365 ^a	3,547	3,705	3,685
26-49	14,492	14,476	14,348	13,889	14,020	14,242	13,937	14,415	14,542
50 or Older	9,466 ^b	9,952 ^b	9,695 ^b	10,661 ^b	10,812 ^b	11,794	12,518	12,059	13,087
GENDER									
Male	8,772 ^b	8,689 ^b	9,044 ^b	9,308 ^a	9,458 ^a	9,836	10,053	10,107	10,473
Female	18,442 ^b	19,250 ^a	18,458 ^b	18,858 ^b	18,888 ^b	19,565	19,949	20,072	20,841
HISPANIC ORIGIN AND RACE									
Not Hispanic or Latino	25,172 ^b	25,919 ^b	25,523 ^b	25,999 ^b	26,320 ^b	27,241	27,951	27,906	28,813
White	21,943 ^b	22,914 ^b	22,512 ^b	22,890 ^b	23,279 ^a	24,647	24,637	24,754	25,261
Black or African American	1,953	2,021	2,052	2,193	1,855	1,711 ^b	2,230	2,021	2,344
American Indian or Alaska Native	227	136	143	169	130	128	124	*	145
Native Hawaiian or Other Pacific Islander	29	*	*	*	58	*	*	*	*
Asian	668	424	422	365	519	370	442	353	545
Two or More Races	352	373	335	317	480	359	464	463	480
Hispanic or Latino	2,043	2,020	1,979 ^a	2,166	2,025	2,160	2,051	2,274	2,501

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Because of revisions to the logical editing of the outpatient mental health treatment/counseling data, beginning with the 2004 NSDUH and retroactively applied to 2002 and 2003, these 2002 and 2003 estimates may differ slightly from 2002 and 2003 estimates published in NSDUH reports prior to the 2004 National Findings Report. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

Table 1.62B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2002-2010

Demographic Characteristic	2002	2003	2004	2005	2006	2007	2008	2009	2010
TOTAL	13.0	13.2	12.8 ^a	13.0	12.9 ^a	13.2	13.4	13.3	13.7
AGE									
18-25	10.5	11.1	10.8	11.2	10.8	10.3	10.8	11.1	10.9
26-49	14.5	14.5	14.4	13.9	14.0	14.3	14.0	14.6	14.8
50 or Older	12.0 ^a	12.3	11.7 ^a	12.5	12.4	13.2	13.6	12.8	13.6
GENDER									
Male	8.7	8.5 ^a	8.8	8.9	8.9	9.2	9.3	9.2	9.5
Female	16.9	17.5	16.6	16.8	16.6	17.0	17.2	17.1	17.7
HISPANIC ORIGIN AND RACE									
Not Hispanic or Latino	13.6 ^a	13.9	13.6 ^a	13.7 ^a	13.8 ^a	14.2	14.4	14.3	14.6
White	14.7 ^b	15.2 ^a	14.9 ^b	15.1 ^a	15.2 ^a	16.0	16.0	16.0	16.2
Black or African American	8.5	8.5	8.5	8.9	7.4	6.8 ^a	8.7	7.7	8.8
American Indian or Alaska Native	17.7	12.6	12.6	15.4	11.9	11.6	13.2	*	13.5
Native Hawaiian or Other Pacific Islander	3.9	*	*	*	7.0	*	*	*	*
Asian	8.3	4.9	4.7	4.0	5.6	3.9	4.5	3.5	5.3
Two or More Races	16.6	17.5	15.8	14.4	21.6	15.6	18.8	19.1	18.1
Hispanic or Latino	8.2	7.8	7.4	7.8	7.0	7.3	6.8	7.3	7.9

*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Because of revisions to the logical editing of the outpatient mental health treatment/counseling data, beginning with the 2004 NSDUH and retroactively applied to 2002 and 2003, these 2002 and 2003 estimates may differ slightly from 2002 and 2003 estimates published in NSDUH reports prior to the 2004 National Findings Report. See Section B.5.2 in Appendix B of the *Results from the 2004 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.