

**Table 1.13C (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
<b>ILLCIT DRUGS<sup>1</sup></b>	583	631	326	343	153	177	159	152	224	230	510	554
Marijuana and Hashish	527	548	273	273	124	142	136	130	183	176	456	478
Cocaine	208	202	127	119	76	58	40	49	83	88	160	165
Crack	105	92	71	71	44	45	25	31	46	42	75	59
Heroin	78	75	41	54	33	37	15	14	19	36	64	51
Hallucinogens	151	170	87	94	39	60	49	42	60	54	124	133
LSD	53	69	29	45	16	27	16	28	17	20	42	48
PCP	19	23	14	21	2	20	13	3	4	3	13	9
Ecstasy	123	121	76	68	33	34	42	34	55	44	100	94
Inhalants	101	109	68	54	19	35	49	29	42	28	75	94
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	360	389	230	240	114	132	114	119	163	157	285	317
Pain Relievers	318	329	198	210	100	122	106	109	132	129	244	258
OxyContin <sup>®</sup>	102	114	71	87	46	69	33	42	43	31	69	71
Tranquilizers	220	205	150	148	74	71	77	94	109	96	160	154
Stimulants <sup>3</sup>	146	155	102	91	55	53	49	43	69	59	109	117
Methamphetamine <sup>3</sup>	100	103	70	53	51	33	30	22	39	35	71	85
Sedatives	78	103	50	62	35	43	22	23	28	39	59	82
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	410	433	254	265	127	143	124	127	180	173	332	345

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.13D (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2009 and 2010**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Any Mental Illness (2009)</b>	<b>Any Mental Illness (2010)</b>	<b>Serious Mental Illness (2009)</b>	<b>Serious Mental Illness (2010)</b>	<b>Moderate Mental Illness (2009)</b>	<b>Moderate Mental Illness (2010)</b>	<b>Mild Mental Illness (2009)</b>	<b>Mild Mental Illness (2010)</b>	<b>No Mental Illness (2009)</b>	<b>No Mental Illness (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	0.26	0.28	0.70	0.74	1.52	1.61	1.42	1.46	0.90	0.95	0.27	0.28
Marijuana and Hashish	0.23	0.24	0.62	0.61	1.37	1.38	1.26	1.29	0.77	0.76	0.24	0.24
Cocaine	0.09	0.09	0.30	0.29	0.88	0.61	0.41	0.50	0.36	0.40	0.09	0.09
Crack	0.05	0.04	0.17	0.17	0.52	0.48	0.25	0.32	0.20	0.19	0.04	0.03
Heroin	0.03	0.03	0.10	0.13	0.39	0.39	0.14	0.15	0.08	0.16	0.03	0.03
Hallucinogens	0.07	0.07	0.21	0.23	0.47	0.64	0.49	0.43	0.26	0.24	0.07	0.07
LSD	0.02	0.03	0.07	0.11	0.19	0.29	0.16	0.29	0.07	0.09	0.02	0.03
PCP	0.01	0.01	0.03	0.05	0.02	0.22	0.13	0.03	0.02	0.02	0.01	0.00
Ecstasy	0.05	0.05	0.18	0.16	0.40	0.36	0.42	0.34	0.24	0.20	0.05	0.05
Inhalants	0.04	0.05	0.16	0.13	0.23	0.37	0.48	0.29	0.19	0.13	0.04	0.05
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.16	0.17	0.52	0.54	1.22	1.29	1.07	1.15	0.68	0.67	0.15	0.17
Pain Relievers	0.14	0.14	0.45	0.48	1.11	1.21	0.99	1.05	0.56	0.57	0.13	0.14
OxyContin <sup>®</sup>	0.04	0.05	0.17	0.21	0.54	0.72	0.33	0.42	0.19	0.14	0.04	0.04
Tranquilizers	0.10	0.09	0.35	0.35	0.84	0.75	0.73	0.92	0.47	0.42	0.09	0.08
Stimulants <sup>3</sup>	0.06	0.07	0.24	0.22	0.63	0.57	0.48	0.43	0.30	0.26	0.06	0.06
Methamphetamine <sup>3</sup>	0.04	0.04	0.17	0.13	0.59	0.36	0.29	0.23	0.17	0.16	0.04	0.05
Sedatives	0.03	0.04	0.12	0.15	0.41	0.45	0.21	0.23	0.12	0.18	0.03	0.04
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.18	0.19	0.57	0.60	1.34	1.39	1.16	1.23	0.75	0.75	0.18	0.18

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.13P (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Any Mental Illness (2009)</b>	<b>Any Mental Illness (2010)</b>	<b>Serious Mental Illness (2009)</b>	<b>Serious Mental Illness (2010)</b>	<b>Moderate Mental Illness (2009)</b>	<b>Moderate Mental Illness (2010)</b>	<b>Mild Mental Illness (2009)</b>	<b>Mild Mental Illness (2010)</b>	<b>No Mental Illness (2009)</b>	<b>No Mental Illness (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	0.4980		0.7900		0.6799		0.8183		0.4187		0.3225	
Marijuana and Hashish	0.6557		0.6345		0.8609		0.7087		0.4499		0.4100	
Cocaine	0.2844		0.2995		0.0827		0.2893		0.4307		0.5694	
Crack	0.3844		0.5591		0.3031		0.3185		0.4619		0.5078	
Heroin	0.7105		0.7206		0.7078		0.8153		0.3585		0.8407	
Hallucinogens	0.8659		0.7008		0.3446		0.9908		0.1264		0.9410	
LSD	0.3065		0.2772		0.5366		0.1433		0.7102		0.5853	
PCP	0.5879		0.9564		0.3127		0.1611		0.9590		0.2853	
Ecstasy	0.1976		0.3169		0.9337		0.7872		0.1855		0.3870	
Inhalants	0.9810		0.6482		0.2984		0.3299		0.5888		0.7046	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.9109		0.8864		0.9340		0.8716		0.4890		1.0000	
Pain Relievers	0.6546		0.9270		0.8266		0.8793		0.5859		0.6415	
OxyContin <sup>®</sup>	0.2656		0.3668		0.4207		0.2103		0.3529		0.4972	
Tranquilizers	0.8927		0.8678		0.7595		0.4940		0.4499		0.7136	
Stimulants <sup>3</sup>	0.3295		0.1271		0.1522		0.3484		0.5754		0.9890	
Methamphetamine <sup>3</sup>	0.1257		0.0177		0.0134		0.1531		0.8113		0.8553	
Sedatives	0.4352		0.3989		0.6149		0.7697		0.6816		0.7416	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.3743		0.7120		0.9597		0.7747		0.2715		0.4176	

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.14C (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2009 and 2010**

Substance	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
<b>PAST YEAR</b>												
Tobacco Products <sup>1</sup>	851	906	476	484	200	235	230	227	342	344	921	982
Cigarettes	829	828	447	442	196	226	218	215	317	304	822	845
Smokeless Tobacco	333	345	137	145	65	68	66	67	102	109	311	325
Cigars	497	504	236	224	89	116	123	105	178	162	451	507
Alcohol	883	953	645	644	270	274	307	281	447	470	1,685	1,750
<b>PAST MONTH</b>												
Tobacco Products <sup>1</sup>	822	840	449	457	197	220	215	217	318	313	859	867
Cigarettes	790	772	419	422	193	214	202	205	289	281	770	770
Smokeless Tobacco	288	310	117	131	55	59	55	57	84	100	271	291
Cigars	358	358	169	169	64	81	103	86	116	126	320	330
Pipe Tobacco	149	147	70	88	18	36	59	40	33	69	133	119
Alcohol	945	974	559	556	210	233	254	237	406	407	1,494	1,502
Binge Alcohol Use <sup>2</sup>	784	748	367	348	149	163	166	154	265	255	843	814
Heavy Alcohol Use <sup>2</sup>	404	421	185	187	80	91	90	82	136	129	378	402

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.14D (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2009 and 2010**

Substance	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
<b>PAST YEAR</b>												
Tobacco Products <sup>1</sup>	0.37	0.40	0.86	0.87	1.89	1.80	1.76	1.75	1.19	1.21	0.40	0.43
Cigarettes	0.36	0.36	0.86	0.83	1.89	1.79	1.75	1.69	1.16	1.13	0.38	0.39
Smokeless Tobacco	0.15	0.15	0.32	0.34	0.78	0.72	0.64	0.68	0.44	0.47	0.16	0.17
Cigars	0.22	0.22	0.53	0.52	1.02	1.16	1.12	1.03	0.74	0.71	0.23	0.24
Alcohol	0.39	0.42	0.84	0.87	1.74	1.83	1.60	1.70	1.17	1.20	0.43	0.45
<b>PAST MONTH</b>												
Tobacco Products <sup>1</sup>	0.36	0.37	0.86	0.86	1.89	1.77	1.76	1.74	1.16	1.18	0.39	0.40
Cigarettes	0.35	0.34	0.83	0.82	1.87	1.74	1.72	1.67	1.09	1.12	0.37	0.37
Smokeless Tobacco	0.13	0.14	0.28	0.31	0.65	0.62	0.53	0.58	0.36	0.43	0.14	0.15
Cigars	0.16	0.16	0.39	0.40	0.75	0.83	0.96	0.85	0.50	0.56	0.17	0.17
Pipe Tobacco	0.07	0.06	0.17	0.21	0.21	0.38	0.57	0.40	0.14	0.31	0.07	0.06
Alcohol	0.42	0.42	0.89	0.93	1.89	1.79	1.67	1.80	1.24	1.31	0.47	0.47
Binge Alcohol Use <sup>2</sup>	0.34	0.33	0.75	0.72	1.57	1.54	1.48	1.46	1.02	1.04	0.38	0.36
Heavy Alcohol Use <sup>2</sup>	0.18	0.18	0.43	0.43	0.92	0.94	0.86	0.82	0.59	0.57	0.20	0.20

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.14P (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Substance	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
<b>PAST YEAR</b>												
Tobacco Products <sup>1</sup>	0.6980		0.8067		0.5103		0.3761		0.7500		0.5984	
Cigarettes	0.3806		0.7720		0.6058		0.2341		0.8820		0.4085	
Smokeless Tobacco	0.6745		0.9577		0.2924		0.9112		0.5184		0.6562	
Cigars	0.5446		0.8495		0.2736		0.2421		0.8566		0.5564	
Alcohol	0.5900		0.7560		0.6661		0.1916		0.5065		0.6444	
<b>PAST MONTH</b>												
Tobacco Products <sup>1</sup>	0.6664		0.7951		0.8123		0.4610		0.9148		0.7359	
Cigarettes	0.4926		0.4722		0.8103		0.3567		0.6851		0.6920	
Smokeless Tobacco	0.6229		0.6771		0.2297		0.9754		0.1990		0.7212	
Cigars	0.9730		0.6038		0.4546		0.3519		0.3268		0.8201	
Pipe Tobacco	0.5939		0.0824		0.0045		0.1842		0.0200		0.6352	
Alcohol	0.9436		0.7769		0.5304		0.2720		0.9887		0.9595	
Binge Alcohol Use <sup>2</sup>	0.1921		0.1877		0.7757		0.2829		0.2211		0.4066	
Heavy Alcohol Use <sup>2</sup>	0.5806		0.5470		0.4449		0.7160		0.2886		0.7516	

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.