

Table 1.15C (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Numbers in Thousands, 2009 and 2010

Dependence or Abuse Status	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)
DEPENDENCE				
None	748	744	302	330
Both Illicit Drugs and Alcohol	78	73	38	46
Illicit Drugs or Alcohol	228	247	119	138
Illicit Drugs	130	150	77	96
Alcohol	201	208	97	108
DEPENDENCE OR ABUSE				
None	721	719	294	322
Both Illicit Drugs and Alcohol	123	98	73	63
Illicit Drugs or Alcohol	281	287	137	155
Illicit Drugs	161	176	96	104
Alcohol	257	251	121	130

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15D (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Standard Errors of Percentages, 2009 and 2010

Dependence or Abuse Status	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)
DEPENDENCE				
None	0.31	0.30	0.14	0.15
Both Illicit Drugs and Alcohol	3.78	3.98	2.83	3.78
Illicit Drugs or Alcohol	1.44	1.52	0.97	1.13
Illicit Drugs	2.25	2.23	1.65	1.98
Alcohol	1.75	1.85	1.07	1.29
DEPENDENCE OR ABUSE				
None	0.31	0.31	0.14	0.15
Both Illicit Drugs and Alcohol	2.73	2.58	2.31	2.26
Illicit Drugs or Alcohol	1.05	1.09	0.62	0.72
Illicit Drugs	1.90	1.85	1.45	1.54
Alcohol	1.18	1.20	0.65	0.75

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.15P (REVISED) Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Dependence or Abuse Status	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)
DEPENDENCE				
None	0.8141		0.1539	
Both Illicit Drugs and Alcohol	0.8665		0.1694	
Illicit Drugs or Alcohol	0.3287		0.0663	
Illicit Drugs	0.6543		0.4548	
Alcohol	0.5301		0.0411	
DEPENDENCE OR ABUSE				
None	0.8011		0.2037	
Both Illicit Drugs and Alcohol	0.8385		0.4462	
Illicit Drugs or Alcohol	0.4881		0.0574	
Illicit Drugs	0.6430		0.8444	
Alcohol	0.8504		0.0291	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16C (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2009 and 2010

Dependence or Abuse Status	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
DEPENDENCE												
Both Illicit Drugs and Alcohol	100	86	78	73	38	46	38	31	54	44	60	47
Illicit Drugs or Alcohol	336	361	228	247	119	138	121	128	145	157	255	254
Illicit Drugs	180	201	130	150	77	96	64	86	79	74	136	125
Alcohol	291	302	201	208	97	108	109	102	132	144	221	218
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	154	125	123	98	73	63	65	40	71	63	95	79
Illicit Drugs or Alcohol	452	460	281	287	137	155	139	144	197	183	372	385
Illicit Drugs	218	237	161	176	96	104	82	94	97	100	158	147
Alcohol	414	409	257	251	121	130	130	123	183	168	351	358

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.16D (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2009 and 2010

Dependence or Abuse Status	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.04	0.04	0.19	0.17	0.46	0.48	0.37	0.32	0.24	0.19	0.03	0.03
Illicit Drugs or Alcohol	0.15	0.16	0.52	0.56	1.32	1.33	1.12	1.21	0.63	0.68	0.14	0.13
Illicit Drugs	0.08	0.09	0.31	0.35	0.89	0.98	0.62	0.84	0.35	0.33	0.07	0.07
Alcohol	0.13	0.13	0.47	0.48	1.09	1.08	1.02	1.00	0.58	0.62	0.12	0.11
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.07	0.05	0.29	0.24	0.84	0.65	0.63	0.41	0.31	0.28	0.05	0.04
Illicit Drugs or Alcohol	0.20	0.20	0.62	0.64	1.48	1.47	1.28	1.34	0.82	0.78	0.20	0.19
Illicit Drugs	0.10	0.10	0.38	0.42	1.09	1.05	0.79	0.93	0.42	0.45	0.09	0.08
Alcohol	0.18	0.18	0.58	0.56	1.33	1.27	1.21	1.18	0.77	0.72	0.18	0.18

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.16P (REVISED) Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Dependence or Abuse Status	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
DEPENDENCE												
Both Illicit Drugs and Alcohol	0.1070		0.1788		1.0000		0.4166		0.0883		0.3812	
Illicit Drugs or Alcohol	0.2901		1.0000		0.6955		0.8882		0.5939		0.1427	
Illicit Drugs	0.6427		0.9865		0.8407		0.4675		0.3814		0.4956	
Alcohol	0.1538		0.5929		0.5214		0.3274		0.4814		0.1325	
DEPENDENCE OR ABUSE												
Both Illicit Drugs and Alcohol	0.0438		0.1353		0.5167		0.2950		0.2150		0.2021	
Illicit Drugs or Alcohol	0.3147		1.0000		0.6249		0.7582		0.4004		0.2124	
Illicit Drugs	0.7064		0.9147		0.4631		0.5014		0.9467		0.5207	
Alcohol	0.1143		0.4515		0.4474		0.5504		0.1692		0.1712	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17C (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	218	237	161	176	96	104	82	94	97	100	158	147
18-25	83	85	49	57	25	31	26	29	33	35	72	69
26-49	159	178	120	135	68	80	59	55	80	89	113	115
50 or Older	123	114	94	94	65	54	48	70	44	28	81	64
MALE	190	187	137	133	75	66	74	78	84	86	139	130
18-25	66	65	35	41	17	19	17	22	26	28	64	58
26-49	137	142	104	104	52	60	53	37	72	76	98	98
50 or Older	110	96	82	72	*	*	*	*	37	24	75	64
FEMALE	113	131	88	111	61	81	37	53	48	50	72	63
18-25	48	51	35	38	18	25	20	18	21	21	33	36
26-49	83	98	63	80	45	53	27	40	36	43	56	53
50 or Older	55	58	46	58	37	50	15	26	*	15	29	*

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17D (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.10	0.10	0.38	0.42	1.09	1.05	0.79	0.93	0.42	0.45	0.09	0.08
18-25	0.25	0.25	0.75	0.80	1.96	1.96	1.48	1.54	0.94	1.00	0.25	0.24
26-49	0.16	0.18	0.53	0.62	1.33	1.45	1.09	1.11	0.68	0.79	0.14	0.15
50 or Older	0.13	0.12	0.68	0.64	2.58	1.83	1.45	2.04	0.54	0.34	0.10	0.08
MALE	0.17	0.17	0.83	0.79	2.40	1.85	1.86	2.03	0.90	0.92	0.15	0.14
18-25	0.39	0.38	1.33	1.47	4.26	3.65	2.62	3.07	1.65	1.90	0.40	0.37
26-49	0.28	0.29	1.15	1.17	2.83	2.81	2.41	2.07	1.41	1.55	0.23	0.24
50 or Older	0.25	0.21	1.67	1.21	*	*	*	*	1.30	0.68	0.19	0.16
FEMALE	0.10	0.11	0.34	0.43	1.08	1.28	0.60	0.84	0.36	0.39	0.08	0.07
18-25	0.29	0.30	0.87	0.90	2.10	2.31	1.71	1.61	1.06	1.07	0.25	0.27
26-49	0.16	0.20	0.48	0.62	1.38	1.55	0.83	1.24	0.55	0.70	0.15	0.14
50 or Older	0.11	0.11	0.51	0.66	2.26	2.59	0.70	1.29	*	0.30	0.07	*

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.17P (REVISED) Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.7064		0.9147		0.4631		0.5014		0.9467		0.5207	
18-25	0.7170		0.0768		0.6031		0.3054		0.2673		0.4087	
26-49	0.8500		0.8231		0.9799		0.9367		0.8882		0.7769	
50 or Older	0.2652		0.4548		0.1564		0.5584		0.4363		0.3306	
MALE	0.2778		0.1963		0.1000		0.8880		0.3281		0.5765	
18-25	0.5577		0.1832		0.1160		0.0048		0.4973		0.1903	
26-49	0.7701		0.5961		1.0000		0.3072		0.7959		0.8293	
50 or Older	0.2661		0.1634		*		*		0.2660		0.6514	
FEMALE	0.3604		0.1158		0.6418		0.3164		0.2887		0.8118	
18-25	0.2001		0.1934		0.0445		0.1980		0.2683		0.6845	
26-49	0.4304		0.2909		0.9440		0.1711		0.5670		0.8041	
50 or Older	0.7634		0.6409		0.9465		0.4376		*		*	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18C (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	414	409	257	251	121	130	130	123	183	168	351	358
18-25	117	125	69	68	27	36	40	35	47	41	123	121
26-49	289	280	186	199	86	110	107	98	129	135	246	228
50 or Older	263	262	149	134	74	59	58	66	112	100	225	232
MALE	347	332	199	177	83	94	104	85	149	123	303	312
18-25	91	99	49	49	18	24	27	26	35	31	99	101
26-49	245	232	147	150	56	78	87	72	106	106	222	202
50 or Older	224	223	119	95	*	*	*	*	92	76	192	208
FEMALE	235	225	155	174	81	90	77	94	105	109	180	150
18-25	67	70	47	47	20	28	29	23	31	28	58	58
26-49	152	149	116	132	63	78	63	72	75	80	109	93
50 or Older	151	140	86	94	48	41	28	55	64	65	126	102

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18D (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.18	0.18	0.58	0.56	1.33	1.27	1.21	1.18	0.77	0.72	0.18	0.18
18-25	0.35	0.37	0.96	0.90	2.06	2.15	1.88	1.79	1.23	1.15	0.37	0.37
26-49	0.29	0.28	0.78	0.85	1.63	1.89	1.80	1.78	1.05	1.15	0.30	0.27
50 or Older	0.28	0.27	1.04	0.90	2.92	2.05	1.75	1.96	1.34	1.16	0.28	0.28
MALE	0.32	0.30	1.14	1.01	2.63	2.52	2.43	2.19	1.51	1.26	0.32	0.31
18-25	0.54	0.57	1.64	1.68	4.19	4.02	3.20	3.22	2.00	2.07	0.56	0.58
26-49	0.50	0.48	1.50	1.54	3.00	3.42	3.43	3.50	1.95	2.02	0.52	0.46
50 or Older	0.51	0.50	2.32	1.58	*	*	*	*	3.01	2.11	0.49	0.52
FEMALE	0.20	0.19	0.59	0.67	1.41	1.41	1.20	1.41	0.76	0.84	0.19	0.16
18-25	0.40	0.41	1.12	1.07	2.28	2.52	2.24	1.96	1.47	1.39	0.41	0.42
26-49	0.30	0.30	0.83	1.01	1.86	2.18	1.86	2.09	1.10	1.30	0.28	0.24
50 or Older	0.30	0.27	0.94	1.05	2.84	2.15	1.33	2.71	1.20	1.30	0.30	0.24

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.18P (REVISED) Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.1143		0.4515		0.4474		0.5504		0.1692		0.1712	
18-25	0.3513		0.1794		0.4820		0.0767		0.2057		0.7098	
26-49	0.3183		0.6787		0.0649		0.5794		0.6622		0.1624	
50 or Older	0.4349		0.3727		0.2110		0.5152		0.3890		0.6645	
MALE	0.2410		0.0802		0.3842		0.0856		0.0719		0.5900	
18-25	0.5448		0.5716		0.5733		0.7096		0.3234		0.7239	
26-49	0.3505		0.7513		0.0551		0.2456		0.4413		0.3942	
50 or Older	0.6946		0.1392		*		*		0.3237		0.8912	
FEMALE	0.2541		0.6684		0.9182		0.3847		0.7155		0.0807	
18-25	0.3555		0.2133		0.7041		0.0317		0.5227		0.6960	
26-49	0.6971		0.3673		0.4872		0.5083		0.9438		0.1718	
50 or Older	0.4008		0.9869		0.3380		0.1195		0.6322		0.3094	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.19C (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	452	460	281	287	137	155	139	144	197	183	372	385
18-25	128	139	77	80	32	43	42	40	53	49	137	134
26-49	319	310	207	225	103	126	114	107	142	150	266	247
50 or Older	274	280	159	157	81	76	66	91	115	103	232	239
MALE	377	362	218	205	96	104	113	111	161	137	323	335
18-25	99	106	54	56	21	26	28	30	40	37	113	111
26-49	273	250	165	170	70	90	93	77	117	121	241	214
50 or Older	232	238	126	116	*	*	*	*	94	78	199	216
FEMALE	257	251	171	197	92	116	82	106	111	113	190	158
18-25	75	78	53	57	24	34	32	26	34	34	64	64
26-49	163	167	127	146	72	90	66	81	81	83	118	101
50 or Older	159	152	93	110	52	62	32	61	67	67	129	102

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.19D (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2009 and 2010

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.20	0.20	0.62	0.64	1.48	1.47	1.28	1.34	0.82	0.78	0.20	0.19
18-25	0.38	0.41	1.03	0.97	2.27	2.29	1.91	1.94	1.34	1.28	0.41	0.42
26-49	0.32	0.31	0.84	0.93	1.88	2.03	1.90	1.91	1.13	1.23	0.33	0.29
50 or Older	0.29	0.29	1.10	1.04	3.16	2.55	1.94	2.59	1.38	1.18	0.28	0.29
MALE	0.34	0.33	1.21	1.15	2.97	2.69	2.58	2.71	1.59	1.41	0.34	0.33
18-25	0.59	0.62	1.72	1.73	4.76	4.03	3.21	3.37	2.13	2.21	0.61	0.63
26-49	0.56	0.51	1.62	1.66	3.60	3.60	3.55	3.65	2.10	2.23	0.56	0.49
50 or Older	0.53	0.53	2.41	1.90	*	*	*	*	3.06	2.15	0.50	0.54
FEMALE	0.22	0.21	0.64	0.74	1.56	1.74	1.28	1.58	0.81	0.87	0.20	0.17
18-25	0.45	0.47	1.20	1.21	2.52	2.78	2.30	2.17	1.61	1.62	0.45	0.45
26-49	0.33	0.33	0.89	1.09	2.05	2.40	1.94	2.34	1.18	1.33	0.31	0.26
50 or Older	0.31	0.29	1.02	1.21	3.10	3.18	1.50	2.97	1.27	1.33	0.31	0.24

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.19P (REVISED) Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Gender/ Age Group	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
TOTAL	0.3147		1.0000		0.6249		0.7582		0.4004		0.2124	
18-25	0.7962		0.8103		0.3469		0.4215		0.9821		0.6456	
26-49	0.5935		0.4744		0.1259		0.7491		0.9855		0.3078	
50 or Older	0.4718		0.5815		0.1761		0.1798		0.3465		0.5543	
MALE	0.4510		0.1504		0.9285		0.3384		0.1969		0.7823	
18-25	0.7938		0.6699		0.8838		0.2365		0.7639		0.7032	
26-49	0.6163		0.9244		0.2146		0.1744		0.9698		0.6395	
50 or Older	0.7068		0.1488		*		*		0.2547		0.8974	
FEMALE	0.4535		0.2395		0.5372		0.1540		0.8287		0.0473	
18-25	0.8175		0.9578		0.2099		0.0355		0.6197		0.5653	
26-49	0.8389		0.2681		0.4191		0.2349		0.8243		0.2051	
50 or Older	0.4570		0.6472		0.6600		0.0856		0.6725		0.1872	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.