

Table 1.32C (REVISED) Type and Location of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2009 and 2010

Type and Location of Mental Health Treatment/Counseling	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
ANY MENTAL HEALTH SERVICES¹	603	616	467	510	275	302	239	240	288	308	467	479
Outpatient	420	448	347	368	234	244	164	183	206	194	264	274
Outpatient Mental Health Clinic or Center	177	197	156	179	102	116	76	90	92	103	81	98
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	314	345	256	267	152	188	118	121	154	144	178	212
Doctor's Office - Not Part of a Clinic	235	204	201	160	140	101	84	100	117	73	128	139
Outpatient Medical Clinic	138	110	114	88	86	69	55	37	50	41	77	65
Partial Day Hospital or Day Treatment Program	65	54	59	46	38	39	25	17	37	20	28	26
School or University Setting/Clinic/Center ²	18	16	16	11	12	6	3	7	10	6	7	11
Some Other Place ³	48	67	41	58	18	16	22	48	29	29	26	34
Inpatient	154	154	121	104	86	72	45	39	75	58	96	118
Prescription Medication	564	557	422	488	264	284	223	225	252	292	438	412
DID NOT RECEIVE MENTAL HEALTH SERVICES	603	616	608	570	167	188	267	248	502	460	1,991	2,084

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple service sources; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.32D (REVISED) Type and Location of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Standard Errors of Percentages, 2009 and 2010

Type and Location of Mental Health Treatment/Counseling	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
ANY MENTAL HEALTH SERVICES¹	0.27	0.27	0.86	0.89	1.68	1.67	1.75	1.75	1.11	1.11	0.24	0.24
Outpatient	0.19	0.20	0.74	0.78	1.97	1.89	1.44	1.57	0.86	0.82	0.14	0.14
Outpatient Mental Health Clinic or Center	0.08	0.09	0.37	0.42	1.19	1.18	0.72	0.90	0.41	0.46	0.04	0.05
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.14	0.15	0.59	0.60	1.58	1.67	1.09	1.16	0.66	0.62	0.10	0.11
Doctor's Office - Not Part of a Clinic	0.10	0.09	0.47	0.37	1.52	1.03	0.81	0.97	0.51	0.33	0.07	0.07
Outpatient Medical Clinic	0.06	0.05	0.27	0.21	0.97	0.72	0.54	0.38	0.22	0.19	0.04	0.03
Partial Day Hospital or Day Treatment Program	0.03	0.02	0.14	0.11	0.45	0.41	0.25	0.17	0.17	0.09	0.02	0.01
School or University Setting/Clinic/Center ²	0.01	0.01	0.04	0.03	0.15	0.06	0.03	0.07	0.04	0.02	0.00	0.01
Some Other Place ³	0.02	0.03	0.10	0.14	0.21	0.17	0.22	0.49	0.13	0.13	0.01	0.02
Inpatient	0.07	0.07	0.29	0.25	0.98	0.77	0.44	0.39	0.33	0.26	0.05	0.06
Prescription Medication	0.25	0.24	0.82	0.90	1.77	1.80	1.72	1.72	1.00	1.09	0.22	0.21
DID NOT RECEIVE MENTAL HEALTH SERVICES	0.27	0.27	0.86	0.89	1.68	1.67	1.75	1.75	1.11	1.11	0.24	0.24

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple service sources; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.32P (REVISED) Type and Location of Mental Health Treatment/Counseling Received in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Type and Location of Mental Health Treatment/Counseling	Total (2009)	Total (2010)	Any Mental Illness (2009)	Any Mental Illness (2010)	Serious Mental Illness (2009)	Serious Mental Illness (2010)	Moderate Mental Illness (2009)	Moderate Mental Illness (2010)	Mild Mental Illness (2009)	Mild Mental Illness (2010)	No Mental Illness (2009)	No Mental Illness (2010)
ANY MENTAL HEALTH SERVICES¹	0.3113		0.0727		0.6930		0.3835		0.3541		0.9905	
Outpatient	0.3027		0.3782		0.4389		0.0905		0.8982		0.4295	
Outpatient Mental Health Clinic or Center	0.4222		0.6628		0.5834		0.1859		0.7266		0.3258	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.1120		0.2368		0.2623		0.8362		0.9468		0.1945	
Doctor's Office - Not Part of a Clinic	0.4896		0.3709		0.1118		0.2162		0.2470		0.9664	
Outpatient Medical Clinic	0.1638		0.1264		0.0682		0.6638		0.7172		0.8725	
Partial Day Hospital or Day Treatment Program	0.3574		0.4336		0.6796		0.6698		0.3921		0.6423	
School or University Setting/Clinic/Center ²	0.8915		0.3178		0.3507		0.3225		0.1852		0.2228	
Some Other Place ³	0.5458		0.4746		0.7863		0.3525		0.8391		0.9951	
Inpatient	0.4511		0.1625		0.1319		0.6997		0.3546		0.6690	
Prescription Medication	0.3515		0.0677		0.9703		0.3697		0.2234		0.8074	
DID NOT RECEIVE MENTAL HEALTH SERVICES	0.3113		0.0727		0.6930		0.3835		0.3541		0.9905	

*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Respondents could indicate multiple service sources; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location was the most commonly reported other location for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.