

Table 1.49C Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2009 and 2010

Illicit Drug or Alcohol Dependence or Abuse Status¹/Level of Mental Illness	Had Serious Thoughts of Suicide (2009)	Had Serious Thoughts of Suicide (2010)	Made Any Suicide Plans (2009)	Made Any Suicide Plans (2010)	Attempted Suicide (2009)	Attempted Suicide (2010)
DEPENDENCE OR ABUSE	130	152	82	60	46	42
Any Mental Illness	123	141	81	58	46	40
Serious Mental Illness	89	94	67	50	38	34
Moderate Mental Illness	53	59	36	15	14	13
Mild Mental Illness	62	83	28	24	21	16
No Mental Illness	43	56	15	15	6	13
NO DEPENDENCE OR ABUSE	272	264	134	155	78	98
Any Mental Illness	227	228	109	124	60	66
Serious Mental Illness	153	154	97	93	55	50
Moderate Mental Illness	112	105	38	54	17	37
Mild Mental Illness	131	121	31	69	17	19
No Mental Illness	144	161	63	93	49	73

*Low precision; no estimate reported.

NOTE: Estimates in this table are based only on responses to suicide and mental illness items in the Mental Health module. Respondents with unknown suicide information were excluded.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.49D Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Standard Errors of Percentages, 2009 and 2010

Illicit Drug or Alcohol Dependence or Abuse Status¹/Level of Mental Illness	Had Serious Thoughts of Suicide (2009)	Had Serious Thoughts of Suicide (2010)	Made Any Suicide Plans (2009)	Made Any Suicide Plans (2010)	Attempted Suicide (2009)	Attempted Suicide (2010)
DEPENDENCE OR ABUSE	0.60	0.71	0.39	0.30	0.22	0.21
Any Mental Illness	1.28	1.35	0.87	0.63	0.52	0.44
Serious Mental Illness	2.58	2.60	2.13	1.67	1.32	1.18
Moderate Mental Illness	2.51	2.66	1.79	0.78	0.75	0.67
Mild Mental Illness	1.44	1.77	0.66	0.57	0.51	0.38
No Mental Illness	0.36	0.49	0.12	0.14	0.05	0.12
NO DEPENDENCE OR ABUSE	0.13	0.13	0.06	0.07	0.04	0.05
Any Mental Illness	0.58	0.58	0.30	0.33	0.17	0.18
Serious Mental Illness	1.60	1.59	1.15	1.03	0.67	0.58
Moderate Mental Illness	1.45	1.38	0.53	0.77	0.24	0.53
Mild Mental Illness	0.61	0.57	0.15	0.32	0.08	0.09
No Mental Illness	0.09	0.09	0.04	0.05	0.03	0.04

*Low precision; no estimate reported.

NOTE: Estimates in this table are based only on responses to suicide and mental illness items in the Mental Health module. Respondents with unknown suicide information were excluded.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.49P Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Illicit Drug or Alcohol Dependence or Abuse Status¹/Level of Mental Illness	Had Serious Thoughts of Suicide (2009)	Had Serious Thoughts of Suicide (2010)	Made Any Suicide Plans (2009)	Made Any Suicide Plans (2010)	Attempted Suicide (2009)	Attempted Suicide (2010)
DEPENDENCE OR ABUSE	0.1416		0.3286		0.7063	
Any Mental Illness	0.1614		0.1526		0.2700	
Serious Mental Illness	0.5283		0.4096		0.5010	
Moderate Mental Illness	0.4095		0.0402		0.3806	
Mild Mental Illness	0.2688		0.8974		0.5217	
No Mental Illness	0.4509		0.6760		0.0735	
NO DEPENDENCE OR ABUSE	0.8326		0.0681		0.3697	
Any Mental Illness	0.8950		0.0674		0.3626	
Serious Mental Illness	0.8129		0.8011		0.9416	
Moderate Mental Illness	0.9097		0.2100		0.1759	
Mild Mental Illness	0.6508		0.0532		0.4077	
No Mental Illness	0.8489		0.4654		0.7222	

*Low precision; no estimate reported.

NOTE: Estimates in this table are based only on responses to suicide and mental illness items in the Mental Health module. Respondents with unknown suicide information were excluded.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2010 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.