110427

Table 1.61C Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
TOTAL	322	362	193	234	257	280
18	32	38	19	23	25	30
19	31	30	22	18	22	24
20	23	37	15	23	19	29
21	28	34	19	19	21	27
22	26	29	21	18	17	22
23	29	24	20	14	22	20
24	25	27	15	18	20	20
25	26	24	17	15	20	19
26-29	86	91	49	57	67	72
30-34	101	95	58	64	80	71
35-39	100	105	59	65	82	84
40-44	112	86	73	46	90	72
45-49	105	108	72	75	70	73
50-54	128	140	77	86	101	112
55-59	118	164	80	65	90	147
60-64	91	116	56	89	73	74
65 or Older	121	140	52	98	110	107

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110427

Table 1.61D Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
TOTAL	0.14	0.16	0.18	0.21	0.22	0.24
18	0.61	0.73	0.73	0.86	1.02	1.19
19	0.66	0.65	0.89	0.80	0.95	1.04
20	0.55	0.76	0.68	0.93	0.92	1.23
21	0.67	0.72	0.90	0.86	0.98	1.13
22	0.62	0.67	0.99	0.83	0.80	1.05
23	0.70	0.59	0.96	0.69	1.03	0.96
24	0.57	0.65	0.78	0.91	0.89	0.93
25	0.67	0.60	0.94	0.78	1.00	0.89
26-29	0.49	0.52	0.58	0.66	0.77	0.84
30-34	0.49	0.46	0.57	0.63	0.79	0.67
35-39	0.48	0.54	0.56	0.68	0.80	0.84
40-44	0.51	0.42	0.71	0.45	0.79	0.68
45-49	0.46	0.47	0.64	0.69	0.59	0.62
50-54	0.57	0.62	0.70	0.79	0.87	0.95
55-59	0.62	0.82	0.87	0.71	0.93	1.37
60-64	0.58	0.69	0.76	1.08	0.90	0.88
65 or Older	0.32	0.36	0.32	0.58	0.51	0.49

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110427

Table 1.61P Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Gender and Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
TOTAL	0.3973		0.3944		0.6779	
18	0.2692		0.6936		0.2295	
19	0.8433		0.3424		0.6131	
20	0.0091		0.0859		0.0541	
21	0.6029		0.2672		0.8379	
22	0.1700		0.7766		0.0241	
23	0.0667		0.0735		0.4145	
24	0.6476		0.6153		0.8580	
25	0.2191		0.7799		0.1599	
26-29	0.6744		0.3703		0.9281	
30-34	0.2882		0.4349		0.0372	
35-39	0.7114		0.8623		0.8118	
40-44	0.0294		0.0211		0.3895	
45-49	0.2433		0.5880		0.2558	
50-54	0.5742		0.7292		0.6567	
55-59	0.2176		0.8968		0.1223	
60-64	0.5157		0.2684		0.8622	
65 or Older	0.6063		0.1053		0.6328	

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110427

Table 1.62C Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Age Group, Demographic Characteristics, and Health Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic/Health Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	322	362	75	83	211	217	229	271
GENDER								
Male	193	234	49	53	132	141	134	172
Female	257	280	57	64	168	158	190	221
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	305	320	67	71	196	192	209	240
White	291	287	60	61	188	169	204	215
Black or African American	105	136	31	35	79	86	61	100
American Indian or Alaska Native	8	11	5	5	6	9	*	*
Native Hawaiian or Other Pacific Islander	14	21	*	*	*	*	*	*
Asian	35	53	16	24	30	30	*	37
Two or More Races	33	69	6	16	25	16	21	*
Hispanic or Latino	130	162	35	43	84	100	96	123
HEALTH INSURANCE[1]								
Private	232	228	52	56	155	147	157	164
Medicaid/CHIP[2]	161	203	42	40	97	122	119	152
Other[3]	176	199	19	25	80	79	154	179
No Coverage	142	161	45	48	109	102	77	110
CURRENT EMPLOYMENT								
Full-Time	195	195	45	43	150	149	114	119
Part-Time	103	115	41	45	75	86	61	61
Unemployed	103	113	33	36	74	75	63	71
Other[4]	236	267	41	47	126	124	187	230

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

[1] Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

[2] CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

[3] Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

[4] The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110427

Table 1.62D Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Age Group, Demographic Characteristics, and Health Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic/Health Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	0.14	0.16	0.22	0.24	0.21	0.22	0.24	0.28
GENDER								
Male	0.18	0.21	0.29	0.31	0.27	0.29	0.31	0.38
Female	0.22	0.24	0.34	0.38	0.33	0.32	0.38	0.43
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.16	0.16	0.24	0.26	0.24	0.23	0.24	0.27
White	0.18	0.18	0.28	0.29	0.28	0.26	0.28	0.29
Black or African American	0.38	0.48	0.61	0.68	0.63	0.66	0.65	1.02
American Indian or Alaska Native	0.79	1.07	2.01	2.42	1.44	1.87	*	*
Native Hawaiian or Other Pacific Islander	1.98	3.12	*	*	*	*	*	*
Asian	0.35	0.50	1.03	1.45	0.57	0.55	*	1.01
Two or More Races	1.32	2.46	1.44	2.72	2.24	1.49	2.26	*
Hispanic or Latino	0.42	0.51	0.57	0.65	0.49	0.59	1.20	1.48
HEALTH INSURANCE[1]								
Private	0.15	0.15	0.27	0.29	0.22	0.22	0.22	0.24
Medicaid/CHIP[2]	0.75	0.89	0.81	0.73	1.07	1.23	1.82	2.05
Other[3]	0.33	0.36	0.88	1.03	1.18	1.15	0.34	0.39
No Coverage	0.38	0.40	0.45	0.48	0.54	0.46	1.10	1.31
CURRENT EMPLOYMENT								
Full-Time	0.17	0.17	0.34	0.34	0.22	0.22	0.31	0.33
Part-Time	0.32	0.34	0.42	0.46	0.61	0.68	0.58	0.53
Unemployed	0.67	0.75	0.69	0.73	1.00	1.08	2.02	1.96
Other[4]	0.35	0.39	0.55	0.58	0.86	0.83	0.41	0.49

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

[1] Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

[2] CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

[3] Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

[4] The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110427

Table 1.62P Serious Psychological Distress in the Past Month among Persons Aged 18 or Older, by Age Group, Demographic Characteristics, and Health Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic/Health Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
TOTAL	0.3973		0.4487		0.5867		0.1364	
GENDER								
Male	0.3944		0.5528		0.9767		0.1482	
Female	0.6779		0.1352		0.4578		0.4523	
HISPANIC ORIGIN AND RACE								
Not Hispanic or Latino	0.7334		0.6387		0.2353		0.1399	
White	0.6417		0.9444		0.1229		0.4813	
Black or African American	0.1380		0.9651		0.6435		0.0977	
American Indian or Alaska Native	0.8978		0.3554		0.6418		*	
Native Hawaiian or Other Pacific Islander	0.3920		*		*		*	
Asian	0.2983		0.8213		0.7060		*	
Two or More Races	0.7245		0.0139		0.3707		*	
Hispanic or Latino	0.2007		0.4966		0.2147		0.7238	
HEALTH INSURANCE[1]								
Private	0.9722		0.7831		0.4469		0.5787	
Medicaid/CHIP[2]	0.9803		0.6327		0.9135		0.9531	
Other[3]	0.1889		0.2910		0.5939		0.1300	
No Coverage	0.8394		0.9973		0.1997		0.1660	
CURRENT EMPLOYMENT								
Full-Time	0.6977		0.6547		0.8102		0.8646	
Part-Time	0.7044		0.3995		0.9471		0.7031	
Unemployed	0.8711		0.8644		0.7430		0.8678	
Other[4]	0.1910		0.4099		0.6345		0.0800	

*Low precision; no estimate reported.

NOTE: Serious Psychological Distress (SPD) is defined for this table as having a score of 13 or higher on the K6 scale during the past 30 days.

[1] Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.
[2] CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.
[3] Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.
[4] The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.