

**Table 1.6C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010**

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
<b>TOTAL</b>	322	362	184	220	265	267
18	20	23	8	16	18	16
19	18	17	11	9	14	14
20	18	22	10	12	14	18
21	18	26	12	13	12	23
22	19	23	13	14	12	17
23	20	22	12	13	16	18
24	18	22	11	13	15	15
25	21	20	11	11	18	16
26-29	85	84	51	47	66	70
30-34	95	107	54	62	76	85
35-39	104	92	58	54	89	75
40-44	97	100	65	61	76	81
45-49	104	116	63	81	77	81
50-54	125	140	85	81	91	115
55-59	104	151	58	63	87	133
60-64	113	106	40	86	107	61
65 or Older	106	106	63	83	86	58

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.6D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2009 and 2010**

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
<b>TOTAL</b>	0.14	0.16	0.17	0.20	0.23	0.23
18	0.39	0.46	0.32	0.61	0.71	0.73
19	0.39	0.39	0.45	0.39	0.64	0.66
20	0.43	0.46	0.48	0.48	0.71	0.81
21	0.43	0.59	0.57	0.59	0.59	1.01
22	0.45	0.53	0.65	0.64	0.60	0.80
23	0.49	0.53	0.59	0.61	0.77	0.85
24	0.43	0.52	0.56	0.63	0.66	0.73
25	0.55	0.49	0.60	0.57	0.92	0.77
26-29	0.50	0.50	0.61	0.55	0.77	0.84
30-34	0.48	0.53	0.54	0.62	0.76	0.81
35-39	0.51	0.47	0.55	0.56	0.88	0.74
40-44	0.46	0.48	0.64	0.60	0.69	0.76
45-49	0.45	0.51	0.57	0.74	0.64	0.69
50-54	0.55	0.63	0.77	0.75	0.78	0.97
55-59	0.54	0.77	0.64	0.69	0.89	1.26
60-64	0.72	0.64	0.54	1.04	1.28	0.72
65 or Older	0.28	0.27	0.38	0.49	0.40	0.26

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.6P (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Age Category	Total (2009)	Total (2010)	Male (2009)	Male (2010)	Female (2009)	Female (2010)
<b>TOTAL</b>	0.0709		0.0990		0.2780	
18	0.0897		0.0115		0.7162	
19	0.5992		0.6486		0.3632	
20	0.3059		0.6771		0.3130	
21	0.0267		0.5475		0.0203	
22	0.1070		0.3157		0.1707	
23	0.6545		0.8522		0.6192	
24	0.9909		0.9287		0.9787	
25	0.3129		0.6888		0.1185	
26-29	0.3843		0.7573		0.4282	
30-34	0.3944		0.2926		0.8165	
35-39	0.5339		0.7837		0.5155	
40-44	0.4408		0.8958		0.3477	
45-49	0.0686		0.1336		0.2623	
50-54	0.3922		0.7032		0.1532	
55-59	0.2257		0.6637		0.2685	
60-64	0.8229		0.1538		0.4733	
65 or Older	0.9922		0.4551		0.4475	

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.7C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Demographic Characteristic</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26-49 (2009)</b>	<b>Aged 26-49 (2010)</b>	<b>Aged 50+ (2009)</b>	<b>Aged 50+ (2010)</b>
<b>TOTAL</b>	322	362	51	59	219	224	218	263
<b>GENDER</b>								
Male	184	220	30	35	126	143	127	158
Female	265	267	41	48	173	165	185	200
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	310	330	48	51	207	208	212	236
White	296	296	42	46	190	182	208	220
Black or African American	89	98	19	21	71	88	47	40
American Indian or Alaska Native	13	28	5	5	12	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	2	*	*	*	*
Asian	44	60	13	9	41	47	*	*
Two or More Races	40	69	6	15	26	20	30	*
Hispanic or Latino	103	126	22	29	79	78	59	96
<b>CURRENT EMPLOYMENT</b>								
Full-Time	188	212	23	32	147	156	107	131
Part-Time	93	109	27	37	81	77	38	65
Unemployed	88	99	25	24	58	69	60	69
Other <sup>1</sup>	221	228	29	30	121	121	179	193

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.7D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010**

<b>Demographic Characteristic</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26-49 (2009)</b>	<b>Aged 26-49 (2010)</b>	<b>Aged 50+ (2009)</b>	<b>Aged 50+ (2010)</b>
<b>TOTAL</b>	0.14	0.16	0.15	0.17	0.22	0.23	0.23	0.27
<b>GENDER</b>								
Male	0.17	0.20	0.18	0.20	0.26	0.29	0.29	0.35
Female	0.23	0.23	0.24	0.29	0.34	0.33	0.37	0.39
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.16	0.17	0.17	0.19	0.25	0.25	0.25	0.27
White	0.18	0.19	0.20	0.21	0.29	0.28	0.28	0.30
Black or African American	0.32	0.36	0.40	0.42	0.55	0.69	0.51	0.42
American Indian or Alaska Native	1.21	2.51	2.28	2.53	2.58	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	1.34	*	*	*	*
Asian	0.43	0.59	0.87	0.60	0.77	0.89	*	*
Two or More Races	1.58	2.47	1.26	2.47	2.36	1.88	3.08	*
Hispanic or Latino	0.33	0.39	0.36	0.44	0.46	0.46	0.75	1.15
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.16	0.18	0.19	0.25	0.22	0.23	0.29	0.36
Part-Time	0.29	0.32	0.28	0.38	0.65	0.62	0.36	0.56
Unemployed	0.57	0.65	0.54	0.51	0.79	0.98	1.88	1.89
Other <sup>1</sup>	0.33	0.33	0.38	0.38	0.83	0.82	0.40	0.42

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.7P (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:  
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Demographic Characteristic</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26-49 (2009)</b>	<b>Aged 26-49 (2010)</b>	<b>Aged 50+ (2009)</b>	<b>Aged 50+ (2010)</b>
<b>TOTAL</b>	0.0709		0.0132		0.3189		0.2378	
<b>GENDER</b>								
Male	0.0990		0.0582		0.2849		0.3317	
Female	0.2780		0.0785		0.6595		0.4365	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.0878		0.0731		0.3478		0.2089	
White	0.3553		0.0314		0.8447		0.2769	
Black or African American	0.0232		0.9606		0.0079		0.9886	
American Indian or Alaska Native	0.2420		0.8803		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*		*	
Asian	0.4149		0.1126		0.4168		*	
Two or More Races	0.9680		0.1598		0.8329		*	
Hispanic or Latino	0.4927		0.0417		0.7269		0.9893	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.2712		0.0232		0.7052		0.3874	
Part-Time	0.1864		0.0886		0.7476		0.2523	
Unemployed	0.3308		1.0000		0.1574		0.8847	
Other <sup>1</sup>	0.4544		0.4934		0.6591		0.5620	

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.8C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Geographic/Socioeconomic Characteristic</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26-49 (2009)</b>	<b>Aged 26-49 (2010)</b>	<b>Aged 50+ (2009)</b>	<b>Aged 50+ (2010)</b>
<b>TOTAL</b>	322	362	51	59	219	224	218	263
<b>GEOGRAPHIC REGION</b>								
Northeast	123	131	25	28	86	88	83	90
Midwest	133	153	23	22	104	90	82	122
South	209	213	26	35	132	133	154	145
West	151	187	31	34	102	117	114	142
<b>COUNTY TYPE</b>								
Large Metro	256	266	43	48	164	173	182	181
Small Metro	161	174	27	36	125	120	101	122
250K – 1 Mil. Pop.	129	153	22	30	104	103	80	106
< 250K Pop.	92	81	16	19	70	62	61	59
Nonmetro	130	172	17	19	88	81	90	143
Urbanized	93	117	10	9	59	61	69	95
Less Urbanized	81	95	13	15	63	52	48	78
Completely Rural	46	84	6	8	21	16	34	*
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	142	165	28	30	99	97	95	127
100-199%	140	158	26	33	101	102	93	114
200% or More	256	262	35	42	167	173	180	190
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>								
Yes	163	193	30	34	113	127	111	144
No	283	299	45	51	183	184	196	213
<b>HEALTH INSURANCE<sup>3</sup></b>								
Private	215	254	37	44	155	165	145	178
Medicaid/CHIP <sup>4</sup>	134	153	29	24	99	86	91	126
Other <sup>5</sup>	184	184	15	19	80	92	163	158
No Coverage	123	148	28	32	95	111	73	94

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>3</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>4</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>5</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.8D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: Standard Errors of Percentages, 2009 and 2010**

Geographic/Socioeconomic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
<b>TOTAL</b>	0.14	0.16	0.15	0.17	0.22	0.23	0.23	0.27
<b>GEOGRAPHIC REGION</b>								
Northeast	0.29	0.31	0.40	0.46	0.48	0.49	0.46	0.49
Midwest	0.27	0.30	0.32	0.29	0.49	0.43	0.39	0.56
South	0.25	0.25	0.22	0.28	0.36	0.37	0.45	0.41
West	0.29	0.35	0.38	0.42	0.43	0.50	0.55	0.66
<b>COUNTY TYPE</b>								
Large Metro	0.21	0.21	0.24	0.25	0.29	0.31	0.38	0.36
Small Metro	0.23	0.24	0.25	0.30	0.41	0.40	0.34	0.41
250K – 1 Mil. Pop.	0.28	0.33	0.32	0.41	0.51	0.51	0.41	0.54
< 250K Pop.	0.39	0.33	0.41	0.41	0.68	0.61	0.60	0.59
Nonmetro	0.34	0.45	0.36	0.39	0.57	0.53	0.51	0.79
Urbanized	0.54	0.72	0.48	0.43	0.88	0.89	0.91	1.29
Less Urbanized	0.46	0.52	0.52	0.64	0.84	0.73	0.61	0.90
Completely Rural	1.15	2.24	1.64	1.97	1.57	1.27	1.53	*
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	0.50	0.53	0.36	0.34	0.76	0.73	1.31	1.45
100-199%	0.32	0.33	0.32	0.38	0.53	0.48	0.53	0.62
200% or More	0.16	0.17	0.21	0.25	0.24	0.26	0.26	0.27
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>								
Yes	0.45	0.46	0.43	0.43	0.67	0.64	0.91	1.00
No	0.14	0.16	0.16	0.18	0.22	0.23	0.23	0.26
<b>HEALTH INSURANCE<sup>3</sup></b>								
Private	0.14	0.17	0.20	0.23	0.22	0.25	0.21	0.26
Medicaid/CHIP <sup>4</sup>	0.67	0.69	0.57	0.46	1.10	0.88	1.48	1.74
Other <sup>5</sup>	0.34	0.34	0.69	0.81	1.18	1.30	0.36	0.34
No Coverage	0.33	0.38	0.29	0.33	0.47	0.52	1.05	1.14

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>3</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>4</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>5</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.8P (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group, Geographic Characteristics, and Socioeconomic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic/Socioeconomic Characteristic	Total (2009)	Total (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26-49 (2009)	Aged 26-49 (2010)	Aged 50+ (2009)	Aged 50+ (2010)
<b>TOTAL</b>	0.0709		0.0132		0.3189		0.2378	
<b>GEOGRAPHIC REGION</b>								
Northeast	0.3498		0.2317		0.4630		0.7206	
Midwest	0.5832		0.2578		0.3891		0.0578	
South	0.3885		0.0009		0.7123		0.7816	
West	0.1551		0.2903		0.0734		0.9105	
<b>COUNTY TYPE</b>								
Large Metro	0.2872		0.0601		0.1062		0.7473	
Small Metro	0.0886		0.0319		0.7298		0.1289	
250K – 1 Mil. Pop.	0.0387		0.0092		0.5492		0.1039	
< 250K Pop.	0.9676		0.8960		0.7536		0.7629	
Nonmetro	0.6278		0.7102		0.2259		0.1275	
Urbanized	0.3667		0.2357		0.9904		0.2076	
Less Urbanized	0.5957		0.7853		0.1381		0.5527	
Completely Rural	0.6972		0.9848		0.2721		*	
<b>POVERTY LEVEL<sup>1</sup></b>								
Less Than 100%	0.4799		0.4840		0.2736		0.9674	
100-199%	0.7116		0.1008		0.1892		0.2535	
200% or More	0.1255		0.0485		0.2165		0.4977	
<b>GOVERNMENT ASSISTANCE<sup>2</sup></b>								
Yes	0.6205		0.3085		0.8788		0.7294	
No	0.1955		0.0428		0.5220		0.3563	
<b>HEALTH INSURANCE<sup>3</sup></b>								
Private	0.0206		0.0070		0.2978		0.0804	
Medicaid/CHIP <sup>4</sup>	0.4601		0.0725		0.2866		0.6545	
Other <sup>5</sup>	0.1716		0.1787		0.0921		0.6159	
No Coverage	0.6518		0.2205		0.8485		0.9774	

\*Low precision; no estimate reported.

NOTE: Some 2009 and 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

<sup>1</sup> Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

<sup>2</sup> Government Assistance is defined as one or more household family members having received Supplemental Security Income (SSI), cash assistance (Temporary Assistance for Needy Families, TANF), noncash assistance, or food stamps.

<sup>3</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>4</sup> CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

<sup>5</sup> Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.