

Table 2.9C Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Total ¹ (2009)	Total ¹ (2010)	MDE (2009)	MDE (2010)	No MDE (2009)	No MDE (2010)
PAST YEAR USE						
Illicit Drugs ²	89	93	37	34	97	102
Marijuana and Hashish	77	83	30	29	77	86
Cocaine	22	22	8	9	20	20
Crack	6	5	4	4	5	3
Heroin	7	8	4	4	6	7
Hallucinogens	38	40	15	16	35	37
LSD	14	17	8	6	11	16
PCP	10	8	4	3	9	7
Ecstasy	29	32	11	14	26	29
Inhalants	42	42	16	17	40	40
Nonmedical Use of Psychotherapeutics ^{3,4}	58	58	27	22	56	57
Pain Relievers	53	54	25	20	50	52
OxyContin [®]	20	24	9	7	17	21
Tranquilizers	26	30	13	12	22	27
Stimulants ⁴	23	24	13	11	20	21
Methamphetamine ⁴	13	15	9	6	9	13
Sedatives	16	14	10	7	12	11
Illicit Drugs Other Than Marijuana ²	71	74	31	29	72	74
PAST MONTH USE						
Daily Cigarette Use ⁵	29	29	11	9	28	27
Heavy Alcohol Use ⁶	30	28	12	9	28	26

*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.9D Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Standard Errors of Percentages, 2009 and 2010

Substance	Total ¹ (2009)	Total ¹ (2010)	MDE (2009)	MDE (2010)	No MDE (2009)	No MDE (2010)
PAST YEAR USE						
Illicit Drugs ²	0.36	0.38	1.46	1.48	0.37	0.40
Marijuana and Hashish	0.31	0.34	1.34	1.36	0.32	0.35
Cocaine	0.09	0.09	0.39	0.46	0.09	0.09
Crack	0.03	0.02	0.19	0.19	0.02	0.01
Heroin	0.03	0.03	0.19	0.20	0.03	0.03
Hallucinogens	0.16	0.16	0.76	0.81	0.15	0.16
LSD	0.06	0.07	0.41	0.29	0.05	0.07
PCP	0.04	0.03	0.20	0.18	0.04	0.03
Ecstasy	0.12	0.13	0.57	0.73	0.12	0.13
Inhalants	0.17	0.17	0.81	0.85	0.17	0.18
Nonmedical Use of Psychotherapeutics ^{3,4}	0.24	0.24	1.22	1.11	0.23	0.24
Pain Relievers	0.22	0.22	1.13	1.02	0.21	0.22
OxyContin [®]	0.08	0.10	0.45	0.39	0.08	0.09
Tranquilizers	0.11	0.12	0.65	0.64	0.10	0.12
Stimulants ⁴	0.09	0.10	0.68	0.56	0.09	0.09
Methamphetamine ⁴	0.05	0.06	0.43	0.31	0.04	0.06
Sedatives	0.06	0.06	0.48	0.39	0.05	0.05
Illicit Drugs Other Than Marijuana ²	0.29	0.30	1.34	1.37	0.29	0.31
PAST MONTH USE						
Daily Cigarette Use ⁵	0.12	0.12	0.54	0.46	0.12	0.12
Heavy Alcohol Use ⁶	0.12	0.11	0.61	0.45	0.12	0.12

*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.9P Substance Use in the Past Year and Past Month among Persons Aged 12 to 17, by Past Year Major Depressive Episode (MDE): P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Total¹ (2009)	Total¹ (2010)	MDE (2009)	MDE (2010)	No MDE (2009)	No MDE (2010)
PAST YEAR USE						
Illicit Drugs ²	0.8272		0.4636		0.6152	
Marijuana and Hashish	0.3793		0.5887		0.5123	
Cocaine	0.9726		0.7094		0.5978	
Crack	0.1848		0.9272		0.1685	
Heroin	0.5574		0.9848		0.5023	
Hallucinogens	0.7009		0.4284		0.9311	
LSD	0.8135		0.0479		0.6867	
PCP	0.3554		0.5333		0.4210	
Ecstasy	0.2161		0.1575		0.4589	
Inhalants	0.2524		0.5159		0.1963	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.5114		0.1617		0.8441	
Pain Relievers	0.2945		0.1208		0.6075	
OxyContin [®]	0.6674		0.1148		0.3800	
Tranquilizers	0.7702		0.7707		0.7636	
Stimulants ⁴	0.8729		0.2979		0.8069	
Methamphetamine ⁴	0.4723		0.5314		0.2108	
Sedatives	0.4986		0.5488		0.6838	
Illicit Drugs Other Than Marijuana ²	0.6104		0.5429		0.4633	
PAST MONTH USE						
Daily Cigarette Use ⁵	0.2642		0.4873		0.3144	
Heavy Alcohol Use ⁶	0.0122		0.0233		0.0482	

*Low precision; no estimate reported.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms.

¹ Estimates in the Total column represent persons aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings Report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.