

**Table 1.15A Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Numbers in Thousands, 2011 and 2012**

<b>Dependence or Abuse Status</b>	<b>Any Mental Illness (2011)</b>	<b>Any Mental Illness (2012)</b>	<b>Serious Mental Illness (2011)</b>	<b>Serious Mental Illness (2012)</b>
<b>DEPENDENCE</b>				
None	36,733	37,855	7,397	7,552
Both Illicit Drugs and Alcohol	695	835	348	368
Illicit Drugs or Alcohol	4,649 <sup>b</sup>	5,858	1,613 <sup>a</sup>	2,018
Illicit Drugs	2,029 <sup>a</sup>	2,473	809	915
Alcohol	3,314 <sup>b</sup>	4,220	1,152	1,471
<b>DEPENDENCE OR ABUSE</b>				
None	34,568	35,305	6,888	6,960
Both Illicit Drugs and Alcohol	1,167	1,480	476	562
Illicit Drugs or Alcohol	6,813 <sup>b</sup>	8,409	2,123 <sup>a</sup>	2,609
Illicit Drugs	2,492 <sup>b</sup>	3,131	951	1,108
Alcohol	5,489 <sup>b</sup>	6,757	1,649 <sup>a</sup>	2,063

\*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.15B Any Mental Illness and Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Dependence or Abuse Status: Percentages, 2011 and 2012**

<b>Dependence or Abuse Status</b>	<b>Any Mental Illness (2011)</b>	<b>Any Mental Illness (2012)</b>	<b>Serious Mental Illness (2011)</b>	<b>Serious Mental Illness (2012)</b>
<b>DEPENDENCE</b>				
None	16.5	16.9	3.3	3.4
Both Illicit Drugs and Alcohol	66.7	70.8	33.4	31.3
Illicit Drugs or Alcohol	45.2 <sup>a</sup>	50.1	15.7	17.3
Illicit Drugs	51.6	54.3	20.6	20.1
Alcohol	44.8 <sup>a</sup>	50.7	15.6	17.7
<b>DEPENDENCE OR ABUSE</b>				
None	16.2	16.5	3.2	3.2
Both Illicit Drugs and Alcohol	52.8	59.0	21.5	22.4
Illicit Drugs or Alcohol	36.1 <sup>b</sup>	40.7	11.2	12.6
Illicit Drugs	46.4	49.6	17.7	17.5
Alcohol	34.9 <sup>b</sup>	40.1	10.5	12.2

\*Low precision; no estimate reported.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.16A Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2011 and 2012**

<b>Dependence or Abuse Status</b>	<b>Total (2011)</b>	<b>Total (2012)</b>	<b>Any Mental Illness (2011)</b>	<b>Any Mental Illness (2012)</b>	<b>Serious Mental Illness (2011)</b>	<b>Serious Mental Illness (2012)</b>	<b>Moderate Mental Illness (2011)</b>	<b>Moderate Mental Illness (2012)</b>	<b>Mild Mental Illness (2011)</b>	<b>Mild Mental Illness (2012)</b>	<b>No Mental Illness (2011)</b>	<b>No Mental Illness (2012)</b>
<b>DEPENDENCE</b>												
Both Illicit Drugs and Alcohol	1,042	1,178	695	835	348	368	153	249	194	218	347	344
Illicit Drugs or Alcohol	10,290 <sup>b</sup>	11,695	4,649 <sup>b</sup>	5,858	1,613 <sup>a</sup>	2,018	1,087 <sup>b</sup>	1,679	1,948	2,161	5,641	5,836
Illicit Drugs	3,934 <sup>a</sup>	4,557	2,029 <sup>a</sup>	2,473	809	915	489 <sup>a</sup>	689	731	869	1,905	2,084
Alcohol	7,397 <sup>a</sup>	8,316	3,314 <sup>b</sup>	4,220	1,152	1,471	752 <sup>b</sup>	1,239	1,411	1,510	4,083	4,096
<b>DEPENDENCE OR ABUSE</b>												
Both Illicit Drugs and Alcohol	2,212	2,506	1,167	1,480	476	562	291	412	401	506	1,044	1,026
Illicit Drugs or Alcohol	18,887 <sup>b</sup>	20,669	6,813 <sup>b</sup>	8,409	2,123 <sup>a</sup>	2,609	1,605 <sup>b</sup>	2,258	3,085	3,542	12,073	12,260
Illicit Drugs	5,374 <sup>b</sup>	6,316	2,492 <sup>b</sup>	3,131	951	1,108	598 <sup>a</sup>	824	943	1,200	2,882	3,185
Alcohol	15,724 <sup>a</sup>	16,859	5,489 <sup>b</sup>	6,757	1,649 <sup>a</sup>	2,063	1,297 <sup>b</sup>	1,847	2,543	2,848	10,235	10,101

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.16B Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2011 and 2012**

Dependence or Abuse Status	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>DEPENDENCE</b>												
Both Illicit Drugs and Alcohol	0.4	0.5	1.7	1.9	3.9	3.8	1.5	2.1	0.9	1.0	0.2	0.2
Illicit Drugs or Alcohol	4.4 <sup>b</sup>	5.0	11.2 <sup>b</sup>	13.4	17.9	21.1	10.5 <sup>b</sup>	14.1	8.8	9.7	2.9	3.0
Illicit Drugs	1.7 <sup>a</sup>	1.9	4.9	5.7	9.0	9.6	4.7	5.8	3.3	3.9	1.0	1.1
Alcohol	3.2 <sup>a</sup>	3.5	8.0 <sup>a</sup>	9.7	12.8	15.4	7.3 <sup>b</sup>	10.4	6.4	6.8	2.1	2.1
<b>DEPENDENCE OR ABUSE</b>												
Both Illicit Drugs and Alcohol	1.0	1.1	2.8	3.4	5.3	5.9	2.8	3.5	1.8	2.3	0.5	0.5
Illicit Drugs or Alcohol	8.1 <sup>b</sup>	8.8	16.5 <sup>b</sup>	19.2	23.6	27.3	15.5 <sup>a</sup>	18.9	14.0	15.9	6.3	6.4
Illicit Drugs	2.3 <sup>b</sup>	2.7	6.0 <sup>a</sup>	7.2	10.5	11.6	5.8	6.9	4.3	5.4	1.5	1.7
Alcohol	6.8	7.2	13.3 <sup>b</sup>	15.5	18.3	21.6	12.6 <sup>a</sup>	15.5	11.5	12.8	5.4	5.3

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.17A Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	5,374 <sup>b</sup>	6,316	2,492 <sup>b</sup>	3,131	951	1,108	598 <sup>a</sup>	824	943	1,200	2,882	3,185
18-25	2,574	2,706	970	1,110	306	283	238	303	426	523	1,604	1,597
26-49	2,235 <sup>b</sup>	2,924	1,207 <sup>a</sup>	1,676	525	660	294	450	388	566	1,028	1,247
50 or Older	565	686	315	345	119	165	67	71	129	110	250	341
<b>MALE</b>	3,615 <sup>a</sup>	4,270	1,336 <sup>b</sup>	1,819	423	581	343	452	571 <sup>a</sup>	787	2,278	2,451
18-25	1,726	1,731	490	585	124	123	114	158	252	304	1,236	1,146
26-49	1,509 <sup>a</sup>	2,000	676 <sup>a</sup>	1,014	249	348	178	258	249	408	833	986
50 or Older	380	539	170	220	*	*	*	36	69	74	210	319
<b>FEMALE</b>	1,760	2,046	1,156	1,312	528	528	256	372	372	413	604	734
18-25	848	976	480	525	182	160	124	145	173	219	368	451
26-49	726	924	531	662	276	312	116	192	139	158	195	262
50 or Older	185	147	145	125	70	55	*	34	60	35	40	22

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.17B Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	2.3 <sup>b</sup>	2.7	6.0 <sup>a</sup>	7.2	10.5	11.6	5.8	6.9	4.3	5.4	1.5	1.7
18-25	7.5	7.8	15.3	16.4	23.6	19.9	14.3	16.6	12.5	14.8	5.7	5.7
26-49	2.3 <sup>b</sup>	3.0	6.1 <sup>a</sup>	8.1	10.6	13.0	5.6	8.0	4.0	5.6	1.3	1.6
50 or Older	0.6	0.7	2.1	2.1	4.3	5.4	1.9	1.6	1.5	1.3	0.3	0.4
<b>MALE</b>	3.2 <sup>a</sup>	3.8	8.4 <sup>a</sup>	10.8	13.1	16.0	9.4	9.6	6.3 <sup>a</sup>	9.2	2.4	2.5
18-25	10.0	10.0	19.7	22.1	27.8	23.8	19.8	23.6	17.2	20.8	8.4	7.8
26-49	3.1 <sup>a</sup>	4.2	9.0 <sup>a</sup>	12.3	14.7	17.6	9.7	11.4	6.3	10.2	2.1	2.5
50 or Older	0.8	1.1	2.9	3.7	*	*	*	2.0	1.9	2.4	0.5	0.8
<b>FEMALE</b>	1.5	1.7	4.5	4.9	9.1	8.9	3.8	5.1	2.9	3.0	0.6	0.8
18-25	5.0	5.7	12.4	12.7	21.4	17.7	11.4	12.5	9.0	10.6	2.8 <sup>a</sup>	3.4
26-49	1.5	1.9	4.3	5.3	8.5	10.1	3.4	5.7	2.4	2.6	0.5	0.7
50 or Older	0.3	0.3	1.6	1.2	4.1	2.8	*	1.3	1.1	0.6	0.1	0.0

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.18A Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	15,724 <sup>a</sup>	16,859	5,489 <sup>b</sup>	6,757	1,649 <sup>a</sup>	2,063	1,297 <sup>b</sup>	1,847	2,543	2,848	10,235	10,101
18-25	4,936	4,956	1,577	1,777	370	446	438	501	768	829	3,359	3,179
26-49	7,867	8,441	2,962 <sup>b</sup>	3,800	909	1,154	686 <sup>b</sup>	1,082	1,367	1,565	4,905	4,641
50 or Older	2,921	3,462	950	1,180	370	463	173	264	407	453	1,971	2,282
<b>MALE</b>	10,040 <sup>a</sup>	11,198	2,800 <sup>b</sup>	3,779	754 <sup>a</sup>	1,140	632 <sup>b</sup>	1,055	1,415	1,585	7,239	7,419
18-25	2,950	2,905	752	834	151	175	181	252	420	407	2,198	2,070
26-49	5,156	5,634	1,503 <sup>b</sup>	2,168	420 <sup>a</sup>	642	303 <sup>b</sup>	614	780	913	3,653	3,466
50 or Older	1,933 <sup>a</sup>	2,660	545	777	*	*	*	189	215	265	1,388	1,883
<b>FEMALE</b>	5,685	5,661	2,688	2,978	895	924	666	792	1,128	1,263	2,996	2,682
18-25	1,986	2,051	825	942	220	271	257	249	348	422	1,161	1,109
26-49	2,711	2,807	1,459	1,632	488	512	383	468	587	652	1,252	1,175
50 or Older	988	802	405	404	187	141	25	74	192	188	583	399

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.18B Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	6.8	7.2	13.3 <sup>b</sup>	15.5	18.3	21.6	12.6 <sup>a</sup>	15.5	11.5	12.8	5.4	5.3
18-25	14.4	14.3	24.8	26.2	28.6	31.3	26.3	27.5	22.6	23.5	12.0	11.4
26-49	8.0	8.6	14.9 <sup>b</sup>	18.3	18.4	22.8	13.1 <sup>b</sup>	19.3	14.0	15.5	6.3	6.0
50 or Older	2.9	3.4	6.3	7.3	13.4	15.0	5.0	5.9	4.6	5.3	2.3	2.6
<b>MALE</b>	9.0 <sup>a</sup>	9.9	17.6 <sup>b</sup>	22.4	23.4 <sup>a</sup>	31.5	17.3	22.4	15.7	18.6	7.5	7.7
18-25	17.2	16.7	30.3	31.5	33.8	33.9	31.4	37.7	28.7	27.9	15.0	14.1
26-49	10.7	11.7	20.1 <sup>b</sup>	26.3	24.9	32.5	16.6 <sup>a</sup>	27.1	19.7	22.8	9.0	8.7
50 or Older	4.1 <sup>a</sup>	5.6	9.2	13.0	*	*	*	10.7	5.9	8.6	3.4	4.5
<b>FEMALE</b>	4.7	4.6	10.5	11.1	15.5	15.5	9.9	11.0	8.7	9.2	3.1	2.8
18-25	11.6	11.9	21.3	22.8	25.8	29.9	23.6	21.6	18.0	20.4	8.8	8.5
26-49	5.4	5.6	11.7	13.0	15.0	16.6	11.3	14.0	10.1	10.7	3.3	3.1
50 or Older	1.8	1.5	4.4	4.0	11.1	7.2	1.1	2.7	3.6	3.4	1.3	0.9

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.19A Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Numbers in Thousands, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	18,887 <sup>b</sup>	20,669	6,813 <sup>b</sup>	8,409	2,123 <sup>a</sup>	2,609	1,605 <sup>b</sup>	2,258	3,085	3,542	12,073	12,260
18-25	6,369	6,535	2,060 <sup>a</sup>	2,338	529	567	546	657	985	1,114	4,308	4,197
26-49	9,217 <sup>a</sup>	10,245	3,629 <sup>b</sup>	4,687	1,153 <sup>a</sup>	1,489	854 <sup>b</sup>	1,309	1,622	1,889	5,588	5,559
50 or Older	3,300	3,888	1,124	1,384	441	553	205	292	478	539	2,177	2,504
<b>MALE</b>	12,105 <sup>b</sup>	13,751	3,459 <sup>b</sup>	4,676	932 <sup>b</sup>	1,378	804 <sup>b</sup>	1,257	1,723	2,042	8,646	9,075
18-25	3,879	3,930	984	1,134	207	232	228 <sup>a</sup>	323	549	579	2,894	2,796
26-49	6,062 <sup>a</sup>	6,860	1,863 <sup>b</sup>	2,665	516 <sup>a</sup>	787	412 <sup>b</sup>	731	935	1,147	4,199	4,195
50 or Older	2,165 <sup>a</sup>	2,961	612	877	*	*	*	202	239	316	1,553	2,084
<b>FEMALE</b>	6,781	6,918	3,354	3,733	1,191	1,232	801	1,001	1,362	1,500	3,427	3,185
18-25	2,490	2,605	1,076	1,204	322	335	318	334	436	535	1,414	1,401
26-49	3,156	3,385	1,766	2,021	637	702	442	578	688	741	1,390	1,364
50 or Older	1,135	928	512	507	233	194	41	89	238	224	623	420

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.19B Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Percentages, 2011 and 2012**

Gender/ Age Group	Total (2011)	Total (2012)	Any Mental Illness (2011)	Any Mental Illness (2012)	Serious Mental Illness (2011)	Serious Mental Illness (2012)	Moderate Mental Illness (2011)	Moderate Mental Illness (2012)	Mild Mental Illness (2011)	Mild Mental Illness (2012)	No Mental Illness (2011)	No Mental Illness (2012)
<b>TOTAL</b>	8.1 <sup>b</sup>	8.8	16.5 <sup>b</sup>	19.2	23.6	27.3	15.5 <sup>a</sup>	18.9	14.0	15.9	6.3	6.4
18-25	18.6	18.9	32.4	34.5	40.8	39.9	32.9	36.0	29.0	31.6	15.4	15.1
26-49	9.4 <sup>a</sup>	10.5	18.2 <sup>b</sup>	22.6	23.3 <sup>a</sup>	29.4	16.3 <sup>b</sup>	23.3	16.6	18.7	7.2	7.2
50 or Older	3.3	3.8	7.4	8.6	16.0	18.0	5.9	6.5	5.4	6.3	2.6	2.9
<b>MALE</b>	10.8 <sup>b</sup>	12.2	21.8 <sup>b</sup>	27.7	29.0 <sup>a</sup>	38.1	22.1	26.7	19.1 <sup>a</sup>	23.9	9.0	9.4
18-25	22.6	22.7	39.6	42.9	46.6	44.8	39.7	48.4	37.5	39.7	19.7	19.0
26-49	12.6 <sup>a</sup>	14.3	24.9 <sup>b</sup>	32.4	30.5	39.8	22.5 <sup>a</sup>	32.4	23.6	28.7	10.3	10.5
50 or Older	4.6 <sup>a</sup>	6.2	10.3	14.7	*	*	*	11.4	6.6	10.2	3.8	5.0
<b>FEMALE</b>	5.6	5.7	13.2	13.9	20.6	20.7	12.0	13.9	10.5	11.0	3.6	3.3
18-25	14.5	15.1	27.8	29.2	37.8	37.0	29.3	28.9	22.6	25.9	10.7	10.7
26-49	6.3	6.8	14.2	16.1	19.5	22.7	13.0	17.3	11.9	12.1	3.7	3.7
50 or Older	2.1	1.7	5.6	5.0	13.8	9.9	1.8	3.3	4.5	4.1	1.4	0.9

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories.

Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.