

**Table 1.33A Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Numbers in Thousands, 2011 and 2012**

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental Illness	Mental Health Treatment or Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment or Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment Only <sup>1</sup>	Mental Health Treatment Only <sup>1</sup>	Substance Use Treatment at a Specialty Facility Only <sup>2</sup>	Substance Use Treatment at a Specialty Facility Only <sup>2</sup>	Mental Health Treatment and Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment and Substance Use Treatment at a Specialty Facility <sup>1,2</sup>
	(2011)	(2012)	(2011)	(2012)	(2011)	(2012)	(2011)	(2012)
<b>TOTAL</b>	32,709 <sup>b</sup>	35,270	30,530 <sup>b</sup>	32,931	1,046	1,124	1,112	1,191
<b>DEPENDENCE OR ABUSE</b>	5,107 <sup>a</sup>	5,812	3,876	4,302	582	692	647	809
Any Mental Illness	3,162 <sup>b</sup>	3,882	2,365 <sup>a</sup>	2,849	248	360	550	668
Serious Mental Illness	1,469	1,680	1,078	1,142	86	120	304	412
Moderate Mental Illness	689 <sup>b</sup>	1,066	553 <sup>b</sup>	833	48	84	88	148
Mild Mental Illness	1,004	1,136	733	874	114	155	158	108
No Mental Illness	1,945	1,930	1,511	1,452	334	332	97	141
<b>NO DEPENDENCE OR ABUSE</b>	27,602 <sup>a</sup>	29,458	26,654 <sup>a</sup>	28,629	464	432	465	382
Any Mental Illness	14,037	14,451	13,633	14,066	105	93	299	290
Serious Mental Illness	4,469	4,487	4,341	4,366	14	38	115	83
Moderate Mental Illness	4,106	4,481	3,995	4,392	22	6	89	81
Mild Mental Illness	5,462	5,483	5,298	5,308	70	49	94	127
No Mental Illness	13,565 <sup>a</sup>	15,007	13,021 <sup>a</sup>	14,563	359	340	166	91

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

<sup>2</sup> Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

**Table 1.33B Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Percentages, 2011 and 2012**

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental Illness	Mental Health Treatment or Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment or Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment Only <sup>1</sup>		Substance Use Treatment at a Specialty Facility Only <sup>2</sup>		Mental Health Treatment and Substance Use Treatment at a Specialty Facility <sup>1,2</sup>	Mental Health Treatment and Substance Use Treatment at a Specialty Facility <sup>1,2</sup>
	(2011)	(2012)	(2011)	(2012)	(2011)	(2012)	(2011)	(2012)
<b>TOTAL</b>	14.1 <sup>a</sup>	15.0	13.2 <sup>a</sup>	14.0	0.4	0.5	0.5	0.5
<b>DEPENDENCE OR ABUSE</b>	27.1	28.2	20.6	20.9	3.1	3.3	3.4	3.9
Any Mental Illness	46.5	46.3	34.8	34.0	3.6	4.3	8.1	7.9
Serious Mental Illness	69.3	64.7	50.9	44.0	4.1	4.6	14.3	15.8
Moderate Mental Illness	43.1	47.2	34.6	36.9	3.0	3.7	5.5	6.6
Mild Mental Illness	32.6	32.2	23.8	24.8	3.7	4.4	5.1	3.0
No Mental Illness	16.2	15.8	12.6	11.9	2.8	2.7	0.8	1.2
<b>NO DEPENDENCE OR ABUSE</b>	12.9 <sup>a</sup>	13.8	12.5 <sup>a</sup>	13.4	0.2	0.2	0.2	0.2
Any Mental Illness	40.7	41.0	39.5	39.9	0.3	0.3	0.9	0.8
Serious Mental Illness	65.0	64.5	63.1	62.8	0.2	0.5	1.7	1.2
Moderate Mental Illness	47.1	46.5	45.8	45.6	0.2	0.1	1.0	0.8
Mild Mental Illness	28.9	29.4	28.0	28.5	0.4	0.3	0.5	0.7
No Mental Illness	7.6 <sup>a</sup>	8.4	7.3 <sup>b</sup>	8.1	0.2	0.2	0.1	0.1

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Mental illness estimates in 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>a</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded.

<sup>2</sup> Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.