

Table 1.51A Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Type of Illicit Drug Use in the Past Year: Numbers in Thousands, 2011 and 2012

Drug	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
ILLCIT DRUGS¹	2,925 ^a	3,475	950	1,164	448	573
Marijuana and Hashish	2,087 ^b	2,701	659 ^a	919	309 ^a	471
Cocaine	467	657	190	225	74	136
Crack	97 ^a	197	53	55	25	63
Heroin	106	*	36	*	24	17
Hallucinogens	464	528	189	215	101	136
LSD	109	146	47	50	20	34
PCP	*	*	*	*	*	*
Ecstasy	297	343	131	128	61	83
Inhalants	130	182	61	75	17	45
Nonmedical Use of Psychotherapeutics ^{2,3}	1,587	1,916	535	726	306	326
Pain Relievers	1,266	1,444	419	559	217	231
OxyContin [®]	270	281	107	131	46	45
Tranquilizers	709	785	236	316	144	134
Stimulants ³	354 ^a	538	125 ^a	260	78	119
Methamphetamine ³	173	292	79	156	64	81
Sedatives	*	105	33	30	28	29
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,872 ^a	2,243	648	808	325	398

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.51B Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Type of Illicit Drug Use in the Past Year: Percentages, 2011 and 2012

Drug	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
ILLCIT DRUGS¹	8.8	9.4	2.8	3.2	1.3	1.6
Marijuana and Hashish	8.0 ^a	9.6	2.5 ^a	3.3	1.2 ^a	1.7
Cocaine	13.0	14.7	5.3	5.0	2.1	3.1
Crack	16.5	22.2	9.2	6.2	4.2	7.1
Heroin	18.9	*	6.4	*	4.3	2.8
Hallucinogens	13.7	14.2	5.6	5.8	3.0	3.7
LSD	15.2	16.2	6.6	5.5	2.7	3.7
PCP	*	*	*	*	*	*
Ecstasy	15.0	14.8	6.6	5.5	3.1	3.6
Inhalants	12.7	17.4	5.9	7.2	1.6	4.3
Nonmedical Use of Psychotherapeutics ^{2,3}	12.4	12.8	4.2	4.9	2.4	2.2
Pain Relievers	13.2	13.0	4.4	5.0	2.3	2.1
OxyContin [®]	19.4	22.0	7.7	10.3	3.3	3.5
Tranquilizers	15.5	14.0	5.1	5.6	3.2	2.4
Stimulants ³	14.8	18.1	5.2	8.7	3.3	4.0
Methamphetamine ³	18.7	27.4	8.5	14.6	6.9	7.6
Sedatives	*	20.9	7.4	5.9	6.3	5.7
ILLCIT DRUGS OTHER THAN MARIJUANA¹	11.5	11.9	4.0	4.3	2.0	2.1

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52A Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Tobacco Product and Alcohol Use in the Past Year and Past Month: Numbers in Thousands, 2011 and 2012

Substance	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
PAST YEAR						
Tobacco Products ¹	4,258	4,614	1,420	1,483	688	775
Cigarettes	3,766	4,127	1,318	1,349	658	689
Smokeless Tobacco	525	567	171	220	74	76
Cigars	1,390	1,546	488	613	223	299
Alcohol	6,377	6,819	1,803	1,932	826	996
PAST MONTH						
Tobacco Products ¹	3,738	4,032	1,267	1,334	600	702
Cigarettes	3,283	3,585	1,178	1,205	574	617
Smokeless Tobacco	297	350	97	115	46	42
Cigars	796	871	308	379	151	175
Pipe Tobacco	224	251	78	110	50	65
Alcohol	4,924	5,254	1,346	1,383	635	758
Binge Alcohol Use ²	2,783	3,136	772	853	407 ^a	563
Heavy Alcohol Use ²	964	1,093	275	319	141	167

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.

Table 1.52B Had Serious Thoughts of Suicide, Made Any Suicide Plans, or Attempted Suicide in the Past Year among Persons Aged 18 or Older, by Tobacco Product and Alcohol Use in the Past Year and Past Month: Percentages, 2011 and 2012

Substance	Had Serious Thoughts of Suicide (2011)	Had Serious Thoughts of Suicide (2012)	Made Any Suicide Plans (2011)	Made Any Suicide Plans (2012)	Attempted Suicide (2011)	Attempted Suicide (2012)
PAST YEAR						
Tobacco Products ¹	5.5	5.9	1.8	1.9	0.9	1.0
Cigarettes	5.9	6.4	2.1	2.1	1.0	1.1
Smokeless Tobacco	5.0	5.1	1.6	2.0	0.7	0.7
Cigars	6.1	6.5	2.1	2.6	1.0	1.3
Alcohol	3.9	4.1	1.1	1.2	0.5	0.6
PAST MONTH						
Tobacco Products ¹	5.7	6.0	1.9	2.0	0.9	1.0
Cigarettes	6.0	6.4	2.2	2.2	1.1	1.1
Smokeless Tobacco	3.9	4.1	1.3	1.4	0.6	0.5
Cigars	6.7	6.8	2.6	3.0	1.3	1.4
Pipe Tobacco	11.7	10.7	4.1	4.7	2.6	2.8
Alcohol	3.8	4.0	1.0	1.0	0.5	0.6
Binge Alcohol Use ²	4.9	5.4	1.4	1.5	0.7	1.0
Heavy Alcohol Use ²	6.2	6.6	1.8	1.9	0.9	1.0

*Low precision; no estimate reported.

NOTE: Respondents with unknown suicide information were excluded.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 and 2012.