

Table 1.69A Any Mental Illness in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2008-2012

Demographic Characteristic	2008¹	2009	2010	2011	2012
TOTAL	39,826 ^b	41,195 ^a	41,417 ^a	41,381 ^a	43,714
AGE					
18-25	6,099 ^b	6,043 ^b	6,151 ^b	6,356 ^a	6,774
26-49	20,697	21,455	20,648	19,936	20,777
50 or Older	13,030 ^b	13,698 ^b	14,618 ^a	15,089	16,163
GENDER					
Male	14,778 ^b	15,614 ^a	16,461	15,895	16,856
Female	25,048 ^a	25,581	24,956 ^a	25,486	26,858
HISPANIC ORIGIN AND RACE					
Not Hispanic or Latino	34,971 ^b	36,321	36,580	36,793	38,049
White	28,286	29,799	29,704	29,552	30,066
Black or African American	4,314	4,113 ^a	4,520	4,364	5,044
American Indian or Alaska Native	163 ^b	219 ^a	187 ^a	357	366
Native Hawaiian or Other Pacific Islander	*	*	*	*	*
Asian	1,300	1,287	1,425	1,473	1,608
Two or More Races	666	757	584	847	709
Hispanic or Latino	4,855 ^a	4,874 ^a	4,837 ^a	4,589 ^b	5,665

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Any Mental Illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Any mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Respondents' AMI status for the 2008 NSDUH is determined based on responses to questions on distress (K6 scale) and impairment (World Health Organization Disability Assessment Schedule [WHODAS] for half the sample and Sheehan Disability Scale [SDS] for the other half). Respondents' AMI status for the 2009-2012 NSDUH are based solely on responses to the WHODAS questions because the SDS questions were dropped. For details, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.69B Any Mental Illness in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2008-2012

Demographic Characteristic	2008 ¹	2009	2010	2011	2012
TOTAL	17.7 ^a	18.1	18.1	17.8	18.6
AGE					
18-25	18.5 ^a	18.0 ^b	18.1 ^b	18.5 ^a	19.6
26-49	20.7	21.6	20.9	20.3	21.2
50 or Older	14.1 ^a	14.5	15.1	15.0	15.8
GENDER					
Male	13.6 ^a	14.2	14.8	14.2	14.9
Female	21.5	21.8	21.1	21.1	22.0
HISPANIC ORIGIN AND RACE					
Not Hispanic or Latino	18.0 ^a	18.5	18.5	18.5	19.0
White	18.3	19.2	19.0	19.0	19.3
Black or African American	16.7	15.6 ^a	16.9	16.3	18.6
American Indian or Alaska Native	17.3 ^a	20.0	17.4 ^a	27.7	28.3
Native Hawaiian or Other Pacific Islander	*	*	*	*	*
Asian	13.3	12.8	13.7	13.1	13.9
Two or More Races	26.5	31.1 ^b	22.0	25.8	20.7
Hispanic or Latino	16.0	15.7	15.2	13.5 ^b	16.3

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Any Mental Illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Any mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Respondents' AMI status for the 2008 NSDUH is determined based on responses to questions on distress (K6 scale) and impairment (World Health Organization Disability Assessment Schedule [WHODAS] for half the sample and Sheehan Disability Scale [SDS] for the other half). Respondents' AMI status for the 2009-2012 NSDUH are based solely on responses to the WHODAS questions because the SDS questions were dropped. For details, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.70A Any Mental Illness in the Past Year among Persons Aged 18 or Older, by Geographic and Socioeconomic Characteristics: Numbers in Thousands, 2008-2012

Geographic/Socioeconomic Characteristic	2008 ¹	2009	2010	2011	2012
TOTAL	39,826 ^b	41,195 ^a	41,417 ^a	41,381 ^a	43,714
GEOGRAPHIC REGION					
Northeast	7,865	7,587	7,890	6,966 ^a	7,723
Midwest	9,075	9,169	9,369	9,020	9,173
South	13,905 ^b	14,604 ^b	14,656 ^a	15,362	16,282
West	8,980 ^b	9,835	9,502	10,033	10,536
COUNTY TYPE					
Large Metro	20,591 ^a	21,396	21,449	20,781 ^a	22,619
Small Metro	12,548 ^a	12,941	13,127	13,668	13,943
250K – 1 Mil. Pop.	8,231	8,623	8,874	9,045	8,899
< 250K Pop.	4,317	4,317	4,252	4,623	5,044
Nonmetro	6,687	6,858	6,842	6,932	7,151
Urbanized	2,765	2,890	2,963	2,986	3,138
Less Urbanized	3,276	3,378	3,305	3,318	3,343
Completely Rural	647	591	574	628	670
POVERTY LEVEL²					
Less Than 100%	6,402 ^b	6,955 ^b	7,323 ^b	8,428 ^b	9,945
100-199%	8,925	8,885	9,756	9,813	9,890
200% or More	24,313	25,148	24,187	23,005	23,669
EDUCATION					
< High School	7,137	6,582 ^a	6,475 ^a	6,567	7,491
High School Graduate	11,838	12,255	12,519	11,838 ^a	13,037
Some College	11,071 ^a	11,796	11,462	11,877	12,307
College Graduate	9,780 ^a	10,563	10,960	11,099	10,880
HEALTH INSURANCE³					
Private	23,775	24,949	23,693	22,565	23,589
Medicaid/CHIP ⁴	5,287 ^b	5,767 ^b	6,084 ^a	6,664	7,085
Other ⁵	8,227 ^b	8,587 ^b	9,815 ^a	9,727 ^b	11,458
No Coverage	7,751	7,446 ^a	7,863	8,095	8,423

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Any Mental Illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Any mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Respondents' AMI status for the 2008 NSDUH is determined based on responses to questions on distress (K6 scale) and impairment (World Health Organization Disability Assessment Schedule [WHODAS] for half the sample and Sheehan Disability Scale [SDS] for the other half). Respondents' AMI status for the 2009-2012 NSDUH are based solely on responses to the WHODAS questions because the SDS questions were dropped. For details, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁵ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.70B Any Mental Illness in the Past Year among Persons Aged 18 or Older, by Geographic and Socioeconomic Characteristics: Percentages, 2008-2012

Geographic/Socioeconomic Characteristic	2008 ¹	2009	2010	2011	2012
TOTAL	17.7 ^a	18.1	18.1	17.8	18.6
GEOGRAPHIC REGION					
Northeast	18.9	18.0	18.7	16.3	18.0
Midwest	18.4	18.4	18.7	18.0	18.2
South	17.0 ^a	17.6	17.4	17.9	18.7
West	17.3 ^a	18.8	17.9	18.6	19.3
COUNTY TYPE					
Large Metro	17.2	17.7	17.5	16.8 ^a	18.0
Small Metro	18.4	18.7	18.8	18.8	19.6
250K – 1 Mil. Pop.	18.2	18.9	19.1	18.4	18.6
< 250K Pop.	18.8 ^a	18.3 ^a	18.1 ^b	19.6	21.6
Nonmetro	18.0	18.4	18.7	19.1	18.7
Urbanized	18.0	17.7	19.1	19.8	18.8
Less Urbanized	18.2	19.6	18.7	19.1	19.3
Completely Rural	17.7	15.6	16.9	16.3	15.4
POVERTY LEVEL²					
Less Than 100%	25.2	25.1	24.7	25.8	26.8
100-199%	21.6	20.2	20.5	20.5	21.8
200% or More	15.5	16.3	16.0	15.2	15.6
EDUCATION					
< High School	20.3	18.9 ^b	18.5 ^b	19.9	21.9
High School Graduate	17.0 ^a	17.6	18.1	17.0 ^a	18.7
Some College	19.3	20.4	19.4	19.3	19.7
College Graduate	15.6	16.3	16.7	16.3	15.9
HEALTH INSURANCE³					
Private	15.3	16.2	15.7	14.9	15.4
Medicaid/CHIP ⁴	28.7	29.9	28.8	28.7	30.5
Other ⁵	15.9 ^b	16.3 ^b	18.1	17.4 ^a	19.5
No Coverage	22.8	20.5	20.4 ^a	21.3	22.3

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Any Mental Illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Any mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Respondents' AMI status for the 2008 NSDUH is determined based on responses to questions on distress (K6 scale) and impairment (World Health Organization Disability Assessment Schedule [WHODAS] for half the sample and Sheehan Disability Scale [SDS] for the other half). Respondents' AMI status for the 2009-2012 NSDUH are based solely on responses to the WHODAS questions because the SDS questions were dropped. For details, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁵ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.71A Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2008-2012

Demographic Characteristic	2008	2009	2010	2011	2012
TOTAL	8,331 ^b	8,383 ^a	9,334	9,011	9,570
AGE					
18-25	1,236 ^a	1,124 ^b	1,334	1,296	1,423
26-49	4,763	4,858	5,132	4,951	5,065
50 or Older	2,332 ^a	2,401 ^a	2,868	2,764	3,082
GENDER					
Male	2,868 ^b	2,847 ^b	3,351	3,218	3,619
Female	5,463	5,535	5,983	5,793	5,951
HISPANIC ORIGIN AND RACE					
Not Hispanic or Latino	7,553	7,486	8,304	8,163	8,038
White	6,515	6,341	6,743	6,899	6,608
Black or African American	657 ^a	719	1,028	743	927
American Indian or Alaska Native	28 ^a	53	87	126	111
Native Hawaiian or Other Pacific Islander	9	*	*	*	17
Asian	220	164	232	214	233
Two or More Races	124	183	196	151	144
Hispanic or Latino	778 ^b	897 ^b	1,030 ^a	848 ^b	1,531

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment. Serious mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.71B Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2008-2012

Demographic Characteristic	2008	2009	2010	2011	2012
TOTAL	3.7	3.7	4.1	3.9	4.1
AGE					
18-25	3.8	3.3 ^b	3.9	3.8	4.1
26-49	4.8	4.9	5.2	5.0	5.2
50 or Older	2.5	2.5	3.0	2.8	3.0
GENDER					
Male	2.6 ^a	2.6 ^a	3.0	2.9	3.2
Female	4.7	4.7	5.1	4.8	4.9
HISPANIC ORIGIN AND RACE					
Not Hispanic or Latino	3.9	3.8	4.2	4.1	4.0
White	4.2	4.1	4.3	4.4	4.2
Black or African American	2.5	2.7	3.9	2.8	3.4
American Indian or Alaska Native	3.0 ^a	4.8	8.1	9.8	8.5
Native Hawaiian or Other Pacific Islander	1.2	*	*	*	1.8
Asian	2.2	1.6	2.2	1.9	2.0
Two or More Races	4.9	7.5	7.4	4.6	4.2
Hispanic or Latino	2.6 ^b	2.9 ^b	3.2	2.5 ^b	4.4

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.72A Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Geographic and Socioeconomic Characteristics: Numbers in Thousands, 2008-2012

Geographic/Socioeconomic Characteristic	2008	2009	2010	2011	2012
TOTAL	8,331 ^b	8,383 ^a	9,334	9,011	9,570
GEOGRAPHIC REGION					
Northeast	1,619	1,476	1,651	1,436	1,569
Midwest	1,968	2,151	2,276	2,109	1,983
South	2,785 ^b	2,827 ^a	3,113	3,108	3,572
West	1,959 ^a	1,928 ^a	2,293	2,357	2,445
COUNTY TYPE					
Large Metro	4,301	4,237	4,700	4,153	4,826
Small Metro	2,692	2,551 ^a	2,972	3,268	3,190
250K – 1 Mil. Pop.	1,771	1,663	2,091	2,092	2,004
< 250K Pop.	921	888	880	1,176	1,186
Nonmetro	1,338	1,594	1,662	1,590	1,553
Urbanized	631	664	758	686	602
Less Urbanized	620	741	700	765	760
Completely Rural	86 ^a	189	203	138	191
POVERTY LEVEL¹					
Less Than 100%	1,486 ^b	1,792 ^b	2,075 ^a	2,481	2,685
100-199%	2,046	1,993	2,242	2,199	2,338
200% or More	4,764	4,551	4,987	4,293	4,503
EDUCATION					
< High School	1,171 ^a	1,363	1,198 ^a	1,552	1,640
High School Graduate	2,726	2,623	3,251	2,399 ^b	3,089
Some College	2,635	2,441	2,698	2,883	2,744
College Graduate	1,799	1,956	2,186	2,177	2,096
HEALTH INSURANCE²					
Private	4,502	4,099	4,765	3,967	4,106
Medicaid/CHIP ³	1,438 ^b	1,680	1,694	2,033	1,965
Other ⁴	1,706 ^b	1,799 ^b	2,204	1,941 ^a	2,569
No Coverage	1,678 ^b	1,710 ^b	1,909	1,972	2,290

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment. Serious mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁴ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.

Table 1.72B Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Geographic and Socioeconomic Characteristics: Percentages, 2008-2012

Geographic/Socioeconomic Characteristic	2008	2009	2010	2011	2012
TOTAL	3.7	3.7	4.1	3.9	4.1
GEOGRAPHIC REGION					
Northeast	3.9	3.5	3.9	3.4	3.7
Midwest	4.0	4.3	4.6	4.2	3.9
South	3.4 ^a	3.4	3.7	3.6	4.1
West	3.8	3.7	4.3	4.4	4.5
COUNTY TYPE					
Large Metro	3.6	3.5	3.8	3.4	3.8
Small Metro	4.0	3.7 ^a	4.2	4.5	4.5
250K – 1 Mil. Pop.	3.9	3.7	4.5	4.3	4.2
< 250K Pop.	4.0	3.8	3.7	5.0	5.1
Nonmetro	3.6	4.3	4.5	4.4	4.1
Urbanized	4.1	4.1	4.9	4.5	3.6
Less Urbanized	3.4	4.3	4.0	4.4	4.4
Completely Rural	2.4	5.0	6.0	3.6	4.4
POVERTY LEVEL¹					
Less Than 100%	5.9 ^a	6.5	7.0	7.6	7.2
100-199%	4.9	4.5	4.7	4.6	5.2
200% or More	3.0	2.9	3.3	2.8	3.0
EDUCATION					
< High School	3.3 ^b	3.9	3.4 ^b	4.7	4.8
High School Graduate	3.9	3.8	4.7	3.4 ^b	4.4
Some College	4.6	4.2	4.6	4.7	4.4
College Graduate	2.9	3.0	3.3	3.2	3.1
HEALTH INSURANCE²					
Private	2.9	2.7	3.2 ^a	2.6	2.7
Medicaid/CHIP ³	7.8	8.7	8.0	8.7	8.5
Other ⁴	3.3 ^a	3.4	4.1	3.5	4.4
No Coverage	4.9 ^a	4.7 ^a	4.9 ^a	5.2	6.1

*Low precision; no estimate reported.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Serious mental illness estimates from 2008 to 2011 may differ from previously published estimates due to revised estimation procedures. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

^a Difference between estimate and 2012 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds. Respondents aged 18 to 22 who were living in a college dormitory were excluded.

² Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

³ CHIP is the Children's Health Insurance Program. Individuals aged 19 or younger are eligible for this plan.

⁴ Other Health Insurance is defined as having Medicare, CHAMPUS, TRICARE, CHAMPVA, the VA, military health care, or any other type of health insurance.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2012.